

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07D 311/58, 335/06, 215/54, 311/92, 493/04, 409/04, 491/04, A61K 31/35</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/47890 (43) International Publication Date: 29 October 1998 (29.10.98)</p>
<p>(21) International Application Number: PCT/US98/07677 (22) International Filing Date: 18 April 1998 (18.04.98) (30) Priority Data: 60/044,485 21 April 1997 (21.04.97) US (71) Applicant (for all designated States except US): G.D. SEARLE &amp; CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CARTER, Jeffery, S. [US/US]; 15321 Grantley Drive, Chesterfield, MO 63017 (US). OBUKOWICZ, Mark, G. [US/US]; 655 N. Kirkwood Road, Kirkwood, MO 63122 (US). DEVADAS, Balekudru [IN/US]; 2175 Parasol Drive, Chesterfield, MO 63017 (US). TALLEY, John, J. [US/US]; 8772 Pine Avenue, Brentwood, MO 63144 (US). BROWN, David, L. [US/US]; 15504 Twingate, Chesterfield, MO 63017 (US). GRANETO, Matthew, J. [US/US]; 352 Hartwell Court, Chesterfield, MO 63017 (US). BERTENSHAW, Stephen, R. [US/US]; 43 Lansdown Lane, Cheshire, CT 06410 (US). ROGIER, Donald, J., Jr. [US/US]; 1828 Westmeade Drive, Chesterfield, MO 63017 (US). NAGARAJAN, Srinivasan, R. [IN/US]; 16209 Forest Meadows Drive, Chesterfield, MO 63005</p>		<p>(US). HANAU, Cathleen, E. [US/US]; 1441 Rankin Drive, St. Louis, MO 63117 (US). HARTMANN, Susan, J. [US/US]; 449 Colony Woods Drive, Kirkwood, MO 63122 (US). LUDWIG, Cindy, L. [US/US]; 1412 Dautel Lane, St. Louis, MO 63146 (US). METZ, Suzanne [US/US]; 525 Westernmill Drive, Chesterfield, MO 63017 (US). (74) Agents: BULOCK, Joseph, W. et al.; G.D. Searle &amp; Co., P.O. Box 5110, Chicago, IL 60680-5110 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: SUBSTITUTED BENZOPYRAN DERIVATIVES FOR THE TREATMENT OF INFLAMMATION</p> <div data-bbox="615 1165 1104 1365" data-label="Chemical-Block"> <p style="text-align: right;">( I ' )</p> </div> <p>(57) Abstract</p> <p>A class of benzopyran derivatives is described for use in treating cyclooxygenase-2 mediated disorders. Compounds of particular interest are defined by Formula (I'), wherein X, A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>, A<sup>4</sup>, R, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are as described in the specification.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**SUBSTITUTED BENZOPYRAN DERIVATIVES  
FOR THE TREATMENT OF INFLAMMATION**

5

**FIELD OF THE INVENTION**

This invention is in the field of anti-inflammatory pharmaceutical agents and specifically relates to compounds, compositions and methods for treating cyclooxygenase-2 mediated disorders, such as inflammation and inflammation-related disorders.

**BACKGROUND OF THE INVENTION**

15 Prostaglandins play a major role in the inflammation process and the inhibition of prostaglandin production, especially production of  $\text{PGG}_2$ ,  $\text{PGH}_2$  and  $\text{PGE}_2$ , has been a common target of antiinflammatory drug discovery. However, common non-steroidal antiinflammatory drugs (NSAIDs) that are active in reducing the prostaglandin-induced pain and swelling associated with the inflammation process are also active in affecting other prostaglandin-regulated processes not associated with the inflammation process. Thus, use of high doses of most common NSAIDs can produce severe side effects, including life threatening ulcers, that limit their therapeutic potential. An alternative to NSAIDs is the use of corticosteroids, which have even more drastic side effects, especially when long term therapy is involved.

Previous NSAIDs have been found to prevent the production of prostaglandins by inhibiting enzymes in the human arachidonic acid/prostaglandin pathway, including the enzyme cyclooxygenase (COX). The recent



discovery of an inducible enzyme associated with inflammation (named "cyclooxygenase-2 (COX-2)" or "prostaglandin G/H synthase II") provides a viable target of inhibition which more effectively reduces inflammation and produces fewer and less drastic side effects.

The references below that disclose antiinflammatory activity, show continuing efforts to find a safe and effective antiinflammatory agent. The novel benzopyran, dihydroquinoline, benzothiopyran and dihydronaphthalene derivatives disclosed herein are such safe and also effective antiinflammatory agents furthering such efforts. The substituted benzopyran, dihydroquinoline, benzothiopyran and dihydronaphthalene derivatives disclosed herein preferably selectively inhibit cyclooxygenase-2 over cyclooxygenase-1.

US Patent No. 5,618,843, to Fisher et al., generically describes acid substituted bicyclic moieties as IIb/IIIA antagonists. WO 94/13659, published June 23, 1994, describes fused benzo compounds for the treatment of CNS disorders. Manrao et al. (J. Indian. Counc. Chem., 12, 38-41 (1996)) describes carboxy coumarinimide derivatives and their antifungal activity. US Patent No. 5,348,976, to Shibata et al., describes amide substituted benzopyrans as antifungals.

WO96/40110, published December 19, 1996, describes benzopyran derivatives as tyrosine kinase modulators. Loiodice et al. (Tetrahedron, 6, 1001-11 (1995)) describe the preparation of 6-chloro-2,3-dihydro-4H-1-benzopyran carboxylic acids.

Clemence et al. (J. Med. Chem., 31, 1453-62, (1988)) describe 4-hydroxy-3-quinolinecarboxylic acids as starting material in the preparation of antiinflammatories. Lazer, et al. (J. Med. Chem., 40, 980-89 (1997)) describe


benzothiopyran carboxylates as starting material in the preparation of antiinflammatories.

Benzopyran-3-carboxylic acids have been described. Gupta et al. (Indian J. Chem., 21B, 344-347 (1982)) describe chromene-3-carboxylic acid as an intermediate in the preparation of centrally acting muscle relaxants. Rene and Royer (Eur. J. Med. Chem. - Chim. Ther., 10, 72-78 (1975)) describe the preparation of chromene-3-carboxylic acid. US Patent No. 4,665,202, to Rimbault et al., describes 2-phenyl substituted flavenes and thioflavenes as 5-lipoxygenase inhibitors. U.S. Patent No. 5,250,547, to Lochead et al., describe benzopyran derivatives as 5-lipoxygenase inhibitors. Satoh et al. [J. Med. Chem., 36, 3580-94 (1993)] describe substituted chromenes as 5-lipoxygenase inhibitors. U.S. Patent No. 5,155,130, to Stanton et al. describes substituted chromenes as 5-lipoxygenase inhibitors, and specifically 6-benzyloxy-2H-benzopyran-3-carboxylic acid as an intermediate.

However, compounds of the current invention have not been described as cyclooxygenase inhibitors.

#### DESCRIPTION OF THE INVENTION

A class of compounds useful in treating cyclooxygenase-mediated disorders is defined by Formula I':



wherein X is selected from O, S, CR<sup>a</sup>R<sup>b</sup> and NR<sup>a</sup>;

- wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, (optionally substituted phenyl)-C<sub>1</sub>-C<sub>3</sub>-alkyl, alkylsulfonyl, phenylsulfonyl, benzylsulfonyl, acyl and carboxy-C<sub>1</sub>-C<sub>6</sub>-alkyl;
- wherein each of R<sup>b</sup> and R<sup>c</sup> is independently selected from
- 5 hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-alkyl;
- oe wherein CR<sup>a</sup>R<sup>b</sup> form a cyclopropyl ring;
- wherein R is selected from carboxyl, aminocarbonyl, C<sub>1</sub>-
- 10 C<sub>6</sub>-alkylsulfonylaminocarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl;
- wherein R<sup>''</sup> is selected from hydrido, phenyl, thienyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl and C<sub>2</sub>-C<sub>6</sub>-alkenyl;
- wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-
- 15 alkyl;
- wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>1</sub>-C<sub>3</sub>-alkyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-
- 20 C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, -O(CF<sub>2</sub>)<sub>2</sub>O-, aryloxy, arylthio, arylsulfinyl, heteroaryloxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, aryl-C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-
- 25 haloalkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-(haloalkyl-C<sub>1</sub>-C<sub>3</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxyimino-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, heteroarylamino, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, cyano, amino, aminosulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, arylaminosulfonyl,
- 30 heteroarylaminosulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted aryl, optionally substituted heteroaryl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,
- 35 heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-


alkoxycarbonyl, formyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylcarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; and

wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least two of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon;

or wherein R<sup>2</sup> together with ring A forms a radical selected from naphthyl, quinolyl, isoquinolyl, quinoliziny, quinoxaliny and dibenzofuryl;

or an isomer or pharmaceutically acceptable salt thereof.

A related class of compounds useful in treating cyclooxygenase-2 mediated disorders is defined by Formula I:



wherein X is selected from O or S or NR<sup>a</sup>;

wherein R<sup>a</sup> is alkyl;

wherein R is selected from carboxyl, aminocarbonyl, alkylsulfonylamino, carbonyl and alkoxy;

wherein R<sup>1</sup> is selected from haloalkyl, alkyl, aralkyl, cycloalkyl and aryl optionally substituted with one or more radicals selected from alkylthio, nitro and alkylsulfonyl; and

wherein R<sup>2</sup> is one or more radicals selected from hydrido, halo, alkyl, aralkyl, alkoxy, aryloxy, heteroaryloxy, aralkyloxy, heteroaralkyloxy, haloalkyl, haloalkoxy, alkylamino, arylamino, aralkylamino, heteroarylamine, heteroarylalkylamine, nitro, amino, aminosulfonyl, alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aralkylaminosulfonyl,

heteroaralkylaminosulfonyl, heterocyclosulfonyl, alkylsulfonyl, optionally substituted aryl, optionally substituted heteroaryl, aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, and alkylcarbonyl;

5 or wherein R<sup>2</sup> together with ring A forms a naphthyl radical;

or an isomer or pharmaceutically acceptable salt thereof.

Compounds of the present invention would be useful for,  
10 but not limited to, the treatment of inflammation in a subject, and for treatment of other cyclooxygenase-2 mediated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, compounds of the  
15 invention would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis. Such compounds of the invention would be useful in the treatment of asthma,  
20 bronchitis, menstrual cramps, preterm labor, tendinitis, bursitis, liver disease including hepatitis, skin-related conditions such as psoriasis, eczema, burns and dermatitis, and from post-operative inflammation including from ophthalmic surgery such as cataract surgery and refractive  
25 surgery. Compounds of the invention also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis. Compounds of the invention would be useful in treating inflammation in such diseases as  
30 migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, scleroderma, rheumatic fever, type I diabetes, neuromuscular junction disease including myasthenia gravis, white matter disease including multiple sclerosis, sarcoidosis, nephrotic syndrome,  
35 Behcet's syndrome, polymyositis, gingivitis, nephritis,

hypersensitivity, swelling occurring after injury including brain edema, myocardial ischemia, and the like. The compounds would also be useful in the treatment of ophthalmic diseases, such as retinitis, conjunctivitis, 5 retinopathies, uveitis, ocular photophobia, and of acute injury to the eye tissue. The compounds would also be useful in the treatment of pulmonary inflammation, such as that associated with viral infections and cystic fibrosis. The compounds would also be useful for the treatment of 10 certain central nervous system disorders, such as cortical dementias including Alzheimer's disease, and central nervous system damage resulting from stroke, ischemia and trauma. The compounds of the invention are useful as anti-inflammatory agents, such as for the treatment of arthritis, 15 with the additional benefit of having significantly less harmful side effects. These compounds would also be useful in the treatment of allergic rhinitis, respiratory distress syndrome, endotoxin shock syndrome, and liver disease. The compounds would also be useful in the treatment of pain, but 20 not limited to postoperative pain, dental pain, muscular pain, and pain resulting from cancer. The compounds would be useful for the treatment of dementias. The term "treatment" includes partial or total inhibition of the dementia, including Alzheimer's disease, vascular dementia, multi- 25 infarct dementia, pre-senile dementia, alcoholic dementia, and senile dementia.

The method above would be useful for, but not limited to, treating and preventing inflammation-related cardiovascular disorders in a subject. The method would be 30 useful for treatment and prevention of vascular diseases, coronary artery disease, aneurysm, arteriosclerosis, atherosclerosis including cardiac transplant atherosclerosis, myocardial infarction, embolism, stroke, thrombosis, including venous thrombosis, angina including 35 unstable angina, coronary plaque inflammation, bacterial-

induced inflammation including *Chlamydia*-induced inflammation, viral induced inflammation, and inflammation associated with surgical procedures such as vascular grafting including coronary artery bypass surgery, 5 revascularization procedures including angioplasty, stent placement, endarterectomy, or other invasive procedures involving arteries, veins and capillaries.

The compounds would be useful for, but not limited to, the treatment of angiogenesis-related disorders in a 10 subject. According to the present invention, the compounds are administered to a subject in need of angiogenesis inhibition. The method would be useful for treatment of neoplasia, including metastasis; ophthalmological conditions such as corneal graft rejection, ocular neovascularization, 15 retinal neovascularization including neovascularization following injury or infection, diabetic retinopathy, macular degeneration, retrolental fibroplasia and neovascular glaucoma; ulcerative diseases such as gastric ulcer; pathological, but non-malignant, conditions such as 20 hemangiomas, including infantile hemangiomas, angiofibroma of the nasopharynx and avascular necrosis of bone; and disorders of the female reproductive system such as endometriosis.

Compounds of the invention would be useful for the 25 prevention or treatment of neoplasia including cancer, such as colorectal cancer, brain cancer, bone cancer, epithelial cell-derived neoplasia (epithelial carcinoma) such as basal cell carcinoma, adenocarcinoma, gastrointestinal cancer such as lip cancer, mouth cancer, esophageal cancer, small bowel 30 cancer and stomach cancer, colon cancer, liver cancer, bladder cancer, pancreas cancer, ovary cancer, cervical cancer, lung cancer, breast cancer and skin cancer, such as squamous cell and basal cell cancers, prostate cancer, renal cell carcinoma, and other known cancers that effect 35 epithelial cells throughout the body. Preferably, neoplasia

is selected from gastrointestinal cancer, liver cancer, bladder cancer, pancreas cancer, ovary cancer, prostate cancer, cervical cancer, lung cancer, breast cancer and skin cancer, such as squamous cell and basal cell cancers. The compounds can also be used to treat the fibrosis which occurs with radiation therapy. The method can be used to treat subjects having adenomatous polyps, including those with familial adenomatous polyposis (FAP). Additionally, the method can be used to prevent polyps from forming in patients at risk of FAP.

The administration of compounds of the present invention may be used alone or in conjunction with additional therapies known to those skilled in the art in the prevention or treatment of neoplasia. Alternatively, the compounds described herein may be used in conjunctive therapy. By way of example, the compounds may be administered alone or in conjunction with other antineoplastic agents or other growth inhibiting agents or other drugs or nutrients.

There are large numbers of antineoplastic agents available in commercial use, in clinical evaluation and in pre-clinical development, which could be selected for treatment of neoplasia by combination drug chemotherapy. Such antineoplastic agents fall into several major categories, namely, antibiotic-type agents, alkylating agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents and a category of miscellaneous agents. Alternatively, other antineoplastic agents, such as metallomatrix proteases (MMP), SOD mimics or  $\alpha, \beta$ , inhibitors may be used.

A first family of antineoplastic agents which may be used in combination with compounds of the present invention consists of antimetabolite-type antineoplastic agents. Suitable antimetabolite antineoplastic agents may be selected from the group consisting of 5-FU-fibrinogen,



acanthifolic acid, aminothiadiaazole, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, 5 dideoxyguanosine, didox, Yoshitomi DMDC, doxifluridine, Wellcome EHNA, Merck & Co. EX-015, fazarabine, floxuridine, fludarabine phosphate, 5-fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku FO-152, isopropyl pyrrolizine, Lilly LY-188011, Lilly LY-264618, methobenzaprim, 10 methotrexate, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, thioguanine, tiazaofurin, Erbamont TIF, trimetrexate, tyrosine kinase inhibitors, 15 tyrosine protein kinase inhibitors, Taiho UFT and uricytin.

A second family of antineoplastic agents which may be used in combination with compounds of the present invention consists of alkylating-type antineoplastic agents. Suitable alkylating-type antineoplastic agents may be selected from 20 the group consisting of Shionogi 254-S, aldo-phosphamide analogues, altretamine, anaxirone, Boehringer Mannheim BBR-2207, bestrabucil, budotitane, Wakunaga CA-102, carboplatin, carmustine, Chinoin-139, Chinoin-153, chlorambucil, cisplatin, cyclophosphamide, American Cyanamid CL-286558, 25 Sanofi CY-233, cyplatate, Degussa D-19-384, Sumimoto DACHP(Myrr)2, diphenylspiromustine, diplatinum cytostatic, Erba distamycin derivatives, Chugai DWA-2114R, ITI E09, elmustine, Erbamont FCE-24517, estramustine phosphate sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230, 30 hepsul-fam, ifosfamide, iproplatin, lomustine, mafosfamide, mitolactol, Nippon Kayaku NK-121, NCI NSC-264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU, prednimustine, Proter PTT-119, ranimustine, semustine, SmithKline SK&F-101772, Yakult Honsha SN-22, spiromustine, Tanabe Seiyaku TA-077,

taumomustine, temozolomide, teroxirone, tetraplatin and trimelamol.

A third family of antineoplastic agents which may be used in combination with compounds of the present invention consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from the group consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II, Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azino-mycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067, Bristol-Myers BMY-25551, Bristol-Myers BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatin-1, Taiho C-1027, calicheomycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko DC92-B, ditrisarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, esperamicin-A1, esperamicin-Alb, Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-900482, glidobactin, gregatin-A, grincamycin, herbimycin, idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, mitoxantrone, SmithKline M-TAG, neoenactin, Nippon Kayaku NK-313, Nippon Kayaku NKT-01, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindamycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomicin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A, sparsomycin, SS Pharmaceutical SS-21020, SS Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin,

thrazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-10028A, Fujisawa WF-3405, Yoshitomi Y-25024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination with compounds of the present invention consists of a miscellaneous family of antineoplastic agents selected from the group consisting of alpha-carotene, alpha-difluoromethyl-arginine, acitretin, Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphetinile, amsacrine, Angiostat, ankinomycin, anti-neoplaston A10, antineoplaston A2, antineoplaston A3, antineoplaston A5, antineoplaston AS2-1, Henkel APD, aphidicolin glycinate, asparaginase, Avarol, baccharin, batracylin, benfluron, benzotript, Ipsen-Beaufour BIM-23015, bisantrene, Bristo-Myers BMY-40481, Vestar boron-10, bromofosfamide, Wellcome BW-502, Wellcome BW-773, caracemide, carmethizole hydrochloride, Ajinomoto CDAF, chlorsulfaquinoxalone, Chemes CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-958, clanfenur, claviridenone, ICN compound 1259, ICN compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol, curaderm, cytochalasin B, cytarabine, cytocytin, Merz D-609, DABIS maleate, dacarbazine, datelliptinium, didemnin-B, dihaematoporphyrin ether, dihydrolenperone, dinaline, distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi Seiyaku DN-9693, elliprabin, elliptinium acetate, Tsumura EPMTc, ergotamine, etoposide, etretinate, fenretinide, Fujisawa FR-57704, gallium nitrate, genkwadaphnin, Chugai GLA-43, Glaxo GR-63178, grifolan NMF-5N, hexadecylphosphocholine, Green Cross HO-221, homoharringtonine, hydroxyurea, BTG ICRF-187, ilmofosine, isoglutamine, isotretinoin, Otsuka JI-36, Ramot K-477, Otsuka K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110, American Cyanamid L-623, leukoregulin, lonidamine, Lundbeck LU-23-112, Lilly LY-186641, NCI (US) MAP, marycin, Merrel Dow MDL-27048, Medco MEDR-340, merbarone, merocyanine

derivatives, methylanilinoacridine, Molecular Genetics MGI-136, minactivin, mitonafide, mitoquidone, mopidamol, motretinide, Zenyaku Kogyo MST-16, N-(retinoyl)amino acids, Nisshin Flour Milling N-021, N-acylated-dehydroalanines, 5 nafazatrom, Taisho NCU-190, nocodazole derivative, Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782, NCI NSC-95580, octreotide, Ono ONO-112, oquizanocine, Akzo Org-10172, pancratistatin, pazelliptine, Warner-Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, 10 Pierre Fabre PE-1001, ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane, procarbazine, proglumide, Invitron protease nexin I, Tobishi RA-700, razoxane, Sapporo Breweries RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-15 49532, Rhone-Poulenc RP-56976, SmithKline SK&F-104864, Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane derivatives, spirogermanium, Unimed, SS Pharmaceutical SS-554, strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Toyama T-20 506, Toyama T-680, taxol, Teijin TEI-0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, Topostin, Teijin TT-82, Kyowa Hakko UCN-01, Kyowa Hakko UCN-1028, ukrain, Eastman Kodak USB-006, vinblastine sulfate, vincristine, vindesine, vinestramide, vinorelbine, 25 vintriptol, vinzolidine, withanolides and Yamanouchi YM-534.

Examples of radioprotective agents which may be used in combination with compounds of the present invention are AD-5, adchnon, amifostine analogues, detox, dimesna, 1-102, MM-159, N-acylated-dehydroalanines, TGF- Genentech, tiprotimod, 30 amifostine, WR-151327, FUT-187, ketoprofen transdermal, nabumetone, superoxide dismutase (Chiron) and superoxide dismutase Enzon.

Besides being useful for human treatment, these compounds are also useful for veterinary treatment of 35 companion animals, exotic animals and farm animals,

including mammals, rodents, and the like. More preferred animals include horses, dogs, and cats.

The present compounds may also be used in co-therapies, partially or completely, in place of  
5 other conventional antiinflammatories, such as together with steroids, NSAIDs, iNOS inhibitors, 5-lipoxygenase inhibitors, LTB<sub>4</sub> receptor antagonists and LTA<sub>4</sub> hydrolase inhibitors.

Suitable LTA<sub>4</sub> hydrolase inhibitors include RP-  
10 64966, (S,S)-3-amino-4-(4-benzyloxyphenyl)-2-hydroxybutyric acid benzyl ester (Scripps Res. Inst.), N-(2(R)-(cyclohexylmethyl)-3-(hydroxycarbamoyl)propionyl)-L-alanine (Searle), 7-(4-(4-ureidobenzyl)phenyl)heptanoic acid (Rhône-Poulenc Rorer), and  
15 3-(3-(1E,3E-tetradecadienyl)-2-oxiranyl)benzoic acid lithium salt (Searle).

Suitable LTB<sub>4</sub> receptor antagonists include, among others, ebselen, linazolast, ontazolast, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo  
20 Denmark compound ETH-615, Merck compound MAFP, Terumo compound TMK-688, Tanabe compound T-0757, Lilly compounds LY-213024, LY-210073, LY223982, LY233469, and LY255283, LY-293111, 264086 and 292728, ONO compounds ONO-LB457, ONO-4057, and ONO-  
25 LB-448, Shionogi compound S-2474, calcitrol, Lilly compounds Searle compounds SC-53228, SC-41930, SC-50605 and SC-51146, Warner Lambert compound BPC 15, SmithKline Beecham compound SB-209247 and SK&F compound SKF-104493. Preferably, the LTB<sub>4</sub> receptor  
30 antagonists are selected from calcitrol, ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo Denmark compound ETH-615, Lilly compound LY-293111, Ono compound ONO-4057, and Terumo compound TMK-688.

Suitable 5-LO inhibitors include, among others, Abbott compounds A-76745, 78773 and ABT761, Bayer Bay-x-1005, Cytomed CMI-392, Eisai E-3040, Scotia Pharmaceutica EF-40, Fujirebio F-1322, Merckle ML-3000, Purdue Frederick PF-5901, 3M Pharmaceuticals R-840, rilopirox, flobufen, linasolast, lonaprolene, masoprocil, ontasolast, tenidap, zileuton, pranlukast, tepoxalin, rilopirox, flezelastine hydrochloride, enazadrem phosphate, and bunaprolast.

10       The present compounds may also be used in combination therapies with opioids and other analgesics, including narcotic analgesics, Mu receptor antagonists, Kappa receptor antagonists, non-narcotic (i.e. non-addictive) analgesics, monoamine uptake  
15       inhibitors, adenosine regulating agents, cannabinoid derivatives, Substance P antagonists, neurokinin-1 receptor antagonists and sodium channel blockers, among others. More preferred would be combinations with compounds selected from morphine, meperidine, codeine,  
20       pentazocine, buprenorphine, butorphanol, dezocine, meptazinol, hydrocodone, oxycodone, methadone, Tramadol [(+) enantiomer], DuP 747, Dynorphine A, Enadoline, RP-60180, HN-11608, E-2078, ICI-204448, acetaminophen (paracetamol), propoxyphene, nalbuphine, E-4018,  
25       filenadol, mirfentanil, amitriptyline, DuP631, Tramadol [(-) enantiomer], GP-531, acadesine, AKI-1, AKI-2, GP-1683, GP-3269, 4030W92, tramadol racemate, Dynorphine A, E-2078, AXC3742, SNX-111, ADL2-1294, ICI-204448, CT-3, CP-99,994, and CP-99,994.

30       The compounds can be used in combination with one or more antihistamines, decongestants, diuretics, antitussive agents or with other agents previously known to be effective in combination with antiinflammatory agents.

The term "prevention" includes either preventing the onset of clinically evident cardiovascular disorders altogether or preventing the onset of a preclinically evident stage of cardiovascular disorder in individuals.

5 This includes prophylactic treatment of those at risk of developing a cardiovascular disorder.

The phrase "therapeutically-effective" is intended to qualify the amount of each agent which will achieve the goal of improvement in disorder severity and the frequency of

10 incidence over treatment of each agent by itself, while avoiding adverse side effects typically associated with alternative therapies.

The present invention preferably includes compounds which selectively inhibit cyclooxygenase-2

15 over cyclooxygenase-1. Preferably, the compounds have a cyclooxygenase-2  $IC_{50}$  of less than about 0.5  $\mu M$ , and also have a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100. Even

20 more preferably, the compounds have a cyclooxygenase-1  $IC_{50}$  of greater than about 5  $\mu M$ . Such preferred selectivity may indicate an ability to reduce the incidence of common NSAID-induced side effects.

A preferred class of compounds consists of those

25 compounds of Formula I wherein X is oxygen or sulfur; wherein R is selected from carboxyl, lower alkyl, lower aralkyl and lower alkoxy carbonyl; wherein  $R^1$  is selected from lower haloalkyl, lower cycloalkyl and phenyl; and wherein  $R^2$  is one or more radicals selected from hydrido, halo, lower

30 alkyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, lower alkylamino, nitro, amino, aminosulfonyl, lower alkylaminosulfonyl, 5- or 6- membered heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl, 5- or 6- membered nitrogen containing heterocyclosulfonyl,

35 lower alkylsulfonyl, optionally substituted phenyl, lower

aralkylcarbonyl, and lower alkylcarbonyl; or wherein R<sup>2</sup> together with ring A forms a naphthyl radical; or an isomer or pharmaceutically acceptable salt thereof.

A more preferred class of compounds consists of those  
5 compounds of Formula I wherein X is oxygen or sulfur;  
wherein R is selected from carboxyl; wherein R<sup>1</sup> is selected  
from lower haloalkyl; and wherein R<sup>2</sup> is one or more radicals  
selected from hydrido, halo, lower alkyl, lower haloalkyl,  
lower haloalkoxy, lower alkylamino, amino, aminosulfonyl,  
10 lower alkylaminosulfonyl, 5- or 6- membered  
heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl,  
lower alkylsulfonyl, 6- membered nitrogen containing  
heterocyclosulfonyl, optionally substituted phenyl, lower  
aralkylcarbonyl, and lower alkylcarbonyl; or wherein R<sup>2</sup>  
15 together with ring A forms a naphthyl radical; or an isomer  
or pharmaceutically acceptable salt thereof.

An even more preferred class of compounds consists of  
those compounds of Formula I wherein R is carboxyl; wherein  
R<sup>1</sup> is selected from fluoromethyl, chloromethyl,  
20 dichloromethyl, trichloromethyl, pentafluoroethyl,  
heptafluoropropyl, difluoroethyl, difluoropropyl,  
dichloroethyl, dichloropropyl, difluoromethyl, and  
trifluoromethyl; and wherein R<sup>2</sup> is one or more radicals  
selected from hydrido, chloro, fluoro, bromo, iodo, methyl,  
25 ethyl, isopropyl, tert-butyl, butyl, isobutyl, pentyl,  
hexyl, methoxy, ethoxy, isopropoxy, tertbutoxy,  
trifluoromethyl, difluoromethyl, trifluoromethoxy, amino,  
N,N-dimethylamino, N,N-diethylamino, N-  
phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-  
30 furylmethyl)aminosulfonyl, nitro, N,N-dimethylaminosulfonyl,  
aminosulfonyl, N-methylaminosulfonyl, N-ethylsulfonyl, 2,2-  
dimethylethylaminosulfonyl, N,N-dimethylaminosulfonyl, N-(2-  
methylpropyl)aminosulfonyl, N-morpholinosulfonyl,  
methylsulfonyl, benzylcarbonyl, 2,2-dimethylpropylcarbonyl,  
35 phenylacetyl and phenyl; or wherein R<sup>2</sup> together with ring A



forms a naphthyl radical; or an isomer or pharmaceutically acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula I wherein R is carboxyl; wherein  
5 R<sup>1</sup> is trifluoromethyl or pentafluorethyl; and wherein R<sup>2</sup> is selected from one or more radicals hydrido, chloro, fluoro, bromo, iodo, methyl, ethyl, isopropyl, tert-butyl, methoxy, trifluoromethyl, trifluoromethoxy, N-phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-furylmethyl)aminosulfonyl, N,N-dimethylaminosulfonyl, N-methylaminosulfonyl, N-(2,2-dimethylethyl)aminosulfonyl, dimethylaminosulfonyl, 2-methylpropylaminosulfonyl, N-morpholinosulfonyl, methylsulfonyl, benzylcarbonyl, and phenyl; or wherein R<sup>2</sup> together with ring A forms a naphthyl  
10 radical; or an isomer or pharmaceutically acceptable salt thereof.

A preferred class of compounds consists of those compounds of Formula I' wherein X is selected from O, S, CR<sup>a</sup>R<sup>b</sup> and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, (optionally substituted phenyl)-C<sub>1</sub>-C<sub>3</sub>-alkyl, acyl and  
20 carboxy-C<sub>1</sub>-C<sub>6</sub>-alkyl; wherein each of R<sup>b</sup> and R<sup>b</sup> is independently selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R is  
25 selected from carboxyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonylaminocarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl; wherein R' is selected from hydrido, phenyl, thienyl and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R<sup>2</sup> is one or more radicals independently  
30 selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>1</sub>-C<sub>3</sub>-alkyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, aryloxy, arylthio,  
35 arylsulfinyl, heteroaryloxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, aryl-

C<sub>1</sub>-C<sub>6</sub>-alkyloxy, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, aryl-C<sub>1</sub>-C<sub>6</sub>-alkoxy-  
 C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-  
 haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-  
 haloalkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-(haloalkyl-C<sub>1</sub>-C<sub>3</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-  
 5 hydroxyalkyl, hydroxyimino-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino,  
 arylamino, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, heteroarylamino,  
 heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, cyano, amino,  
 aminosulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, arylaminosulfonyl,  
 heteroarylaminosulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl,  
 10 heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, heterocyclisulfonyl,  
 C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally  
 substituted aryl, optionally substituted heteroaryl, aryl-  
 C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,  
 heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-  
 15 alkoxycarbonyl, formyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylcarbonyl and C<sub>1</sub>-C<sub>6</sub>-  
 alkylcarbonyl; and wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup>  
 are independently selected from carbon and nitrogen with the  
 proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon;  
 or wherein R<sup>2</sup> together with ring A forms a naphthyl or  
 20 quinolyl radical; or an isomer or pharmaceutically  
 acceptable salt thereof.

A more preferred class of compounds consists of those  
 compounds of Formula I' wherein X is selected from O, S and  
 NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl and  
 25 (optionally substituted phenyl)methyl; wherein R' is  
 selected from hydrido and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R is  
 carboxyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl;  
 wherein R<sup>2</sup> is one or more radicals independently selected  
 from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl,  
 30 halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl,  
 phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-C<sub>3</sub>-  
 alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkylthio, C<sub>1</sub>-C<sub>3</sub>-alkylsulfinyl,  
 phenyloxy, phenylthio, phenylsulfinyl, C<sub>1</sub>-C<sub>3</sub>-haloalkyl-C<sub>1</sub>-C<sub>3</sub>-  
 hydroxyalkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyloxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-  
 35 haloalkyl, C<sub>1</sub>-C<sub>3</sub>-haloalkoxy, C<sub>1</sub>-C<sub>3</sub>-haloalkylthio, C<sub>1</sub>-C<sub>3</sub>-

hydroxyalkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxyimino-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, cyano, amino, aminosulfonyl, N-alkylaminosulfonyl, N-arylamino sulfonyl, N-heteroarylamino sulfonyl, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkylsulfonyl, 5- to 8-membered heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- to 9-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, phenylcarbonyl, 4-chlorophenylcarbonyl, 4-hydroxyphenylcarbonyl, 4-trifluoromethylphenylcarbonyl, 4-methoxyphenylcarbonyl, aminocarbonyl, formyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl, benzofurylphenyl, or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula I' wherein X is selected from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, (4-chloromethyl)benzyl, (4-methoxy)benzyl, and (4-cyano)benzyl, (4-nitro)benzyl; wherein R is carboxyl; wherein R' is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, nitro,

cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, optionally substituted phenylcarbonyl, aminocarbonyl, formyl and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl, or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I' there is a subclass of chromene compounds wherein X is O; wherein R is carboxyl; wherein R<sup>2</sup> is selected from hydrido and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>6</sub>-perfluoroalkyl; wherein R<sup>3</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy, 5- or 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5- or 6-membered heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-heteroarylamino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, nitro, amino, aminosulfonyl, N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-arylaminosulfonyl, N-heteroarylaminosulfonyl, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered heterocyclisulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl;

wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.

- 5       An even more preferred class of compounds consists of those compounds of Formula I' wherein X is O; wherein R is carboxyl; wherein R' is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals
- 10       independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio,
- 15       methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl,
- 20       methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl,
- 25       phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro,
- 30       benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; and wherein one of the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> is nitrogen and the other three are carbon; or an isomer or pharmaceutically acceptable salt thereof.

Another even more preferred class of compounds consists of those compounds of Formula I' wherein X is O; wherein R is carboxyl; wherein R" is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I' there is another subclass of benzothiopyran compounds wherein X is S; wherein R is carboxyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>6</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy,

5- or 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5- or 6-membered heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, N-phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-heteroarylamino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, amino, aminosulfonyl, N-alkylaminosulfonyl, N-arylaminosulfonyl, N-heteroarylaminosulfonyl, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula I' wherein X is S; wherein R is carboxyl; wherein R' is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-

(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and  
5 methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or  
10 pharmaceutically acceptable salt thereof.

Within Formula I' there is a third subclass of dihydroquinoline compounds wherein X is NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, acyl and carboxy-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R is carboxyl; wherein R<sup>1</sup>  
15 is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy, 5- or 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5- or 6-membered  
20 heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, N-phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-heteroarylamino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, amino, aminosulfonyl, N-alkylaminosulfonyl, N-arylaminosulfonyl, N-heteroarylaminosulfonyl, N-(phenyl-C<sub>1</sub>-  
25 C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered heterocyclisulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring  
30 atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.




An even more preferred class of compounds consists of those compounds of Formula I' wherein X is NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, (4-chloromethyl)benzyl, (4-methoxy)benzyl, (4-cyano)benzyl, and (4-nitro)benzyl; wherein R is carboxyl; wherein R<sup>n</sup> is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I' there is a fourth subclass of compounds wherein X is selected from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl,

acyl and carboxy- $C_1$ - $C_3$ -alkyl; wherein R is selected from carboxyl; wherein  $R^1$  is selected from  $C_1$ - $C_3$ -perfluoroalkyl; wherein the A ring atoms  $A^1$ ,  $A^2$ ,  $A^3$  and  $A^4$  are independently selected from carbon and nitrogen with the proviso that at least three of  $A^1$ ,  $A^2$ ,  $A^3$  and  $A^4$  are carbon; and wherein  $R^2$  together with ring A forms a naphthyl or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula I' wherein X is selected from O, S and  $NR^a$ ; wherein  $R^a$  is selected from hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, (4-chloromethyl)benzyl, (4-methoxy)benzyl, and (4-cyano)benzyl, (4-nitro)benzyl; wherein R is carboxyl; wherein  $R'$  is selected from hydrido and ethenyl; wherein  $R^1$  is selected from trifluoromethyl and pentafluoroethyl; wherein the A ring atoms  $A^1$ ,  $A^2$ ,  $A^3$  and  $A^4$  are independently selected from carbon and nitrogen with the proviso that at least three of  $A^1$ ,  $A^2$ ,  $A^3$  and  $A^4$  are carbon; or wherein  $R^2$  together with ring A forms a naphthyl, or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I there is a subclass of compounds of high interest represented by Formula II:




25

wherein X is selected from O,  $NR^a$  and S;  
 wherein  $R^2$  is lower haloalkyl;  
 wherein  $R^3$  is selected from hydrido, and halo;

wherein R<sup>4</sup> is selected from hydrido, halo, lower alkyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, and  
5 5- or 6- membered nitrogen containing heterocyclosulfonyl;  
wherein R<sup>5</sup> is selected from hydrido, lower alkyl, halo, lower alkoxy, and aryl; and  
wherein R<sup>6</sup> is selected from hydrido, halo, lower alkyl, lower alkoxy, and aryl;  
10 or an isomer or pharmaceutically acceptable salt thereof.

A class of compounds of particular interest consists of those compounds of Formula II wherein R<sup>2</sup> is trifluoromethyl or pentafluoroethyl; wherein R<sup>3</sup> is selected from hydrido,  
15 chloro, and fluoro; wherein R<sup>4</sup> is selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, trifluoromethoxy, methoxy, benzylcarbonyl, dimethylaminosulfonyl, isopropylaminosulfonyl, methylaminosulfonyl, benzylaminosulfonyl,  
20 phenylethylaminosulfonyl, methylpropylaminosulfonyl, methylsulfonyl, and morpholinosulfonyl; wherein R<sup>5</sup> is selected from hydrido, methyl, ethyl, isopropyl, tert-butyl, chloro, methoxy, diethylamino, and phenyl; and wherein R<sup>6</sup> is selected from hydrido, chloro, bromo, fluoro, methyl, ethyl,  
25 tert-butyl, methoxy, and phenyl; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I there is a subclass of compounds of high interest represented by Formula IIa:



wherein  $R^3$  is selected from hydrido, lower alkyl, lower hydroxyalkyl, lower alkoxy and halo;

- 5        wherein  $R^4$  is selected from hydrido, halo, lower alkyl, lower alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower
- 10       haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydroxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen containing
- 15       heterocyclosulfonyl;


wherein  $R^5$  is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and

- wherein  $R^6$  is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl,
- 20       phenylalkynyl, lower alkyl, lower alkoxy, formyl and phenyl; or an isomer or pharmaceutically acceptable salt thereof.

- A class of compounds of particular interest consists of those compounds of Formula IIa wherein  $R^3$  is selected from
- 25       hydrido, and chloro; wherein  $R^4$  is selected from chloro, methyl, tert-butyl, methylthio, trifluoromethyl, difluoromethyl, pentafluoromethyl, trifluoromethylsulfide, trifluoromethoxy, cyano, substituted or unsubstituted phenylcarbonyl, and substituted or unsubstituted phenyl;

wherein  $R^3$  is selected from hydrido, methyl, tert-butyl, chloro; and wherein  $R^6$  is selected from hydrido, chloro, thienyl, hydroxyiminomethyl, substituted or unsubstituted phenylethynyl, and substituted or unsubstituted phenyl; or  
 5 an isomer or pharmaceutically acceptable salt thereof.

Within Formula I there is a subclass of compounds of high interest represented by Formula IIb



10

wherein  $R^3$  is selected from hydrido, lower alkyl, lower hydroxyalkyl, lower alkoxy and halo;

wherein  $R^4$  is selected from hydrido, halo, lower alkyl, lower alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydroxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen containing heterocyclosulfonyl;


wherein  $R^5$  is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and

wherein  $R^6$  is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl, phenylalkynyl, lower alkyl, lower alkoxy, formyl and phenyl;

or an isomer or pharmaceutically acceptable salt thereof.

A class of compounds of particular interest consists of those compounds of Formula IIb wherein R<sup>3</sup> is selected from hydrido, and chloro; wherein R<sup>4</sup> is selected from chloro, methyl, tert-butyl, methylthio, trifluoromethyl, difluoromethyl, pentafluoromethyl, trifluoromethylsulfide, trifluoromethoxy, cyano, substituted or unsubstituted phenylcarbonyl, and substituted or unsubstituted phenyl; wherein R<sup>5</sup> is selected from hydrido, methyl, tert-butyl, chloro; and wherein R<sup>6</sup> is selected from hydrido, chloro, thienyl, hydroxyiminomethyl, substituted or unsubstituted phenylethynyl, and substituted or unsubstituted phenyl; or an isomer or pharmaceutically acceptable salt thereof.

Within Formula I there is a subclass of compounds of high interest represented by Formula IIc:



wherein R<sup>a</sup> is selected from hydrido and lower alkyl; wherein R<sup>3</sup> is selected from hydrido, lower alkyl, lower hydroxyalkyl, lower alkoxy and halo; wherein R<sup>4</sup> is selected from hydrido, halo, lower alkyl, lower alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower

aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydroxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen containing heterocyclosulfonyl;

5        wherein R<sup>5</sup> is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and

         wherein R<sup>6</sup> is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl, phenylalkynyl, lower alkyl, lower alkoxy, formyl and phenyl;

10       or an isomer or pharmaceutically acceptable salt thereof.

         A class of compounds of particular interest consists of those compounds of Formula IIc wherein R<sup>3</sup> is selected from hydrido, and chloro; wherein R<sup>4</sup> is selected from chloro,  
15       methyl, tert-butyl, methylthio, trifluoromethyl, difluoromethyl, pentafluoromethyl, trifluoromethylsulfide, trifluoromethoxy, cyano, substituted or unsubstituted phenylcarbonyl, and substituted or unsubstituted phenyl; wherein R<sup>5</sup> is selected from hydrido, methyl, tert-butyl,  
20       chloro; and wherein R<sup>6</sup> is selected from hydrido, chloro, thienyl, hydroxyiminomethyl, substituted or unsubstituted phenylethynyl, and substituted or unsubstituted phenyl; or an isomer or pharmaceutically acceptable salt thereof.

         A family of specific compounds of particular interest  
25       within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as follows:

6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic  
         acid;

7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

30       7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic  
         acid;

2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

7-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-  
35       carboxylic acid;

- 8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-ethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 7-isopropoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6,8-bis(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;



- 7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dimethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6-nitro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- ethyl 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate;
- 6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6,8-difluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 35

- 8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 7-(N,N-diethylamino)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[[ (phenylmethyl) amino] sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(dimethylamino) sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-aminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-(methylamino) sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6-[(4-morpholino) sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(1,1-dimethylethyl) aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(2-methylpropyl) aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-[[ (phenylmethyl) amino] sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-N,N-diethylaminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-(2,2-dimethylpropylcarbonyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 6,8-dichloro-7-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 5 6-[[ (2-furanylmethyl) amino] sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-[(phenylmethyl) sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-[[ (phenylethyl) amino] sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-bromo-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5,6-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-hydroxymethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5,6,7-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6,7,8-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 5,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5-methoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-benzoyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-chlorobenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-(4-chlorophenoxy)-2-trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- 30 6-(4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(3-chloro-4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(hydroxyimino)methyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(hydroxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 8-(1H-benzimidazol-2-yl)-6-chloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 7-(1,1-dimethylethyl)-2-(pentafluoroethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-8-(methoxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(benzyloxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-chloro-8-ethynyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-chloro-8-(2-furanyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(5-chloro-1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 6-chloro-8-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(3,3-dimethyl-1-butynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-chloro-8-[(4-chlorophenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(4-methoxyphenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 6-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-8-[(4-methylthio)phenyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(4-methylsulfonyl)phenyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-fluorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6,8-diiodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(5-chloro-2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 6-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-bromophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-(ethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-(4-methoxyphenyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-2-(trifluoromethyl)-4-ethenyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-2-(trifluoromethyl)-4-phenyl-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-4-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(2,2,2-trifluoro-1-hydroxyethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,8-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 20 6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,7-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 25 8-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 30 7-chloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,7-dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;

- 2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzothiopyran-3-carboxylic acid;  
6,8-dichloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 5
- 6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6,8-dichloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 10
- 6,7-difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 15
- 6-bromo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-(trifluoromethyl)-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 20
- 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 25
- 6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(trifluoromethyl)phenyl]methyl]-3-quinolinecarboxylic acid;  
6-chloro-1-[(4-chlorophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 30
- 6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(methoxy)phenyl]methyl]-3-quinolinecarboxylic acid;  
6-chloro-1-[(4-cyanophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 35
- 6-chloro-1,2-dihydro-1-[(4-nitrophenyl)methyl]-2-(trifluoromethyl)-3-quinolinecarboxylic acid;



6-chloro-1,2-dihydro-1-ethyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;

6-chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylic acid;

5

2-trifluoromethyl-2H-naphtho[1,2-b]pyran-3-carboxylic acid;

2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;

2-trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid;

5-(hydroxymethyl)-8-methyl-2-(trifluoromethyl)-2H-

10

pyrano[2,3-c]pyridine-3-carboxylic acid;

6-(trifluoromethyl)-6h-1,3-dioxolo[4,5-g][1]benzopyran-7-carboxylic acid; and

3-(trifluoromethyl)-3H-benzofuro[3,2-f][1]benzopyran-2-carboxylic acid.

15

A preferred family of specific compounds of particular interest within Formulas I and I' consists of compounds as follows:

20 (S)-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

(S)-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

(S)-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

25 (S)-2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

(S)-7-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

30 (S)-6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

(S)-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

(S)-6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- (S)-6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-ethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic  
5 acid;
- (S)-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-7,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-7-isopropoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic  
25 acid;
- (S)-7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6,8-bis(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-7-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic  
35 acid;

- (S)-6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6,8-dimethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-nitro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-ethyl 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate;
- (S)-6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 (S)-8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6,8-difluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- (S)-8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-7-(N,N-diethylamino)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[[ (phenylmethyl) amino] sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[(dimethylamino) sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6-aminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(methylamino) sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[(4-morpholino) sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-6-[(1,1-dimethylethyl) aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[(2-methylpropyl) aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 (S)-6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-chloro-6-[[ (phenylmethyl) amino] sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-N,N-diethylaminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(2,2-dimethylpropylcarbonyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- (S)-6,8-dichloro-7-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 5 (S)-6-[[ (2-furanylmethyl)amino]sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[(phenylmethyl)sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-[[ (phenylethyl)amino]sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-6-bromo-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-5,6-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 (S)-6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid ;
- (S)-6-hydroxymethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-5,6,7-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- (S)-6,7,8-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-(methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-5,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzothiopyran-3-carboxylic acid;
- 15 (S)-6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-5-methoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-6-benzoyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(4-chlorobenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 (S)-6-(4-hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-chloro-6-(4-chlorophenoxy)-2-trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- (S)-6-(3-chloro-4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-8-chloro-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-6-chloro-8-[(hydroxyimino)methyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(hydroxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-8-(1H-benzimidazol-2-yl)-6-chloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-7-(1,1-dimethylethyl)-2-(pentafluoroethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(methoxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-6-chloro-8-(benzyloxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-ethynyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 (S)-6-chloro-8-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(2-furanyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 (S)-6-chloro-8-(5-chloro-1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 35

- (S)-6-chloro-8-(3,3-dimethyl-1-butynyl)-2-(trifluoromethyl)-  
2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-[(4-chlorophenyl)ethynyl]-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-chloro-8-[(4-methoxyphenyl)ethynyl]-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- (S)-6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-  
10 benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-[(4-methylthio)phenyl]-2-(trifluoromethyl)-  
2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6-chloro-8-[(4-methylsulfonyl)phenyl]-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- (S)-6-bromo-8-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-  
20 carboxylic acid;
- (S)-6-(4-fluorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-  
3-carboxylic acid;
- (S)-6-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- 25 (S)-8-chloro-6-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- (S)-6,8-diiodo-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- (S)-6-(5-chloro-2-thienyl)-2-(trifluoromethyl)-2H-1-  
30 benzopyran-3-carboxylic acid;
- (S)-6-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid;
- (S)-6-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-  
3-carboxylic acid;



- (S)-6-(4-bromophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-(ethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-8-(4-methoxyphenyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 (S)-6-chloro-2-(trifluoromethyl)-4-ethenyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-2-(trifluoromethyl)-4-phenyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-4-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6-(2,2,2-trifluoro-1-hydroxyethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 20 (S)-6,8-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 25 (S)-6,7-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-8-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 30 (S)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-7-chloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 35

- (S)-6,7-dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6,8-dichloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 10 (S)-6,8-dichloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- (S)-6,7-difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- (S)-6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 15 (S)-6-bromo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- (S)-1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 20 (S)-6-(trifluoromethyl)-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- (S)-6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- (S)-6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 25 (S)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(trifluoromethyl)phenyl]methyl]-3-quinolinecarboxylic acid;
- (S)-6-chloro-1-[(4-chlorophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;
- 30 (S)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(methoxy)phenyl]methyl]-3-quinolinecarboxylic acid;
- (S)-6-chloro-1-[(4-cyanophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;

- (S)-6-chloro-1,2-dihydro-1-[(4-nitrophenyl)methyl]-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
(S)-6-chloro-1,2-dihydro-1-ethyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
5 (S)-6-chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylic acid;
- (S)-2-trifluoromethyl-2H-naphtho[1,2-b]pyran-3-carboxylic acid;  
10 (S)-2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;  
(S)-2-trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid; and  
(S)-5-(hydroxymethyl)-8-methyl-2-(trifluoromethyl)-2H-  
15 pyrano[2,3-c]pyridine-3-carboxylic acid.

The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido  
20 radicals may be attached to a carbon atom to form a methylene (-CH<sub>2</sub>-) radical. Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylsulfonyl", it embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one  
25 to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like. Most preferred are  
30 lower alkyl radicals having one to three carbon atoms. The term "alkenyl" embraces linear or branched radicals having at least one carbon-carbon double bond of two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkenyl radicals are "lower  
35 alkenyl" radicals having two to about six carbon atoms.

Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The term "alkynyl" denotes linear or branched radicals having two to about twenty carbon atoms or, preferably, two to about  
5 twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about ten carbon atoms. Most preferred are lower alkynyl radicals having two to about six carbon atoms. Examples of such radicals include propargyl, butynyl, and the like. The terms "alkenyl" and  
10 "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. The term "halo" means halogens such as fluorine, chlorine, bromine or iodine atoms. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms  
15 is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or  
20 more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" embraces radicals having 1-6 carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl,  
25 heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. "Perfluoroalkyl" means alkyl radicals having all hydrogen atoms replaced with fluoro atoms. Examples include trifluoromethyl and  
30 pentafluoroethyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms  
35 and one or more hydroxyl radicals. Examples of such

radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. The term "cyanoalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one cyano radicals. More preferred cyanoalkyl radicals are "lower cyanoalkyl" radicals having one to six carbon atoms and one cyano radical. Examples of such radicals include cyanomethyl. The terms "alkoxy" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one or two rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Said "aryl" group may have 1 to 3 substituents such as lower alkyl, hydroxy, halo, haloalkyl, nitro, cyano, alkoxy and lower alkylamino. The term "heterocyclyl" embraces saturated, partially saturated and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclic radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl]; saturated 3 to 6-membered

heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl]. Examples of partially saturated heterocyclyl radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole. Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals, include unsaturated 5 to 6 membered heteromonocyclyl group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl [e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl]; unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g., tetrazolo [1,5-b]pyridazinyl]; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, 2-furyl, 3-furyl, etc.; unsaturated 5 to 6-membered heteromonocyclic group containing a sulfur atom, for example, 2-thienyl, 3-thienyl, etc.; unsaturated 5- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl]; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzoxazolyl, benzoxadiazolyl]; unsaturated 5 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl]; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl] and the like. The term also embraces radicals where heterocyclic radicals are fused with aryl radicals. Examples of such fused bicyclic radicals

include benzofuran, benzothiophene, and the like. Said "heterocyclyl" group may have 1 to 3 substituents such as lower alkyl, hydroxy, oxo, amino and lower alkylamino. Preferred heterocyclic radicals include five to ten membered fused or unfused radicals. More preferred examples of heteroaryl radicals include benzofuryl, 2,3-dihydrobenzofuryl, benzothienyl, indolyl, dihydroindolyl, chromanyl, benzopyran, thiochromanyl, benzothiopyran, benzodioxolyl, benzodioxanyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl, and pyrazinyl. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals  $-SO_2-$ . "Alkylsulfonyl" embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. "Haloalkylsulfonyl" embraces haloalkyl radicals attached to a sulfonyl radical, where haloalkyl is defined as above. More preferred haloalkylsulfonyl radicals are "lower haloalkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower haloalkylsulfonyl radicals include trifluoromethylsulfonyl. The term "arylalkylsulfonyl" embraces aryl radicals as defined above, attached to an alkylsulfonyl radical. Examples of such radicals include benzylsulfonyl and phenylethylsulfonyl. The terms "sulfamyl," "aminosulfonyl" and "sulfonamidyl," whether alone or used with terms such as "N-alkylaminosulfonyl", "N-arylaminosulfonyl", "N,N-dialkylaminosulfonyl" and "N-alkyl-N-arylaminosulfonyl", denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide ( $-SO_2NH_2$ ). The term "alkylaminosulfonyl" includes "N-alkylaminosulfonyl" and "N,N-dialkylaminosulfonyl" where sulfamyl radicals are substituted, respectively, with one alkyl radical, or two

alkyl radicals. More preferred alkylaminosulfonyl radicals are "lower alkylaminosulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylaminosulfonyl radicals include N-methylaminosulfonyl, N-ethylaminosulfonyl and N-methyl-N-ethylaminosulfonyl. The terms "N-arylamino-sulfonyl" and "N-alkyl-N-arylamino-sulfonyl" denote sulfamyl radicals substituted, respectively, with one aryl radical, or one alkyl and one aryl radical. More preferred N-alkyl-N-arylamino-sulfonyl radicals are "lower N-alkyl-N-arylsulfonyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower N-alkyl-N-arylamino-sulfonyl radicals include N-methyl-N-phenylaminosulfonyl and N-ethyl-N-phenylaminosulfonyl. Examples of such N-aryl-amino-sulfonyl radicals include N-phenylaminosulfonyl. The term "arylalkylaminosulfonyl" embraces aralkyl radicals as described above, attached to an aminosulfonyl radical. The term "heterocyclylamino-sulfonyl" embraces heterocyclyl radicals as described above, attached to an aminosulfonyl radical. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes  $\text{-CO}_2\text{H}$ . The term "carboxyalkyl" embraces radicals having a carboxy radical as defined above, attached to an alkyl radical. The term "carbonyl", whether used alone or with other terms, such as "alkylcarbonyl", denotes  $\text{-(C=O)-}$ . The term "acyl" denotes a radical provided by the residue after removal of hydroxyl from an organic acid. Examples of such acyl radicals include alkanoyl and aroyl radicals. Examples of such lower alkanoyl radicals include formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, trifluoroacetyl. The term "aroyl" embraces aryl radicals with a carbonyl radical as defined above. Examples of aroyl include benzoyl, naphthoyl, and the like and the aryl in said aroyl may be additionally substituted. The term "alkylcarbonyl" embraces radicals having a carbonyl radical substituted with



an alkyl radical. More preferred alkylcarbonyl radicals are "lower alkylcarbonyl" radicals having one to six carbon atoms. Examples of such radicals include methylcarbonyl and ethylcarbonyl. The term "haloalkylcarbonyl" embraces radicals having a carbonyl radical substituted with an haloalkyl radical. More preferred haloalkylcarbonyl radicals are "lower haloalkylcarbonyl" radicals having one to six carbon atoms. Examples of such radicals include trifluoromethylcarbonyl. The term "arylcarbonyl" embraces radicals having a carbonyl radical substituted with an aryl radical. More preferred arylcarbonyl radicals include phenylcarbonyl. The term "heteroarylcarbonyl" embraces radicals having a carbonyl radical substituted with a heteroaryl radical. The term "arylalkylcarbonyl" embraces radicals having a carbonyl radical substituted with an arylalkyl radical. More preferred arylcarbonyl radicals include benzylcarbonyl. The term "heteroarylalkylcarbonyl" embraces radicals having a carbonyl radical substituted with a heteroarylalkyl radical. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. Preferably, "lower alkoxycarbonyl" embraces alkoxy radicals having one to six carbon atoms. Examples of such "lower alkoxycarbonyl" ester radicals include substituted or unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl. The term "aminocarbonyl" when used by itself or with other terms such as "aminocarbonylalkyl", "N-alkylaminocarbonyl", "N-arylaminocarbonyl", "N,N-dialkylaminocarbonyl", "N-alkyl-N-arylaminocarbonyl", "N-alkyl-N-hydroxyaminocarbonyl" and "N-alkyl-N-hydroxyaminocarbonylalkyl", denotes an amide group of the formula  $-C(=O)NH_2$ . The terms "N-alkylaminocarbonyl" and "N,N-dialkylaminocarbonyl" denote aminocarbonyl radicals which have been substituted with one alkyl radical and with two alkyl radicals, respectively. More preferred are "lower

alkylaminocarbonyl" having lower alkyl radicals as described above attached to an aminocarbonyl radical. The terms "N-arylamino carbonyl" and "N-alkyl-N-arylamino carbonyl" denote aminocarbonyl radicals substituted, respectively, with one  
5 aryl radical, or one alkyl and one aryl radical. The term "N-cycloalkylaminocarbonyl" denoted aminocarbonyl radicals which have been substituted with at least one cycloalkyl radical. More preferred are "lower cycloalkylaminocarbonyl" having lower cycloalkyl radicals of three to seven carbon  
10 atoms, attached to an aminocarbonyl radical. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. The term "alkylaminoalkyl" embraces aminoalkyl radicals having the nitrogen atom substituted with an alkyl radical. The term "heterocyclylalkyl" embraces heterocyclic-  
15 substituted alkyl radicals. More preferred heterocyclylalkyl radicals are "5- or 6- membered heteroarylalkyl" radicals having alkyl portions of one to six carbon atoms and a 5- or 6- membered heteroaryl radical. Examples include such radicals as pyridylmethyl and  
20 thienylmethyl. The term "aralkyl" embraces aryl-substituted alkyl radicals. Preferable aralkyl radicals are "lower aralkyl" radicals having aryl radicals attached to alkyl radicals having one to six carbon atoms. Examples of such radicals include benzyl, diphenylmethyl and phenylethyl.  
25 The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The term "arylalkenyl" embraces aryl-substituted alkenyl radicals. Preferable arylalkenyl radicals are "lower arylalkenyl" radicals having aryl radicals attached to  
30 alkenyl radicals having two to six carbon atoms. Examples of such radicals include phenylethenyl. The aryl in said arylalkenyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The term "arylalkynyl" embraces aryl-substituted alkynyl radicals.  
35 Preferable arylalkynyl radicals are "lower arylalkynyl"

radicals having aryl radicals attached to alkynyl radicals having two to six carbon atoms. Examples of such radicals include phenylethynyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. An example of "alkylthio" is methylthio, ( $\text{CH}_3\text{-S-}$ ). The term "haloalkylthio" embraces radicals containing a haloalkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. An example of "haloalkylthio" is trifluoromethylthio. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent  $\text{-S(=O)-}$  atom. The term "arylsulfinyl" embraces radicals containing an aryl radical, attached to a divalent  $\text{-S(=O)-}$  atom. The term "haloalkylsulfinyl" embraces radicals containing a haloalkyl radical, of one to ten carbon atoms, attached to a divalent  $\text{-S(=O)-}$  atom. The terms "N-alkylamino" and "N,N-dialkylamino" denote amino groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. More preferred alkylamino radicals are "lower alkylamino" radicals having one or two alkyl radicals of one to six carbon atoms, attached to a nitrogen atom. Suitable "alkylamino" may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like. The term "arylamino" denotes amino groups which have been substituted with one or two aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical. The term "heteroarylamino" denotes amino groups which have been substituted with one or two heteroaryl radicals, such as N-thienylamino. The "heteroarylamino" radicals may be further substituted on the

heteroaryl ring portion of the radical. The term "aralkylamino" denotes amino groups which have been substituted with one or two aralkyl radicals, such as N-benzylamino. The "aralkylamino" radicals may be further substituted on the aryl ring portion of the radical. The terms "N-alkyl-N-arylamino" and "N-aralkyl-N-alkylamino" denote amino groups which have been substituted with one aralkyl and one alkyl radical, or one aryl and one alkyl radical, respectively, to an amino group. The term "arylthio" embraces aryl radicals of six to ten carbon atoms, attached to a divalent sulfur atom. An example of "arylthio" is phenylthio. The term "aralkylthio" embraces aralkyl radicals as described above, attached to a divalent sulfur atom. An example of "aralkylthio" is benzylthio. The term "aralkylsulfonyl" embraces aralkyl radicals as described above, attached to a divalent sulfonyl radical. The term "heterocyclylsulfonyl" embraces heterocyclyl radicals as described above, attached to a divalent sulfonyl radical. The term "aryloxy" embraces aryl radicals, as defined above, attached to an oxygen atom. Examples of such radicals include phenoxy. The term "aralkoxy" embraces oxy-containing aralkyl radicals attached through an oxygen atom to other radicals. More preferred aralkoxy radicals are "lower aralkoxy" radicals having phenyl radicals attached to lower alkoxy radical as described above.

The present invention comprises a pharmaceutical composition comprising a therapeutically-effective amount of a compound of Formula I in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of treating cyclooxygenase-2 mediated disorders, such as inflammation, in a subject, the method comprising treating the subject having or susceptible to such

disorder with a therapeutically-effective amount of a compound of Formula I.

Also included in the family of compounds of Formula I are the stereoisomers thereof. Compounds of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or nonracemic mixtures thereof. Accordingly, some of the compounds of this invention may be present in racemic mixtures which are also included in this invention. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active base and then separation of the mixture of diastereoisomers by crystallization, followed by liberation of the optically active bases from these salts. Examples of appropriate bases are brucine, strychnine, dehydroabietylamine, quinine, cinchonidine, ephedrine,  $\alpha$ -methylbenzylamine, amphetamine, deoxyphedrine, chloramphenicol intermediate, 2-amino-1-butanol, and 1-(1-naphthyl)ethylamine. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of Formula I can likewise be obtained by utilizing optically active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt. Additional methods for resolving optical isomers, known to those skilled in the art may be used, for example, those discussed by J. Jaques et al in Enantiomers.

Racemates, and Resolutions, John Wiley and Sons, New York (1981).

Also included in the family of compounds of Formula I and I' are the amide protected acids thereof. Thus primary and secondary amines can be reacted with the chromene-3-carboxylic acids of Formula I and I' to form amides which can be useful as prodrugs. Preferred amines heterocyclicamines, including optionally substituted aminothiazoles, optionally substituted amino-isoxazoles, and optionally substituted aminopyridines; aniline derivatives; sulfonamides; aminocarboxylic acids; and the like. Additionally, 1-acyldihydroquinolines can behave as prodrugs for the 1H-dihydroquinolines.


Also included in the family of compounds of Formula I and I' are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, salicylic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic,

stearic, algenic,  $\beta$ -hydroxybutyric, salicyclic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I or I' include metallic salts, such as salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc, or salts made from organic bases including primary, secondary and tertiary amines, substituted amines including cyclic amines, such as caffeine, arginine, diethylamine, N-ethyl piperidine, histidine, glucamine, isopropylamine, lysine, morpholine, N-ethyl morpholine, piperazine, piperidine, triethylamine, trimethylamine. All of these salts may be prepared by conventional means from the corresponding compound of the invention by reacting, for example, the appropriate acid or base with the compound of Formula I or I'.


#### GENERAL SYNTHETIC PROCEDURES

The compounds of the invention can be synthesized according to the following procedures of Schemes 1-16, wherein the  $R^1$ - $R^6$  substituents are as defined for Formulas I-II, above, except where further noted.

#### SCHEME 1



Synthetic Scheme 1 illustrates the general method for the preparation of a wide variety of substituted 2H-1-benzopyran derivatives 3 and 4. In step 1, a representative ortho-hydroxybenzaldehyde (salicylaldehyde) derivative 1 is condensed with an acrylate derivative 2 in the presence of base, such as potassium carbonate in a solvent such as dimethylformamide, to afford the desired 2H-1-benzopyran ester 3. An alternative base-solvent combination for this condensation includes an organic base such as triethylamine and a solvent such as dimethyl sulfoxide. In step 2 the ester is hydrolyzed to the corresponding acid, such as by treatment with aqueous base (sodium hydroxide) in a suitable solvent such as ethanol to afford after acidification the substituted 2H-1-benzopyran-3-carboxylic acid 4.


**SCHEME 2**

Synthetic Scheme 2 shows the general method for functionalizing selected 2H-1-benzopyrans. Treatment of the 2H-1-benzopyran carboxylic acid 4 or ester 3 with an electrophillic agent makes a 6-substituted 2H-1-benzopyran 5. A wide variety of electrophillic agents react selectively with 2H-1-benzopyrans 4 in the 6-position to provide new analogs in high yield. Electrophillic reagents such as halogen (chlorine or bromine) give the 6-halo derivatives. Chlorosulfonic acid reacts to afford the 6-position sulfonyl chloride that can further be converted to a sulfonamide or sulfone. Friedel-Crafts acylation of 4 provides 6-acylated 2H-1-benzopyrans in good to excellent yield. A number of other electrophiles can be used to selectively react with these 2H-1-benzopyrans in a similar



manner. A 6-position substituted 2H-1-benzopyran can react with an electrophilic reagent at the 8-position using similar chemistries to that described for electrophilic substitution of the 6-position. This yields an 2H-1-  
 5 benzopyran which is substituted at both the 6 and 8 positions.

## SCHEME 3




Synthetic Scheme 3 illustrates a second general synthesis of substituted 2H-1-benzopyran-3-carboxylic acids which allows substitution at position 4 of the 2H-1-  
 15 benzopyran. In this case a commercially or synthetically available substituted ortho-hydroxy acetophenone 6 is treated with two or more equivalents of a strong base such as lithium bis(trimethylsilyl)amide in a solvent such as tetrahydrofuran (THF), followed by reaction with diethyl


carbonate to afford the beta-keto ester 7. Ester 7 is condensed with an acid chloride or anhydride in the presence of a base such as potassium carbonate in a solvent such as toluene with heat to afford 4-oxo-4H-1-benzopyran 8.

- 5 Reduction of the olefin can be accomplished by a variety of agents including sodium borohydride ( $\text{NaBH}_4$ ) in solvent mixtures such as ethanol and tetrahydrofuran (THF), or by use of triethylsilane in a solvent such as trifluoroacetic acid, or by catalytic reduction using palladium on charcoal
- 10 and hydrogen gas in a solvent such as ethanol to yield the new beta-keto ester 9 (two tautomeric structures shown). Acylation of the oxygen of the ketone enolate in the presence of a base such as 2,6-di-tert-butyl-4-methylpyridine, an acylating agent such as
- 15 trifluoromethanesulfonic anhydride, and using a solvent such as methylene chloride yields the enol-triflate 10. Triflate 10 can be reduced with reagents such as tri-n-butyltin hydride, lithium chloride and a palladium (0) catalyst such as tetrakis(triphenylphosphine)palladium (0) in a solvent
- 20 such as tetrahydrofuran to yield 2H-1-benzopyran ester 11 where  $\text{R}''$  is hydrogen. The ester 11 can be saponified with a base such as 2.5 N sodium hydroxide in a mixed solvent such as tetrahydrofuran-ethanol-water (7:2:1) to yield the desired substituted 2H-1-benzopyran-3-carboxylic acid.
- 25 To incorporate a carbon fragment  $\text{R}^1$  one can treat triflate 10 with reagents known to undergo "cross-coupling" chemistries such as tributylethynyltin, lithium chloride and a palladium(0) catalyst such as tetrakis(triphenylphosphine)palladium (0) in a solvent such
- 30 as tetrahydrofuran to yield 2H-1-benzopyran ester 11 where  $\text{R}^1$  is a vinyl moiety. The ester 6 can be saponified with a base such as 2.5 N sodium hydroxide in a mixed solvent such as tetrahydrofuran-ethanol-water (7:2:1) to yield the desired 4-vinyl-2H-1-benzopyran-3-carboxylic acid (12,  $\text{R}'' =$
- 35  $\text{CH}_2\text{CH}-$ ). Similarly triflate 10 can be converted under similar conditions using tri-n-butylphenyltin to 2H-1-benzopyran where  $\text{R}^1 = \text{phenyl}$  and by hydrolysis of the ester

converted to the carboxylic acid 12 where R' = phenyl.  
 Using a similar strategy, substituents which be incorporated  
 as substituent R' can be substituted olefins, substituted  
 aromatics, substituted heteroaryl, acetylenes and substituted  
 5 acetylenes.

**SCHEME 4**

Synthetic Scheme 4 shows an alternative general  
 procedure for the preparation of 4-oxo-4H-1-benzopyran 8.  
 Treatment of an ortho-fluorobenzoyl chloride with an  
 10 appropriately substituted beta-keto ester 14 with a base  
 such as potassium carbonate in a solvent such as toluene  
 provides 4-oxo-4H-1-benzopyran 8. 4-oxo-4H-1-benzopyran 8  
 can be converted to 2H-1-benzopyran 12 as described in  
 Scheme 3.

**SCHEME 5**

Y = Br, I, CF<sub>3</sub>SO<sub>3</sub>

Synthetic Scheme 5 shows a general method for  
 substitution of the aromatic ring of the 2H-1-benzopyran.


20 This can be accomplished through organo-palladium mediated  
 "cross-coupling" chemistries using a palladium (0) catalyst  
 to couple benzopyran 15 at position Y, where Y is iodide,  
 bromide or triflate, with an acetylene, olefin, nitrile, or  
 aryl coupling agent. Substituted acetylenes as the coupling  
 25 agent will provide the corresponding substituted acetylene.

Substituted aryl moieties can be incorporated using arylboronic acids or esters; nitriles can be incorporated by use of zinc (II) cyanide. The resulting ester **16** can be converted to carboxylic acid **17** as described in Scheme 1.

5 Another approach to substitution of the aryl moiety of the benzopyran **15** is to convert Y, where Y is iodide or bromide, to a perfluoroalkyl moiety. Exemplary of this transformation is the conversion of **15** (Y = iodide) to **16** (R<sup>2'</sup> = pentafluoroethyl) using a potassium  
10 pentafluoropropionate and copper (I) iodide in hexamethylphosphoramide (HMPA). The resulting ester **16** can be converted to carboxylic acid **15** as described in Scheme 1.

A similar method adds substitution of the aromatic ring in dihydroquinoline-3-carboxylates. This can be  
15 accomplished through organopalladium couplings with aryl iodides, bromides, or triflates and various coupling agents (R. F. Heck, *Palladium Reagents in Organic Synthesis*. Academic Press 1985). When using a suitable palladium  
catalyst such as tetrakis(triphenyl-phosphine)palladium(0) in  
20 this reaction, coupling agents such as alkynes provide disubstituted alkynes, phenyl boronic acids afford biphenyl compounds, and cyanides produce arylcyano compounds. A number of other palladium catalysts and coupling reagents  
could be used to selectively react with appropriately  
25 substituted dihydroquinoline-3-carboxylates in a similar manner.

#### SCHEME 6



30 Synthetic Scheme 6 shows a general synthetic route for conversion of a commercially or synthetically available


substituted phenol into a substituted salicylaldehyde. Several different methods which utilize formaldehyde or a chemically equivalent reagent are described in detail below.

Reaction of an appropriately substituted phenol **18** in  
5 basic media with formaldehyde (or chemical equivalent) will yield the corresponding salicylaldehyde **1**. The intermediate, ortho-hydroxymethylphenol **19**, will under appropriate reaction conditions be oxidized to the salicylaldehyde **1** in situ. The reaction commonly employs  
10 ethyl magnesium bromide or magnesium methoxide (one equivalent) as the base, toluene as the solvent, paraformaldehyde (two or more equivalents) as the source of formaldehyde, and employs hexamethylphosphoramide (HMPA) or *N,N,N',N'*-tetramethylethylenediamine (TMEDA). (See:  
15 Casiraghi, G. et al., J.C.S. Perkin I, **1978**, 318-321.)

Alternatively an appropriately substituted phenol **18** may react with formaldehyde under aqueous basic conditions to form the substituted ortho-hydroxybenzyl alcohol **19** (See:  
a) J. Leroy and C. Wakselman, J. Fluorine Chem., **40**, 23-32  
20 (1988). b) A. A. Moshfegh, et al., Helv. Chim. Acta., **65**, 1229-1232 (1982)). Commonly used bases include aqueous potassium hydroxide or sodium hydroxide. Formalin (38% formaldehyde in water) is commonly employed as the source of formaldehyde. The resulting ortho-hydroxybenzyl alcohol **19**  
25 can be converted to the salicylaldehyde **1** by an oxidizing agent such as manganese (IV) dioxide in a solvent such as methylene chloride or chloroform (See: R-G. Xie, et al., Synthetic Commun. **24**, 53-58 (1994)).


An appropriately substituted phenol **18** can be treated  
30 under acidic conditions with hexamethylenetetramine (HMTA) to prepare the salicylaldehyde **1** (Duff Reaction; See: Y. Suzuki, and H. Takahashi, Chem. Pharm. Bull., **31**, 1751-1753 (1983)). This reaction commonly employs acids such as acetic acid, boric acid, methanesulfonic acid, or  
35 trifluoromethanesulfonic acid. The source of formaldehyde commonly used is hexamethylenetetramine.

## SCHEME 7




Synthetic Scheme 7 shows the Reimer-Tiemann reaction in which an commercially or synthetically available appropriately substituted phenol 18 will under basic conditions react with chloroform to yield a substituted salicylaldehyde 1 (See: Cragoe, E.J.; Schultz, E.M., U.S. Patent 3 794 734, 1974).

## SCHEME 8




Synthetic Scheme 8 shows the conversion of a commercially or synthetically available appropriately substituted salicylic acid 21 to its respective salicylaldehyde 1 via an intermediate 2-hydroxybenzyl alcohol 19. Reduction of the salicylic acid 21 can be accomplished with a hydride reducing agent such as borane in a solvent such as tetrahydrofuran. Treatment of the intermediate 2-hydroxybenzyl alcohol 19 with an oxidizing agent such as manganese (IV) oxide in a solvent such as methylene chloride or chloroform provides salicylaldehyde 1.

## SCHEME 9



Synthetic Scheme 9 illustrates a general synthetic method for preparation of a wide variety of substituted 2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acids (25). In step 1, an appropriately commercially or synthetically available substituted thiophenol 22 is ortho-metallated with a base such as n-butyllithium employing TMEDA (*N,N,N',N'*-tetramethylethylenediamine) followed by treatment with dimethylformamide to provide the 2-mercaptobenzaldehyde 23. Condensation of the 2-mercaptobenzaldehyde 23 with an acrylate 2 in the presence of base provides ester 24 which can be saponified in the presence of aqueous base to afford the substituted 2H-1-benzothiopyran-3-carboxylic acids 25.


## SCHEME 10



Synthetic Scheme 10 shows a method for preparing a substituted 2-mercaptobenzaldehyde from an appropriate commercially or synthetically available substituted salicylaldehyde. In step 1, the phenolic hydroxyl of salicylaldehyde 1 is converted to the corresponding O-aryl thiocarbamate 26 by acylation with an appropriately substituted thiocarbamoyl chloride such as *N,N*-dimethylthiocarbamoyl chloride in a solvent such as dimethylformamide using a base such as triethylamine. In Step 2, O-aryl thiocarbamate 26 rearranges to S-aryl thiocarbamate 27 when heated sufficiently such as to 200 °C using either no solvent or a solvent such as *N,N*-dimethylaniline (See: A. Levai, and P. Sebok, Synth. Commun., 22 1735-1750 (1992)). Hydrolysis of S-aryl thiocarbamate 27 with a base such as 2.5 N sodium hydroxide in a solvent mixture such as tetrahydrofuran and ethanol yields the substituted 2-mercaptobenzaldehyde 23 which can be converted to the substituted 2H-1-benzothiopyran-3-carboxylic acids 25 as described in Scheme 9.




## SCHEME 11




Synthetic Scheme 11 illustrates the general method for the preparation of a wide variety of dihydroquinoline-3-carboxylic acid derivatives 30. R<sup>2</sup> represents the aromatic substitution of commercially and synthetically available 2-aminobenzaldehydes 28. The 2-amino-benzaldehyde derivative 28, where R<sup>2</sup> represents various substitutions, is condensed with an acrylate derivative 2 in the presence of a base such as potassium carbonate, triethylamine, or diazabicyclo[2.2.2]undec-7-ene in solvents such as dimethylformamide to afford the dihydroquinoline-3-carboxylate esters 29. The ester 29 can be saponified to the corresponding acid, such as by treatment with aqueous inorganic base such as 2.5 N sodium hydroxide in a suitable solvent such as ethanol to afford after acidification the desired dihydroquinoline-3-carboxylic acid 30.

## SCHEME 12




Synthetic Scheme 12 illustrates the preparation of dihydroquinoline-3-carboxylic acid **30** from 2-aminobenzoic acids **31**.  $R^2$  represents the aromatic substitution of commercially and synthetically available 2-aminobenzoic acids **31**. Reduction of the representative 2-aminobenzoic acid **31** to the desired 2-aminobenzyl alcohol **32** was accomplished with a hydride reducing agent such as borane in a solvent such as tetrahydrofuran. Treatment of the desired 2-aminobenzyl alcohol **32** with an oxidizing agent such as manganese(IV)oxide in a solvent such as methylene chloride provides the representative 2-aminobenzaldehydes **28**. (C. T. Alabaster, et al. *J. Med. Chem.* 31, 2048-2056 (1988)) The 2-aminobenzaldehydes were converted to the desired dihydroquinoline-3-carboxylic acid **30** as described in Scheme 11.

## SCHEME 13



Synthetic Scheme 13 illustrates the general method for the preparation of a wide variety of dihydroquinoline-3-carboxylic acid derivatives **30** from isatins **33**.  $R^2$  represents the aromatic substitution of commercially and synthetically available isatins **33**. A representative isatin **33** was treated with basic peroxide generated from hydrogen peroxide and a base such as sodium hydroxide to afford the desired representative 2-aminobenzoic acids **31**. (M. S. Newman and M. W. Lougue, J. Org. Chem., 36, 1398-1401 (1971)) The 2-aminobenzoic acids **31** are subsequently converted to the desired dihydroquinoline-3-carboxylic acid derivatives **30** as described in synthetic Scheme 12.


**SCHEME 14**

15

Synthetic Scheme 14 is another general method for the preparation of dihydroquinoline-3-carboxylic acid derivatives **30**. In step 1, an appropriate commercially or synthetically available substituted aniline **34** can be treated with an acylating reagent such as pivaloyl chloride

20

yielding an amide **35**. The *ortho*-dianion of amide **35** is prepared by treating amide **35** with organo-lithium bases such as *n*-butyllithium or *tert*-butyllithium in tetrahydrofuran at low temperature. The dianion is quenched with dimethylformamide to afford the acylated-2-amino-benzaldehydes **36**. (J. Turner, *J. Org. Chem.*, 48, 3401-3408 (1983)) Reaction of these aldehydes in the presence of bases such as lithium hydride with an acrylate followed by work up with aqueous inorganic bases and hydrolysis, such as by treatment with aqueous base (sodium hydroxide) in a suitable solvent such as ethanol affords, after acidification, a dihydroquinoline-3-carboxylic acid **30**.

**SCHEME 15**


Synthetic Scheme 15 shows a general method for alkylation of the nitrogen of dihydroquinoline-3-carboxylate ester derivatives **29**. The step involves treatment of dihydroquinoline-3-carboxylate ester derivatives **29** with alkyl halides such as iodoethane in the presence of phase transfer catalysts such as tetrabutylammonium iodide, and a base such as caustic (50% aqueous sodium hydroxide) in a solvent such as dichloromethane. These conditions afford the N-alkylated dihydroquinoline-3-carboxylate esters **37**.

Saponification of 37 with aqueous base provides N-alkylated-dihydroquinoline-3-carboxylic acid derivatives 38.

The following examples contain detailed descriptions of the methods of preparation of compounds of Formulas I-  
5 II. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for  
10 illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated. All compounds showed NMR spectra consistent with their assigned structures.

The following abbreviations are used:

- 15 HCl - hydrochloric acid
- MgSO<sub>4</sub> - magnesium sulfate
- Na<sub>2</sub>SO<sub>4</sub> - sodium sulfate
- DMF - dimethylformamide
- THF - tetrahydrofuran
- 20 NaOH - sodium hydroxide
- EtOH - ethanol
- K<sub>2</sub>CO<sub>3</sub> - potassium carbonate
- CDCl<sub>3</sub> - deuterated chloroform
- CD<sub>3</sub>OD - deuterated methanol
- 25 Et<sub>2</sub>O - diethyl ether
- EtOAc - ethyl acetate
- NaHCO<sub>3</sub> - sodium bicarbonate
- KHSO<sub>4</sub> - potassium sulfate
- NaBH<sub>4</sub> - sodium borohydride
- 30 TMEDA - tetramethylethylenediamine
- HMTA - hexamethylenetetraamine
- DMSO - dimethyl sulfoxide
- HMPA hexamethyl phosphoric triamide

**EXAMPLE 1**

5       **6-Chloro-2-trifluoromethyl-2H-1-benzopyran-3-**  
          **carboxylic acid**

Step 1. Preparation of ethyl 6-chloro-2-  
trifluoromethyl-2H-1-benzopyran-3-carboxylate.


10       A mixture of 5-chlorosalicylaldehyde (20.02 g,  
0.128 mole) and ethyl 4,4,4-trifluorocrotonate (23.68  
g, 0.14 mole) was dissolved in anhydrous DMF, warmed to  
60 °C and treated with anhydrous K<sub>2</sub>CO<sub>3</sub> (17.75 g, 0.128  
15       mole). The solution was maintained at 60 °C for 20  
hours, cooled to room temperature, and diluted with  
water. The solution was extracted with ethyl acetate.  
The combined extracts were washed with brine, dried  
over anhydrous MgSO<sub>4</sub>, filtered and concentrated in vacuo  
to afford 54.32 g of an oil. The oil was dissolved in  
20       250 mL of methanol and 100 mL of water, whereupon a  
white solid formed that was isolated by filtration,  
washed with water and dried in vacuo, to afford the  
ester as a yellow solid (24.31 g, 62%): mp 62-64 °C. <sup>1</sup>H  
NMR (CDCl<sub>3</sub>/90 MHz) 7.64 (s, 1H), 7.30-7.21 (m, 2H),  
25       6.96 (d, 1H, J = Hz), 5.70 (q, 1H, J = Hz), 4.30 (q,  
2H, J = 7.2 Hz), 1.35 (t, 3H, J = 7.2 Hz).

Step 2. Preparation of 6-chloro-2-  
trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

30       A solution of the ester from Step 1 (13.02 g,  
42 mmole) was dissolved in 200 mL of methanol and  
20 mL of water, treated with lithium hydroxide  
(5.36 g, 0.128 mole) and stirred at room  
temperature for 16 hours. The reaction mixture  
35       was acidified with 1.2 N HCl, whereupon a solid

formed that was isolated by filtration. The solid was washed with 200 mL of water and 200 mL of hexanes and dried in vacuo to afford the title compound as a yellow solid (10.00 g, 85%): mp 181-184 °C.

## EXAMPLE 2



### 6-(Methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

#### Step 1. Preparation of 5-(methylthio)salicylaldehyde.

Ethyl magnesium bromide (38 mL of a 3.0 M solution in diethyl ether, 113.8 mmole) was chilled with an ice-water bath. To the chilled solution was added a solution of 4-(methylthio)phenol (15.95 g, 113.8 mmole) in diethyl ether (30 mL) over 0.15 hour during which time gas was evolved. The reaction was held at 0 °C for 0.5 hour, at room temperature for 0.5 hour, and the addition funnel replaced with a distillation head. Toluene (250 mL) and the diethyl ether were distilled out of the reactor. The reaction was cooled, toluene (250 mL) and hexamethylphosphoramide (HMPA) (19.8 mL, 20.4 g, 113.8 mmole) were added, and the resulting mixture was stirred for 0.25 hours. The distillation head was replaced with a condenser and paraformaldehyde (8.5 g, 284.4 mmole) was added. The reaction was heated to 90 °C for 3 hours. The reaction mixture was cooled to room temperature, was acidified with 1N HCl and the layers separated. The organic phase was washed with water, and with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo to yield a solid. This solid was purified by silica chromatography (hexanes-ethyl acetate, 5:1) yielding the salicylaldehyde as a yellow

crystalline solid (6.01 g) of suitable purity to be used in the next reaction without further purification.

Step 2. Preparation of ethyl 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

5-Methylthiosalicylaldehyde (Step 1) (2.516 g, 14.96 mmole) was added to dimethylformamide (3.5 mL), potassium carbonate (2.27 g, 16.45 mmole) and ethyl 4,4,4-trifluorocrotonate (3.3 mL, 3.8 g, 22.4 mmole). The mixture was heated to 65 °C for 3 h. The reaction was cooled to room temperature, poured into H<sub>2</sub>O (50 mL), and extracted with diethyl ether (2 X 75 mL). The combined ethereal phases were washed with aqueous NaHCO<sub>3</sub> solution (3 X 50 mL), aqueous 2 N HCl solution (3 X 50 mL), and brine (3 X 50 mL), dried over MgSO<sub>4</sub>, filtered, diluted with isooctane and partially concentrated in vacuo causing the precipitation of the ethyl ester (2.863 g, 60 %) as a yellow powder: mp 87.8-89.6 °C This ester was of suitable purity to use without further purification.


Step 3. Preparation of 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester (Step 2) was hydrolyzed to form the carboxylic acid via a method similar to that described in Example 1, Step 2: mp 166.3-167.9 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.87 (s, 1H), 7.43 (d, 1H, J = 2.2 Hz), 7.33 (dd, 1H, J = 8.5, 2.4 Hz), 6.98 (d, 1H, J = 8.5 Hz), 5.79 (q, 1H, J = 7.0 Hz), 2.48 (s, 3H). FABLRMS m/z 291 (M+H). ESHRMS m/z 289.0152 (M-H, Calc'd 289.0146). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>3</sub>S<sub>1</sub>: C, 49.66; H, 3.13; S, 11.05. Found: C, 49.57; H, 3.02; S, 11.37.

### EXAMPLE 3



82




**7-Methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5

3-Methylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 202.1-203.1 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.84 (s, 1H), 7.12 (d, 1H, *J* = 8.3 Hz), 6.82 (m, 2H), 5.65 (q, 1H, *J* = 6.8 Hz), 2.35 (s, 3H). FABLRMS *m/z* 259 (M+H). FABHRMS *m/z* 259.0576 (M+H, Calc'd 259.0582). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>3</sub>: C, 55.82; H, 3.51. Found: C, 55.93; H, 3.59.

15

**EXAMPLE 4**




**2,7-bis(Trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

20


3-(Trifluoromethyl)phenol was converted to the title compound by a procedure similar to that described in Example 2: mp 190.3-193.5 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.98 (s, 1H), 7.73 (d, 1H, *J* = 7.9 Hz), 7.46 (d, 1H, *J* = 7.9 Hz), 7.36 (s, 1H), 5.93 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 313 (M+H). FABHRMS *m/z* 313.0267 (M+H, Calc'd 313.0299). Anal. Calc'd for C<sub>12</sub>H<sub>6</sub>F<sub>6</sub>O<sub>3</sub>: C, 46.17; H, 1.94. Found: C, 46.25; H, 2.00.

30

**EXAMPLE 5**

5       **7-Bromo-2-trifluoromethyl-2H-1-benzopyran-3-**  
          **carboxylic acid**

3-Bromophenol was converted to the title compound by a procedure similar to that described  
10 in Example 2: mp 198.4-199.5 °C. <sup>1</sup>H NMR (acetone-  
d<sub>6</sub>/300 MHz) 7.89 (s, 1H), 7.43 (d, 1H, *J* = 8.1  
Hz), 7.31 (s, 1H), 7.30 (d, 1H, *J* = 8.1 Hz), 5.84  
(q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 323 (M+H).  
Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>BrF<sub>3</sub>O<sub>3</sub>: C, 40.90; H, 1.87.  
15 Found: C, 41.00; H, 1.85.

**EXAMPLE 6**

20


**6-Chloro-7-methyl-2-trifluoromethyl-2H-1-**  
          **benzopyran-3-carboxylic acid**

4-Chloro-3-methylphenol was converted to the  
25 title compound by a procedure similar to that  
described in Example 2: mp 207.5-209.3 °C. <sup>1</sup>H NMR  
(CDCl<sub>3</sub>/300 MHz) 7.77 (s, 1H), 7.23 (s, 1H), 6.90  
(s, 1H), 5.65 (q, 1H, *J* = 6.8 Hz), 2.37 (s, 3H).  
FABLRMS *m/z* 292 (M+H). FABHRMS *m/z* 299.0287  
30 (M+Li, Calc'd 299.0274). Anal. Calc'd for

$C_{12}H_8ClF_3O_3$ : C, 49.25; H, 2.76; Cl, 12.11. Found:  
C, 49.37; H, 2.82; Cl, 12.17.

### EXAMPLE 7

5



#### 6-(4-Methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid


10

4-(4-Methoxyphenyl)phenol was converted to the title compound by a procedure similar to that described in Example 2: mp 181.7-182.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.87 (s, 1H), 7.11 (m, 1H), 7.02 (m, 2H), 6.98 (m, 4H), 5.81 (q, 1H,  $J = 7.0$  Hz), 3.80 (s, 3H). FABLRMS  $m/z$  365 (M-H). FABHRMS  $m/z$  367.0809 (M+H, Calc'd 367.0793). Anal. Calc'd for  $C_{18}H_{13}F_3O_5$ : C, 59.02; H, 3.58. Found: C, 59.10; H, 3.61.

15

### EXAMPLE 8

20



#### 6-Chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

25

##### Step 1. Preparation of 4-tert-butylsalicylaldehyde.

A five liter three-neck round bottom flask equipped with overhead mechanical stirrer and condenser was charged with trifluoroacetic acid (2.4 L). A mixture of 3-tert-butylphenol (412 g, 2.8 mole) and HMTA (424 g, 3.0 mole) was

30

added portion-wise causing an exotherm. With cooling, the temperature was maintained under 80 °C. The reaction was heated at 80 °C for one hour, then cooled, and water (2 L) added. After 0.5 hour additional water (4 L) was added and the mixture was extracted with ethyl acetate (6 L). The organic extract was washed with water and brine. The resulting organic phase was divided into 2 L volumes and each diluted with water (1 L), and solid NaHCO<sub>3</sub> added until the mixture was neutralized. The organic phases were isolated and combined, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding an oil. This oil was distilled at 95 °C (0.8 mm) yielding the desired salicylaldehyde as an oil (272.9 g, 56 %) which was of sufficient purity to be used without further purification.

Step 2. Preparation of ethyl 7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

A one liter three-neck flask was charged with 4-tert-butylsalicylaldehyde (Step 1) (100.0 g, 0.56 mole), dimethylformamide (110 mL), and potassium carbonate (79.9 g, 0.58 mole) causing the temperature of the mixture to rise to 40 °C. Ethyl 4,4,4-trifluorocrotonate (118.0 g, 0.70 mole) in dimethylformamide (110 mL) was added and the mixture heated to 60 °C at which time the reaction temperature rose to 70 °C. The reaction was cooled to 60 °C, maintained at 60 °C (with added heating) for 8.5 hours and cooled to room temperature. Ethyl acetate (600 mL) and 3 N HCl (600 mL) were added, mixed, and the layers separated. The aqueous phase was extracted with ethyl acetate and the organic phases were combined. The combined organic phases were washed with brine-water (1:1), brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo, yielding a semi-solid. Hexane (600 mL) was added with mixing and the mixture was filtered. The filtrate was washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding a solid. This solid was dissolved in hot ethanol (600 mL). Water

(190 mL) was added which induced crystallization. Filtration of the mixture and drying of the product provided the desired ester as a crystalline solid (131.3 g, 71%): mp 91.0-94.9 °C. This material was of suitable purity to be  
5 used in subsequent steps without further purification.

Step 3. Preparation of ethyl 6-chloro-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


10 A one liter three-neck flask equipped with mechanical stirrer and gas inlet tube was charged with the ester (Step 2) (100 g, 0.3 mole) and acetic acid (300 mL). While cooling (water bath) the reaction mixture, chlorine gas (37.6 g, 0.53 mole) was added which caused the temperature  
15 to rise to 48 °C. After stirring for two hours, the reaction was cooled in an ice-water bath to 15 °C. Zinc powder (19.5 g, 0.3 mole) was added in one portion which caused the temperature to rise to 72 °C. After cooling to room temperature additional zinc powder (5.0 g, 0.08 mole)  
20 was added and the mixture was stirred for 0.5 hour longer. The crude mixture was filtered through diatomaceous earth and was concentrated in vacuo yielding an oil. The oil was dissolved in ethyl acetate (700 mL) washed with brine-water (1:1, 1 L) and brine (0.5 L). The resulting aqueous phase  
25 was extracted with ethyl acetate (700 mL). This ethyl acetate phase was washed with brine-water (1:1, 1 L) and brine (0.5 L). The combined organic phases were dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding the title compound as a yellow oil (116 g, 106 %). This material,  
30 which contained some entrained ethyl acetate, was of suitable purity to be used in subsequent steps without further purification.

Step 4. Preparation of 6-chloro-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

35 To a solution of the ester (Step 3) (116 g, 0.3 mole) in methanol (500 mL) and tetrahydrofuran

(500 mL) in a one liter flask was added aqueous sodium hydroxide (2.5 N, 240 mL, 0.6 mole). After stirring overnight, the pH of the solution was adjusted to 1 with concentrated hydrochloric acid and the solution was extracted with ethyl acetate. The ethyl acetate phase was dried over  $\text{MgSO}_4$ , filtered and concentrated in vacuo yielding a solid. This solid was dissolved in hot ethanol (500 mL). Water (500 mL) was added and upon cooling to room temperature crystals formed which were collected by vacuum filtration. The crystals were washed with ethanol-water (3:7, 3 X 200 mL) and dried providing the title acid as a crystalline solid (91.6 g, 91 %): mp 194.9-196.5 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.86 (s, 1H), 7.52 (s, 1H), 7.12 (s, 1H), 5.83 (q, 1H,  $J = 7.1$  Hz), 1.48 (s, 9H). Anal. Calc'd for  $\text{C}_{15}\text{H}_{14}\text{ClF}_3\text{O}_3$ : C, 53.83; H, 4.22; Cl, 10.59. Found: C, 53.92; H, 4.24; Cl, 10.50.

### EXAMPLE 9




#### 6-(3-Chloro-4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

To a stirred solution of chlorine in acetic acid (3.5 mL of 0.24 M solution, 0.84 mmol) was added 6-(4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (0.31 g, 0.85 mmol) (Example 7). After 1 hour additional chlorine in acetic acid (1.5 mL of 0.24 M solution, 0.36 mmol) was added. After three additional

hours additional chlorine in acetic acid (0.25 ml of 0.25 M solution, 0.06 mmol) was added. After 2.5 hours the reaction was quenched with aqueous 10% sodium bisulfite solution and the resulting mixture extracted with ethyl acetate. The organic phase was washed with water, brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo* yielding a brown oil. The oil was dissolved in a minimum of hexanes which induced crystallization. Vacuum filtration of the mixture provided the title compound as yellow crystals (0.18 g, 53%): mp 205–207 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.89 (s, 1H), 6.97–7.18 (m, 6H), 5.83 (q, 1H,  $J = 7.0$  Hz), 3.90 (s, 3H). FABLRMS  $m/z$  400 ( $M^+$ ). FABHRMS  $m/z$  399.0249 ( $M-H$ , Calc'd 399.0247). Anal. Calc'd for  $\text{C}_{18}\text{H}_{12}\text{ClF}_3\text{O}_5$ : C, 53.95; H, 3.02; Cl, 8.85. Found: C, 53.78; H, 3.08; Cl, 8.98.

### EXAMPLE 10



#### 2-Trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

##### Step 1. Preparation of ethyl 2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

The ester was prepared from salicylaldehyde by a procedure similar to the method described in Example 1, Step 1: bp 107 °C 2mm.  $^1\text{H}$ NMR (acetone- $d_6$ /300 MHz) 7.89 (s, 1H), 7.52–7.38 (m, 2H), 7.09 (dt, 1  $J = 1.0$ , 7.7 Hz), 7.03 (d, 1H,  $J = 8.3$  Hz), 5.84 (q, 1H,  $J = 7.3$  Hz), 4.39–4.23 (m, 2H), 1.33 (t, 3H,  $J = 7.0$  Hz). FABLRMS  $m/z$  273 ( $M+H$ ). ESHRMS ( $m/z$  273.0720 ( $M+H$  Calc'd 273.0739).


Step 2. Preparation of 2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The acid was prepared from the ethyl ester (Step 1) by a procedure similar to the method described in Example 1,

- 5 Step 2: mp 152.2-153.3 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.89 (s, 1H), 7.39-7.49 (m, 2H), 7.11-7.01 (m, 2H), 5.81 (q H-F, 1H,  $J = 7.2$  Hz). FABHRMS  $m/z$  245.0422 (M+H, Calc'd 245.0426). Anal. Calc'd for  $\text{C}_{11}\text{H}_7\text{F}_3\text{O}_3$ : C, 54.11; H, 2.89. Found: C, 54.22; H, 2.97.

10

**EXAMPLE 11**



- 15 **6,8-Dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of 3,5-dichloro-4-methylsalicylaldehyde.

- 20 2,4-Dichloro-3-methylphenol (25.0 g, 141.2 mmol) was added to methanesulfonic acid (100 mL). With stirring, hexamethylenetetramine (HMTA) (39.8g, 282.4 mmol) and additional methanesulfonic acid (100 mL) was added portion-wise during which time the reaction began to froth and
- 25 exotherm. The resulting mixture was heated to 100 °C for 3 hours. The crude ocher colored suspension was cooled to 50 °C and poured over a mechanically stirred mixture of ice-water (2 L). A yellow precipitate was formed which was collected by vacuum filtration. This solid was purified by
- 30 flash chromatography (silica, hexanes-methylene chloride, 9:10) yielding the salicylaldehyde as a pale yellow powder



(6.17 g, 21%; mp 94.0-95.1 °C) of suitable purity to use without further purification.

5     Step 2. Preparation of ethyl 6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


A mixture of 3,5-dichloro-4-methylsalicylaldehyde (Step 1) (5.94 g, 29.0 mmol) and ethyl 4,4,4-trifluorocrotonate (7.67 g, 45.6 mmol) dissolved in anhydrous DMSO (10 mL) was treated with triethylamine (5.88 g, 58.1 mmol). The reaction  
10 was stirred at 85 °C for 49 hours then cooled in ice and filtered to give an orange solid. The solid was dissolved in ethyl acetate (100 mL), washed with 3 N HCl (2 x 50 mL), saturated NaHCO<sub>3</sub>, washed with brine, dried over MgSO<sub>4</sub>, and concentrated in vacuo to give a yellow solid (8.63 g, 84%):  
15 mp 117.1-119.5 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.63 (s, 1H), 7.17 (s, 1H), 5.80 (q, 1H, J = 6.6 Hz), 4.33 (m, 2H), 2.48 (s, 3H), 1.35 (t, 3H, J = 7.1 Hz).

20     Step 3. Preparation of 6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 2 (8.39 g 23.6 mmol) was dissolved in THF (30 mL) and ethanol (20 mL), treated with 2.5 N sodium hydroxide (20 mL, 50 mmol), and stirred at room temperature for 3.5 hours. The reaction mixture was  
25 concentrated in vacuo, acidified with 3 N HCl, filtered, and recrystallized from ethanol/ water to yield a yellow solid (6.0 g, 78%): mp 229.9-230.9 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.58 (s, 1H), 6.00 (q, 1H, J = 6.8 Hz), 2.50 (s, 3H). FABLRMS m/z 325 (M-H). FABHRMS m/z 324.9636 (M-H,  
30 Calc'd 324.9646). Anal. Calc'd for C<sub>17</sub>H<sub>7</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 44.07; H, 2.16; Cl, 21.68. Found: C, 44.06; H, 2.21; Cl, 21.74.

**EXAMPLE 12**

91




**7-(1,1-Dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5

Ethyl 7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 8, Step 2) was hydrolyzed to the carboxylic acid via a procedure similar to that described in Example 1, Step 2: mp 165.6-166.8 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.86 (s, 1H), 7.38 (d, 1H, *J* = 8.1 Hz), 7.15 (dd, 1H, *J* = 1.8 Hz, and *J* = 7.8 Hz), 7.05 (bs, 1H), 5.79 (q <sub>H-F</sub>, 1H, *J* = 7.2 Hz), 1.32 (s, 9H). FABHRMS *m/z* 301.1033 (M+H, Calc'd 301.1051). Anal. Calc'd for C<sub>15</sub>H<sub>15</sub>F<sub>3</sub>O<sub>3</sub>: C, 60.00; H, 5.04. Found: C, 59.80; H, 5.10.

15

**EXAMPLE 13**




20

**6-Bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5-Bromosalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 189.6-190.9 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.89 (s, 1H), 7.70 (d, 1H, *J* = 2.1 Hz), 7.55 (dd, 1H, *J* = 2.4 Hz, and *J* = 8.7 Hz), 7.02 (d, 1H, *J* = 8.7 Hz), 5.86 (q <sub>H-F</sub>, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 322.9519 (M+H, Calc'd

322.9531). Anal. Calc'd for  $C_{11}H_6BrF_3O_3$ : C, 40.90; H, 1.87; Br, 24.73. Found: C, 40.87; H, 1.92; Br, 24.80.

5


**EXAMPLE 14**

10

**8-Chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

2-Chlorophenol was converted to the title compound by a procedure similar to that described in Example 2: mp 224.5-225.6 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.91 (s, 1H), 7.49 (m, 2H), 7.11 (t, 1H,  $J = 7.8$  Hz), 5.96 (q  $H-F$ , 1H,  $J = 7.2$  Hz). FABHRMS  $m/z$  279.0027 ( $M+H$ , Calc'd 279.0036). Anal. Calc'd for  $C_{11}H_6ClF_3O_3$ : C, 47.42; H, 2.17. Found: C, 47.33; H, 2.17.

20

**EXAMPLE 15**

25


**8-Bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

2-Bromo-4-chlorosalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 227.8-228.9 °C.

<sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.65 (dd, 2H, *J* = 2.4 and *J* = 28.8 Hz), 6.00 (q H-F, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 356.9134 (M+H, Calc'd 356.9141). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>BrClF<sub>3</sub>O<sub>3</sub>: C, 36.96; H, 1.41. Found: C, 37.05; H, 1.33.

10

### EXAMPLE 16




### 15. 6-Trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

5-(Trifluoromethoxy)salicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 118.4-119.5 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.95 (s, 1H), 7.54 (d, 1H, *J* = 2.1 Hz), 7.39 (dd, 1H, *J* = 2.4 Hz, and *J* = 9.0 Hz), 7.02 (d, 1H, *J* = 9.0 Hz), 5.88 (q H-F, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 329.0228 (M+H, Calc'd 329.0249). Anal. Calc'd for C<sub>12</sub>H<sub>6</sub>F<sub>6</sub>O<sub>4</sub>: C, 43.92; H, 1.84. Found: C, 43.84; H, 1.87.

25

### EXAMPLE 17

30



**8-Fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5

3-Fluorosalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 197.7-210.1 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.30 (m, 2H),

10 7.11 (m 1H), 5.93 (q <sub>H-F</sub>, 1H, *J* = 7.2 Hz).


FABHRMS *m/z* 263.0341 (M+H, C<sub>11</sub>H<sub>6</sub>F<sub>4</sub>O<sub>3</sub> Calc'd

263.0331). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>F<sub>4</sub>O<sub>3</sub>: C, 50.40;

H, 2.31. Found: C, 50.48; H, 2.25.

15

**EXAMPLE 18**



**5,7-Dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

20


4,6-Dichlorosalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 190.1-191.2 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 8.01 (s, 1H), 7.3 (bs, 1H),

25 7.16 (bs, 1H), 5.94 (q <sub>H-F</sub>, 1H, *J* = 7.2 Hz).

FABHRMS *m/z* 312.9636 (M+H, Calc'd 312.9646).

Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 42.20; H, 1.61.

Found: C, 42.27; H, 1.56.

**EXAMPLE 19**

5


**7,8-Dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3,4-Dichlorophenol was converted to the title compound by a procedure similar to that described in Example 2: mp 219.5-220.9 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.51 (d, 1H, *J* = 8.4 Hz), 7.34 (d, 1H, *J* = 8.4 Hz), 6.02 (q *H*-F, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 318.9709 (M+Li, C<sub>11</sub>H<sub>5</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub> Calc'd 318.9728). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 42.20; H, 1.61. Found: C, 42.15; H, 1.68.

15

**EXAMPLE 20**

20


**7-Isopropoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

25

2,4-Dihydroxybenzaldehyde was alkylated to prepare 4-(1-methylethyloxy)salicylaldehyde. This salicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 161-163 °C. <sup>1</sup>H NMR


(CD<sub>3</sub>OD/300 MHz) 7.73 (s, 1H), 7.21 (d, 1H,  $J = 8.5$  Hz), 6.57 (dd, 1H,  $J = 8.5, 2.2$  Hz). FABHRMS  $m/z$  301.0688 ( $M-H^+$ , C<sub>11</sub>H<sub>12</sub>F<sub>3</sub>O<sub>4</sub> requires 301.0687). Anal. Calc'd for C<sub>11</sub>H<sub>13</sub>F<sub>3</sub>O<sub>4</sub>: C, 55.63; H, 4.34. Found: C, 55.72; H, 4.34.

5

**EXAMPLE 21**

10      **8-Phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

2-Phenylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 171.6-175.0 °C. <sup>1</sup>H NMR (acetone-  
15      d<sub>6</sub>/300 MHz) 7.95 (s, 1H), 7.46 (m, 7H), 7.18 (t, 1H,  $J = 7.5$  Hz), 5.81 (q  $H-F$ , 1H,  $J = 7.2$  Hz). FABHRMS  $m/z$  327.0816 ( $M+Li$ , Calc'd 327.0820). Anal. Calc'd for C<sub>17</sub>H<sub>11</sub>F<sub>3</sub>O<sub>3</sub>: C, 63.76; H, 3.46.  
20      Found: C, 63.52; H, 3.55.


**EXAMPLE 22**

25

7,8-Dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

2,3-Dimethylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 245.2-247.3 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.83 (s, 1H), 7.17 (d, 1H, *J* = 7.8 Hz), 6.89 (d, 1H, *J* = 7.8 Hz), 5.82 (q *H*-F, 1H, *J* = 7.2 Hz), 2.30 (s, 3H), 2.17 (s, 3H). Anal. Calc'd for C<sub>13</sub>H<sub>11</sub>F<sub>3</sub>O<sub>3</sub> + 1.56 % H<sub>2</sub>O: C, 56.46; H, 4.18. Found: C, 56.46; H, 4.15.

EXAMPLE 23




6,8-bis(1,1-Dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

3,5-Di-tert-butylsalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 171.6-175.0 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.65 (s, 1H), 7.34 (d, 1H, *J* = 2.4 Hz), 7.15 (d, 1H, *J* = 2.4 Hz), 6.02 (q *H*-F, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 363.1743 (M+Li, Calc'd 363.1759). Anal. Calc'd for C<sub>19</sub>H<sub>23</sub>BrF<sub>3</sub>O<sub>3</sub>: C, 64.03; H, 6.50. Found: C, 64.13; H, 6.49.

EXAMPLE 24





5 **6-Iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

Step 1: Preparation of 2-hydroxy-5-iodobenzyl alcohol.

A solution of 5-iodosalicylic acid (25.0 g, 94.6 mmol) in tetrahydrofuran (500 mL) was cooled to 0°C. With vigorous mixing, borane-methyl sulfide complex (15.1 ml of 10 M solution, 151.0 mmol) was added drop-wise over 0.25 hours. The solution was warmed to room temperature and then heated at reflux for 4 h. A white precipitate formed during the reflux. The solution was cooled to room temperature and 10% aqueous hydrochloric acid (100 mL) was added over 15 min and the solution stirred at room temperature for 2 h. The precipitate dissolved and the solvent was concentrated in vacuo to a volume of approximately 200 mL. The solution was poured into ethyl acetate (300 mL) and washed with water (2 x 200 mL), saturated sodium bicarbonate (2 x 200 mL), and saturated ammonium chloride (2 x 200 mL). The organic layer was dried over sodium sulfate and concentrated in vacuo. The 2-hydroxy-5-iodobenzyl alcohol was isolated as a white solid (21.3 g, 85.2 mmol) from hexanes. (90% yield): mp 105-110°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 8.21 (s, 1H), 7.30-7.33 (m, 2H), 6.57 (d, 1H, J = 8.3 Hz), 4.97 (bs, 1H), 4.62 (s, 2H). EIHMS m/z = 249.9492 (M<sup>+</sup>, Calc'd 249.9491).

Step 2: Preparation of 2-hydroxy-5-iodobenzaldehyde

To a stirred solution of 2-hydroxy-5-iodobenzyl alcohol (43.5 g, 174.0 mmol) in acetone (700 mL) was added 85% activated manganese(IV) oxide (5 micron, 50 g, 494.0 mmol) and the solution stirred at room temperature for 16 hours. The manganese oxide was removed by filtration through diatomaceous earth and the filtrate concentrated in vacuo. The product was purified by flash silica chromatography (0-

20% ethyl acetate in hexanes) The 2-hydroxy-5-iodobenzaldehyde was obtained as a greenish-yellow solid (24.3 g, 58%). A small amount of the 2-hydroxy-5-iodobenzaldehyde was recrystallized from methanol/water to afford an analytical sample and the remainder of the compound was used without further purification: mp 99-101 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 9.83 (s, 1H), 7.79 (d, 1H, J = 2.2 Hz), 7.77 (dd, 1H, J = 8.7 Hz, J = 2.2 Hz), 6.81 (d, 1H, J = 8.7 Hz). ESHRMS 246.9229 (M-H Calc'd 246.9256).

Step 3: Preparation of ethyl 6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate.


A mixture of 5-iodosalicylaldehyde (16.2 g, 65.3 mmol), ethyl 4,4,4-trifluorocrotonate (22.4 g, 133 mmol) and triethylamine (50 ml, 395 mmol) were combined, stirred at 70°C for 8 h and then heated at reflux for 48 h. The solution was poured into ethyl acetate (300 mL) and washed with 1N hydrochloric acid (3 x 200 mL). The aqueous layers were combined and extracted with ethyl acetate (1 x 100 mL). The combined ethyl acetate extracts were washed with saturated ammonium chloride (2 x 200 mL), dried over magnesium sulfate and concentrated *in vacuo* yielding a dark red oil. This oil was purified by flash chromatography using ethyl acetate-hexanes (3:7) yielding a red oil. Crystallization of this oil from hexanes yielded the title compound as light red crystals (8.3 g, 31%): mp 105-106 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.63 (s, 1H), 7.58 (dd, 2H, J = 8.6, J = 2.1 Hz), 7.54 (d, 1H, J = 2.1 Hz), 6.77 (d, 1H, J = 8.6 Hz), 5.70 (q, 1H, J = 6.7 Hz), 4.20-4.38 (m 2H), 1.35 (t, 3H, J = 7.2 Hz). ESHRMS 415.9926 (M+NH<sub>4</sub><sup>+</sup> Calc'd 396.9746)

Step 4: Preparation of 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

Hydrolysis of the ester (Step 3), using a procedure similar to Example 1, Step 2, yielded the carboxylic acid: mp 168-170 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.57 (s, 1H), 7.70 (d, 1H, J = 2.2 Hz), 7.64 (dd, 1H, J = 8.5, 2.2 Hz), 6.79

(d, 1H,  $J = 8.5$  Hz) 5.78 (q, 1H,  $J = 7.0$  Hz). ESHRMS  $m/z$  368.9222 (Calc'd for M-H 368.9235). Anal. Calc'd for  $C_{11}H_6F_3IO_3$ : C, 35.70; H, 1.63. Found C, 35.67; H, 1.63.

5


**EXAMPLE 25**

10

**7-(1-Methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3-(1-Methylethyl)phenol was converted to the title compound by a procedure similar to that described in Example 2: mp 158.3-159.7 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.86 (s, 1H), 7.37 (d, 1H,  $J = 7.8$  Hz), 7.00 (d, 1H,  $J = 7.8$  Hz), 6.91 (s, 1H), 5.78 (q, 1H,  $J = 6.9$  Hz), 2.93 (m, 1H), 1.24 (d, 6H,  $J = 6.9$  Hz). FABLRMS  $m/z$  287 (M+H). Anal. Calc'd for  $C_{14}H_{13}F_3O_3$ : C, 58.74; H, 4.58. Found: C, 57.37; H, 4.49.

20


**EXAMPLE 26**

25

**7-Phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3-Phenylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 209.4-211.7 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.74 (m, 2H), 7.47 (m, 5H), 7.33 (s, 1H), 5.86 (q, 1H, *J* = 7.2 Hz). FABLRMS *m/z* 321 (M+H). Anal. Calc'd for C<sub>17</sub>H<sub>11</sub>F<sub>3</sub>O<sub>3</sub>: C, 63.76; H, 3.46. Found: C, 64.17; H, 3.61.

10


**EXAMPLE 27**

15

**6-Chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

4-Chloro-3-ethylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 170.7-172.1 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.78 (s, 1H), 7.26 (s, 1H), 6.90 (s, 1H), 5.67 (q, 1H, *J* = 6.9 Hz), 2.73 (q, 2H, *J* = 7.8 Hz), 1.24 (t, 3H, *J* = 7.8 Hz). FABLRMS *m/z* 307 (M+H). Anal. Calc'd for C<sub>13</sub>H<sub>10</sub>F<sub>3</sub>O<sub>3</sub>: C, 50.92; H, 3.29. Found: C, 51.00; H, 3.33.

25


**EXAMPLE 28**

8-Ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

5        2-Ethylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 185.4-186.8 °C. <sup>1</sup>H NMR (acetone-  
d<sub>6</sub>/300 MHz) 7.85 (s, 1H), 7.28 (d, 2H, *J* = 7.5 Hz), 7.00 (t, 1H, *J* = 7.5 Hz), 5.84 (q, 1H, *J* =  
10    7.2 Hz), 2.65 (m, 2H), 1.18 (t, 3H, *J* = 7.5 Hz). FABLRMS *m/z* 273 (M+H). Anal. Calc'd for C<sub>13</sub>H<sub>11</sub>F<sub>3</sub>O<sub>3</sub>: C, 57.36; H, 4.07. Found: C, 57.15; H, 4.11.

15

**EXAMPLE 29**




20

6-Chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

8-Ethyl-2-(trifluoromethyl)2H-1-benzopyran-3-carboxylic acid (Example 28) (0.68 g, 2.5 mmol) was dissolved in trimethylphosphate (5 mL) and was  
25    treated with sulfuryl chloride (0.35 g, 2.62 mmol) at 0 °C. After stirring at 0 °C for 45 minutes and 1 hour at room temperature, the reaction was diluted with cold water (15 mL). The resulting oily mixture was extracted with hexanes-ethyl  
30    acetate. The organic phase was washed with brine, dried, and concentrated *in vacuo* yielding the title compound as a solid (0.9 g, 117 %):

mp 197.2-199.1 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz)  
7.86 (s, 1H), 7.38 (d, 1H,  $J = 2.7$  Hz), 7.30 (d,  
1H,  $J = 2.4$  Hz), 5.88 (q, 1H,  $J = 7.2$  Hz), 2.65  
(m, 2H), 1.19 (t, 3H,  $J = 7.5$  Hz). FABLRMS  $m/z$   
5 307 (M+H). Anal. Calc'd for  $\text{C}_{13}\text{H}_{10}\text{ClF}_3\text{O}_3$ : C,  
50.92; H, 3.29. Found: C, 51.00; H, 3.23.

### EXAMPLE 30




10

#### 6-Chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

15 7-Phenyl-2-(trifluoromethyl)-2H-1-benzopyran-  
3-carboxylic acid (Example 26) was converted to  
the title compound by a procedure similar to that  
described in Example 29: mp 185.3-187.8 °C.  $^1\text{H}$   
NMR (acetone- $d_6$ /300 MHz) 7.94 (s, 1H), 7.68 (s,  
20 1H), 7.47 (m, 5H), 7.06 (s, 1H), 5.87 (q, 1H,  $J =$   
6.9 Hz). FABLRMS  $m/z$  355 (M+H). Anal. Calc'd  
for  $\text{C}_{17}\text{H}_{10}\text{ClF}_3\text{O}_3$ : C, 57.56; H, 2.84. Found: C,  
58.27; H, 3.11.

25

### EXAMPLE 31




**6,7-Dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3,4-Dichlorophenol was converted to the title compound  
5 by a procedure similar to that described in Example 2: mp  
196.1-198.3 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.90 (s, 1H),  
7.74 (s, 1H), 7.30 (s, 1H), 5.88 (q, 1H, J = 6.9 Hz).  
FABLRMS m/z 314 (M+H). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C,  
42.20; H, 1.61. Found: C, 42.31; H, 1.65.

10


**EXAMPLE 32**



15 **6,8-Dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3,5-Dichlorosalicylaldehyde was converted to the title  
compound by a procedure similar to that described in Example  
20 11, Steps 2 & 3: mp 212.8-216.8 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz)  
7.77 (s, 1H), 7.41 (d, 1H, J = 2.4 Hz), 7.18 (d, 1H, J = 2.2  
Hz), 5.82 (q, 1H, J = 6.7 Hz). FABLRMS m/z 311 (M-H).  
FABHRMS m/z 312.9644 (M+H, Calc'd 312.9646). Anal. Calc'd  
for C<sub>11</sub>H<sub>5</sub>F<sub>3</sub>Cl<sub>2</sub>O<sub>3</sub>: C, 42.20; H, 1.61. Found: C, 42.50; H,  
25 1.71.

**EXAMPLE 33**




**6,8-Dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5

3,5-Dibromosalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 225-226 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.76 (s, 1H), 7.74 (d, 1H, *J* = 2.2 Hz), 7.55 (d, 1H, *J* = 2.2 Hz), 5.91 (q H-F, 1H, *J* = 7.2 Hz). FABHRMS *m/z* 400.8648 (M+H<sup>+</sup>, Calc'd 400.8636). Anal. Calcd for C<sub>11</sub>H<sub>5</sub>Br<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 32.87; H, 1.25. Found: C, 33.47; H, 1.38.

**EXAMPLE 34**

15



**6,8-Dimethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

20

4,6-Dimethoxysalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 215-217 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.95 (s, 1H), 6.18-6.20 (m, 2H), 5.65 (q H-F, 1H, *J* = 7.2 Hz), 3.87 (s, 1H), 3.81 (s, 1H). FABHRMS *m/z* 303.0497 (M-H<sup>+</sup>, Calc'd


25



303.0380). Anal. Calc'd for  $C_{13}H_{11}F_3O_5$ : C, 51.33; H, 3.64. Found: C, 51.19; H, 3.71.

### EXAMPLE 35

5



#### Ethyl 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate

10 Step 1. Preparation of ethyl 6-nitro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


A mixture of 5-nitrosalicylaldehyde (4.80g, 28.7 mmol) and ethyl 4,4,4-trifluorocrotonate (6.6 g, 39.4 mol) in anhydrous DMF was warmed to 60 °C and treated  
15 with anhydrous  $K_2CO_3$  (3.90 g, 28.9 mol). The solution was maintained at 60 °C for 20 hours, cooled to room temperature, diluted with water, and extracted with ethyl acetate. The organic extracts were washed with brine, dried over anhydrous  $MgSO_4$ , filtered and  
20 concentrated in vacuo to afford an oil. The oil was dissolved in diethyl ether (5 mL). Hexanes was added until the solution became cloudy. Upon standing at room temperature overnight the ester was obtained as yellow crystals (0.856 g, 7% yield). This material was of  
25 sufficient purity to be used in subsequent steps without further purification.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 8.15-8.19 (m, 2H), 7.74 (s, 1H), 7.09 (d, 1H,  $J = 8.9$  Hz), 5.81 (q, 1H,  $J = 5.8$  Hz), 4.29-4.39 (m, 2H), 1.35 (t, 3H,  $J = 6.0$  Hz),

30

Step 2. Preparation of ethyl 6-amino-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

The ester (Step 1) (0.345 g, 1.08 mmol) was stirred in ethanol (10.0 mL) with 10% palladium on charcoal (15 mg) with hydrogen at 1 atmosphere for 1 hour. The catalyst was removed by filtration and the solvent removed *in vacuo* to afford the title compound as an orange-yellow solid (0.298 g, 95 %): mp 111-115 °C. ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 7.69 (s, 1H), 6.69-6.74 (m, 3H), 5.65 (q H-F, 1H,  $J = 7.2\text{ Hz}$ ), 4.26-4.37 (m, 2H), 1.34 (t, 3H,  $J = 7\text{ Hz}$ ). FABHRMS  $m/z$  288.0860 ( $\text{M}+\text{H}^+$ ,  $\text{C}_{13}\text{H}_{13}\text{F}_3\text{NO}_3$  requires 288.0847). Anal. Calc'd for  $\text{C}_{13}\text{H}_{12}\text{F}_3\text{NO}_3$ : C, 54.36; H, 4.21; N, 4.88. Found: C, 54.46; H, 4.27; N, 4.83.


### EXAMPLE 36



#### 6-Amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

Ethyl 6-amino-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 35, Step 2) was hydrolyzed to the carboxylic acid (title compound) by a procedure similar to that described in Example 1, Step 2: mp 126-133 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 6.81-6.90 (m, 3H), 5.66 (q H-F, 1H,  $J=7.2\text{ Hz}$ ). FABHRMS  $m/z$  260.0535 ( $\text{M}+\text{H}^+$ ,  $\text{C}_{11}\text{H}_9\text{F}_3\text{NO}_5$  requires 260.0534).

### EXAMPLE 37




**6-Nitro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5 Ethyl 6-nitro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 35, Step 1) was hydrolyzed to the carboxylic acid (title compound) by a procedure similar to that described in Example 1, Step 2: mp 187-189 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 8.34 (d, 1H, *J* = 2.6 Hz), 8.27 (dd, 1H, *J* =  
10 8.7, 2.6 Hz), 7.90 (s, 1H), 7.09 (s, 1H, *J* = 8.7 Hz), 5.81 (q H-F, 1H, *J* = 7.2 Hz). EIHRMS *m/z* 289.0177 (Calc'd 289.0198). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>F<sub>3</sub>NO<sub>5</sub>: C, 45.69; H, 2.09; N 4.84 Found: C, 45.71; H, 2.08; N 4.75.

15

**EXAMPLE 38**




**6-Chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

20

4-Chloro-2-methylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 231.9-233.2 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.76 (s, 1H), 7.19 (d, 1H, *J* = 1.8 Hz), 7.09 (d, 1H, *J* = 2.4 Hz), 5.72  
25 (q, 1H, *J* = 6.9 Hz), 2.24 (s, 3H). <sup>19</sup>F NMR (CDCl<sub>3</sub>/282 MHz) -79.2 (d, *J* = 6.5 Hz). FABLRMS *m/z* 299 (M+Li). FABHRMS *m/z*

293.0196 (M+H, Calc'd 293.0192). Anal. Calc'd for  $C_{12}H_8ClF_3O_3$ : C, 49.25; H, 2.76. Found: C, 49.37; H, 2.86.


### EXAMPLE 39



#### 8-Chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

2-Chloro-4-methylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 226.4-227.4 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 7.79 (s, 1H), 7.23 (d, 1H,  $J = 1.4$  Hz), 6.97 (d, 1H,  $J = 1.4$  Hz), 5.77 (q, 1H,  $J = 6.8$  Hz), 2.29 (s, 3H).  $^{19}F$  NMR ( $CDCl_3$ /282 MHz) -79.1 (d,  $J = 7.3$  Hz). FAB/MS  $m/z$  291 (M-H). EI/MS  $m/z$  292.0118 (M+,  $C_{12}H_8ClF_3O_3$  Calc'd 292.0114).

### EXAMPLE 40



#### 8-Chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid


2-Chloro-4-methoxyphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 204.5-206.9 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 7.78 (s, 1H),

6.98 (d, 1H,  $J = 2.8$  Hz), 6.71 (d, 1H,  $J = 2.8$  Hz), 5.74 (q, 1H,  $J = 6.9$  Hz), 3.79 (s, 3H). FABLRMS  $m/z$  326 ( $M+NH_4$ ).

EIHRMS  $m/z$  308.0053 ( $M+$  Calc'd 308.0063). Anal. Calc'd for  $C_{12}H_8ClF_3O_4$ : C, 46.70; H, 2.61. Found: C, 46.60; H, 2.68.

5

### EXAMPLE 41




### 10 6,8-Difluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

2,4-Difluorophenol was converted to the title compound by a procedure similar to that described in Example 2: mp

15 207-211 °C.  $^1H$  NMR ( $CDCl_3$ ) 7.63 (s, 1H), 6.89-6.72 (m, 2H), 5.69 (q, 1H,  $J = 6.7$  Hz). Anal. Calc'd for  $C_{11}H_5O_3F_5$ : C, 47.16; H, 1.80. Found: C, 47.28; H, 1.87.

### EXAMPLE 42

20




### 25 6-Bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

4-Bromo-2-chlorophenol was converted to the title compound by a procedure similar to that described in Example 2: mp 220.7-221.7 °C.  $^1H$  NMR ( $CDCl_3$ ) 7.58 (s, 1H), 7.44 (d, 1H,  $J = 2.2$  Hz), 7.22 (d, 1H,  $J = 2.2$  Hz), 5.74 (q, 1H,  $J = 6.8$

Hz). Anal. Calc'd for  $C_{11}H_5O_3F_3BrCl$ : C, 36.96; H, 1.41.  
Found: C, 37.03; H, 1.44.

### EXAMPLE 43

5




#### 8-Bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

10

2-Bromo-4-fluorophenol was converted to the title compound by a procedure similar to that described in Example 2: mp  $>300^{\circ}C$ .  $^1H$  NMR ( $CDCl_3$ ) 7.58 (s, 1H), 7.22 (dd, 1H,  $J$  = 6.3, 3 Hz), 6.88 (dd, 1H,  $J$  = 6.1, 3.1 Hz), 5.72 (q, 1H,  $J$  = 6.7 Hz). Anal. Calc'd for  $C_{11}H_5O_3F_4Br$ : C, 38.74; H, 1.48.  
Found: C, 38.82; H, 1.56.

### EXAMPLE 44

20




#### 8-Bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

25

2-Bromo-4-methylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp  $237-238^{\circ}C$ .  $^1H$  NMR ( $CDCl_3$ ) 7.59 (s, 1H), 7.27 (m, 1H), 6.91 (d, 1H,  $J$  = 1.4 Hz), 5.69 (q, 1H,  $J$  = 6.9 Hz), 2.20 (s, 3H). Anal. Calc'd for  $C_{12}H_8O_3F_3Br$ : C, 42.76; H, 2.39.  
Found: C, 43.34; H, 2.56.

30


**EXAMPLE 45**

5

**8-Bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

2-Bromo-5-fluorophenol was converted to the title  
10 compound by a procedure similar to that described in Example  
2: mp 221.7-223.3 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) 7.81 (s, 1H), 7.38  
(dd, 1H, *J* = 7.3, 5.8 Hz), 6.58 (t, 1H, *J* = 8.9 Hz), 5.71 (q,  
1H, *J* = 6.7 Hz). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>O<sub>3</sub>F<sub>4</sub>Br: C, 38.74; H,  
1.48. Found: C, 38.70; H, 1.54.

15

**EXAMPLE 46**

20


**6-Chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

4-Chloro-2-fluorophenol was converted to the title  
compound by a procedure similar to that described in Example  
25 2: mp 190.8-193.0 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.77 (s, 1H),  
7.19 (d of d, 1H, *J* = 2.2 and 9.7 Hz), 7.07 (t, 1H, *J* = 1.8  
Hz), 5.76 (q, 1H, *J* = 6.7 Hz). FABLRMS *m/z* 295 (M-H).

EIHRMS  $m/z$  295.9876 ( $M^+$  Calc'd 295.9863). Anal. Calc'd for  $C_{11}H_5ClF_4O_3$ : C, 44.54; H, 1.70. Found: C, 44.36; H, 1.85.

#### EXAMPLE 47

5



#### 6-Bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid


10

4-Bromo-2-methoxysalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp dec. at 244 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.71 (s, 1H), 7.18 (d, 1H,  $J = 2.2$  Hz), 7.11 (d, 1H,  $J = 2.2$  Hz), 5.77 (q  $H-F$ , 1H,  $J=7.2$  Hz), 3.84 (s, 3H). FABLRMS  $m/z$  351 (m-H). Anal. Calc'd for  $C_{12}H_8BrF_3O_5$ : C, 40.82; H, 2.28. Found: C, 40.83; H, 2.30.

15

#### EXAMPLE 48

20



#### 7-(N,N-Diethylamino)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

25

4-(N,N-Diethylamino)salicylaldehyde was converted to the title compound by a procedure similar to that described




in Example 1: mp 214.4-215.4 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 7.67 (s, 1H), 7.06 (d, 1H,  $J = 8.6\text{ Hz}$ ), 6.34 (dd, 1H,  $J = 8.6, 2.3\text{ Hz}$ ), 5.60 (q  $\text{H-F}$ , 1H,  $J=7.2\text{ Hz}$ ), 3.38 (q, 4H,  $J = 7.1\text{ Hz}$ ), 1.16 (t, 6H,  $J = 7.1\text{ Hz}$ ). ESLRMS  $m/z$  316 ( $\text{M} + \text{H}$ ).

5 FABHRMS  $m/z$  316.1145 ( $\text{M} + \text{H}^+$ , Calc'd 316.1161). Anal. Calc'd for  $\text{C}_{15}\text{H}_{16}\text{F}_3\text{NO}_3$ : C, 57.14; H, 5.11; N, 4.44. Found: C, 57.14; H, 5.08; N, 4.44.

### EXAMPLE 49

10



#### 6-[[[(Phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

15

##### Step 1. Preparation of ethyl 6-chlorosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate.

Chlorosulfonic acid (50.0 mL) was cooled to 15 °C and ethyl 2-trifluoromethyl-2H-1-benzopyran-3-carboxylate (Example 10, Step 2) (6.21 g, 22.83 mmol) was added. After stirring at -15 °C for 1 hour, the solution was warmed to room temperature and stirred for 16 hours. The solution was added dropwise onto ice (500 mL) with vigorous stirring and extracted with diethyl ether (2 x 250 mL). The ether layers were combined, washed with water (2 x 250 mL), saturated sodium bicarbonate (2 x 250 mL), and brine (2 x 250 mL). Hexanes (50 mL) were added and the solution was dried over sodium sulfate. The solvent was removed in vacuo to afford the ester as a yellow solid (7.41 g, 87%): mp 97.2-98.4 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz) 7.97 (dd, 1H,  $J = 8.6, 2.2\text{ Hz}$ ), 7.92 (d, 1H,  $J = 2.2\text{ Hz}$ ), 7.73 (s, 1H), 7.17

(d, 1H,  $J = 2.2$  Hz), 5.82 (q H-F, 1H,  $J=7.2$  Hz), 4.28-4.39 (m, 2H), 1.35 (t, 3H,  $J = 7.0$  Hz). FABLRMS  $m/z$  376 ( $M+Li^+$ ).


Step 2. Preparation of ethyl 6-

5 [[ (phenylmethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate.

The sulfonyl chloride from Step 1 (451.0 mg, 1.22 mmol) and benzylamine (600 mg, 5.62 mmol) were mixed in diethyl ether (25 mL) for 1 hour at room  
10 temperature. The solution was washed with 1N HCl (2 x 25 mL), saturated sodium bicarbonate (2 x 25 mL), and brine (2 x 25 mL). The solution was dried over sodium sulfate and dried in vacuo. The aminosulfonyl was obtained by crystallization from  
15 hexanes (431 mg, 84%): mp 128.2-131.9 °C.  $^1H$  NMR ( $CDCl_3$ , 300 MHz) 7.76 (dd, 1H,  $J = 8.4, 2.2$  Hz), 7.70 (d, 1H,  $J = 2.2$  Hz), 7.67 (s, 1H), 7.12-7.30 (m, 5H), 7.05 (d, 1H,  $J = 8.4$  Hz), 5.78 (q H-F, 1H,  $J=7.2$  Hz), 4.68 (m, 2H), 4.19-4.32 (m, 2H), 1.37  
20 (t, 3H,  $J = 7.0$  Hz). FABLRMS  $m/z$  442 ( $M + H^+$ ).  
• FABHRMS  $m/z$  442.0936 ( $M + H^+$ ,  $C_{20}H_{19}F_3NO_5S$  Calc'd 442.0916).

25 Step 3. Preparation of 6-[[ (phenylmethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The acid was converted from the ester (step 2) via the method similar to that described in Example 1, step 2: mp 223.3-224.4 °C  $^1H$  NMR ( $CD_3OD/300$  MHz)  
30 7.31-7.80 (m, 3H), 7.15-7.25 (m, 5H), 7.06 (d, 1H,  $J = 8.3$  Hz), 5.87 (q H-F, 1H,  $J=7.2$  Hz), 4.11 (s, 2H).  
FABLRMS  $m/z$  420 ( $M + Li^+$ ). FABHRMS  $m/z$  414.0589 ( $M + H^+$  Calc'd 414.0623). Anal. Calc'd for  $C_{18}H_{14}F_3NO_5S$ : C, 52.30; H, 3.41; N, 3.39. Found: C, 5.16; H, 3.44; N, 3.32.

**EXAMPLE 50**


5

**6-[(Dimethylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar  
10 procedure to that described in Example 49: mp 201.2-  
202.5 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.90 (s, 1H), 7.82  
(d, 1H, *J* = 2.2 Hz), 7.76 (dd, 1H, *J* = 8.6, 2.2 Hz),  
7.19 (d, 1H, *J* = 8.6 Hz), 5.91 (q H-F, 1H, *J*=7.2  
Hz), 2.70 (s, 6H). FABLRMS *m/z* 352 (M + H<sup>+</sup>).  
15 FABHRMS *m/z* 352.0466 (M + H<sup>+</sup> Calc'd 352.0467).  
Anal. Calc'd for C<sub>13</sub>H<sub>12</sub>F<sub>3</sub>NO<sub>5</sub>S: C, 44.45; H, 3.44; N,  
3.99. Found: C, 4.42; H, 3.45; N, 3.96.

**EXAMPLE 51**

20

**6-Aminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**


25

The title compound was prepared by a similar procedure  
to that described in Example 49: mp 187.9-189.8 °C. <sup>1</sup>H NMR  
(CD<sub>3</sub>OD/300 MHz) 7.58-7.88 (m, 3H), 7.12 (d, *J* = 8.3 Hz),  
5.87 (q H-F, 1H, *J*=7.2 Hz). FABLRMS *m/z* 324 (M + H<sup>+</sup>).

FABHRMS  $m/z$  324.0156 ( $M + H^+$  Calc'd 324.0154). Anal. Calc'd for  $C_{11}H_8F_3NO_5S \cdot 0.74 H_2O$ : C, 39.26; H, 2.84; N, 4.16.

Found: C, 39.33; H, 2.82; N, 4.11.

5


**EXAMPLE 52**

10

**6-(Methylamino)sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 207.6-208.6 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.83-7.97 (m, 3H), 7.19 (d, 1H,  $J = 8.5$  Hz), 5.91 (q  $H-F$ , 1H,  $J=7.2$  Hz), 3.11 (s, 3H). FABLRMS  $m/z$  338 ( $M + H^+$ ). FABHRMS  $m/z$  338.0331 ( $M + H^+$  Calc'd 338.0310). Anal. Calc'd for  $C_{12}H_{11}F_3NO_5S$ : C, 42.73; H, 2.99; N, 4.15. Found: C, 42.91; H, 3.06; N, 4.04.

20

**EXAMPLE 53**


25

**6-[(4-Morpholino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 215.2-219.3 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.88 (s, 1H), 7.81 (d, 1H,  $J = 2.2$  Hz), 7.74

(dd, 1H,  $J = 8.6, 2.2$  Hz), 5.90 (q H-F, 1H,  $J=7.2$  Hz), 3.54-3.70 (m, 4H), 2.94-2.97 (m, 4H). FABLRMS  $m/z$  394 ( $M + H^+$ ). FABHRMS 394.0567 ( $M + H^+$ ,  $C_{15}H_{15}F_3NO_6S$  Calc'd 394.0572).

5


**EXAMPLE 54**

10

**6-[(1,1-Dimethylethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 229.3-233.5 °C  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.82-7.87 (m, 3H), 7.12 (d, 1H,  $J = 8.6$  Hz), 5.87 (q H-F, 1H,  $J=7.2$  Hz), 1.18 (s, 9H). FABLRMS  $m/z$  380 ( $M + H^+$ ). Anal. Calc'd for  $C_{15}H_{16}F_3NO_5S$ : C, 47.49; H, 4.25; N, 3.69. Found: C, 47.95; H, 4.48; N, 3.55.

20

**EXAMPLE 55**


25

**6-[(2-Methylpropyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 190.6-192.4 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.77-7.84 (m, 3H), 7.13 (d, 1H,  $J = 8.4$  Hz),

5.86 (q H-F, 1H,  $J=7.2$  Hz), 2.64 (d, 2H,  $J = 6.8$  Hz), 1.66 (sept, 1H,  $J = 6.6$  Hz), 0.84 (d, 6H,  $J = 6.6$  Hz). FABLRMS  $m/z$  380 ( $M + H^+$ ). Anal. Calc'd for  $C_{15}H_{16}F_3NO_5S$ : C, 47.49; H, 4.25; N, 3.69. Found: C, 47.61; H, 3.34; N, 3.55.

5

**EXAMPLE 56**

10 **6-Methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of 6-chlorosulfonyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

15 To chlorosulfonic acid (50.0 mL) chilled to  $-15^{\circ}\text{C}$  was added 2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 10) (4.0 g, 16.7 mmol). After stirring at  $-15^{\circ}\text{C}$  for 1 hour, the solution was warmed to room temperature and stirred for 16 hours. The resulting solution was added

20 dropwise over ice (100 mL) with two diethyl ether (2 x 75 mL) extractions. The diethyl ether layers were combined, washed with water (2 x 75 mL), and brine (2 x 75 mL), dried over sodium sulfate and concentrated *in vacuo*. The resulting solids were triturated with hexane-ethyl acetate (9:1, 100


25 mL). The 6-chlorosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid was isolated as a white solid: mp 169-174.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 8.18 (d, 1H,  $J = 2.7$  Hz), 8.06 (dd, 1H,  $J = 8.7, 2.7$  Hz), 7.93 (s, 1H), 7.28 (d, 1H,  $J = 8.7$  Hz), 6.00 (q, 1H,  $J = 6.6$  Hz). EIHRMS  $m/z$

30 324.9977 ( $M^+$ , Calcd 324.9994).

Step 2. Preparation of 6-methylsulfonyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

A slurry of the chlorosulfonyl intermediate (Example 49, Step 1) (493 mg, 1.44 mmol), sodium bicarbonate (362 mg, 4.32 mmol), and sodium bisulfite (181 mg, 1.44 mmol) in water (1.5 mL) was heated to 60 °C for 1.5 h, followed by the addition of bromoacetic acid (212 mg, 1.55 mmol). The resulting suspension was heated to reflux, followed by the addition of sodium hydroxide solution (50% NaOH soln., 0.10 mL) and water (3.0 mL). The solution was refluxed for 8 hours, cooled to room temperature, and acidified to pH 1 with 1N aqueous hydrochloric acid. The solution was extracted with ethyl acetate (2 x 25 mL). The combined ethyl acetate layers were washed with 1N aqueous hydrochloric acid (2 x 25 mL), water (2 x 25 mL), and brine (2 x 25 mL), dried over sodium sulfate, filtered and concentrated in vacuo yielding the title compound as an off white solid. (231 mg, 50 % yield): mp 208.3-212.4°C. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz) 7.97 (d, 1H, 2.2 Hz), 7.91 (1H, dd, J = 8.7, 2.2 Hz), 7.19 (d, 1H, J = 8.7 Hz), 5.91 (q H-F, 1H, J=7.2 Hz), 3.11 (s, 1H) HRLRMS m/z 321 (M - H) FABLRMS m/z 321 (M - H). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>5</sub>S\*0.61 H<sub>2</sub>O: C, 43.26; H, 3.09. Found: C, 43.24; H, 3.09.

### EXAMPLE 57




**8-Chloro-6-[[[(phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 167.0-173.8 °C <sup>1</sup>H NMR

(CD<sub>3</sub>OD/300 MHz) 7.78 (s, 1H), 7.72 (d, 1H,  $J = 2.0$  Hz), 7.64 (d, 1H,  $J = 2.0$  Hz), 7.44 (s, 1H), 7.15-7.23 (m, 5H), 6.01 (q H-F, 1H,  $J = 7.2$  Hz), 4.08-4.15 (m, 2H). FABLRMS  $m/z$  454 (M + Li<sup>+</sup>). Anal. Calc'd for C<sub>16</sub>H<sub>13</sub>ClF<sub>3</sub>NO<sub>5</sub>S: C, 48.28; H, 2.93; N, 3.13. Found: C, xx; H, xx; N, xx.

### EXAMPLE 58




10           **6-N,N-Diethylaminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

The title compound was prepared by a similar procedure to that described in Example 49: mp 238-240 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.88 (s, 1H), 7.85 (d, 1H,  $J = 2.2$  Hz), 7.79 (dd, 1H,  $J = 8.5, 2.2$  Hz), 7.14 (d, 1H,  $J = 8.5$  Hz), 5.88 (q H-F, 1H,  $J = 7.2$  Hz), 3.24 (q, 2H,  $J = 7.3$  Hz), 1.11 (t, 3H,  $J = 7.3$  Hz). FABHRMS  $m/z$  380.0763 (M+H<sup>+</sup>, Calc'd 380.0780). Anal. Calc'd for C<sub>15</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>4</sub>S: C, 47.49; H, 4.25; H, 3.69.

20   Found: C, 47.62; H, 4.30; N, 3.72.

### EXAMPLE 59





**6-Phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**


5    Step 1. Preparation of ethyl 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate.

2-Trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (Example 10) (1.32 g, 4.85 mmol) was cooled to 0 °C in dichloromethane (50 mL).  
10 Aluminum chloride (2.58 g, 19.5 mmol) was added and a dark red solution resulted. A solution of phenylacetyl chloride (1.8 g, 12.1 mmol) in dichloromethane (10.0 mL) was added dropwise over 40 minutes. The solution was warmed to room  
15 temperature and stirred for 16 hours. The solution was poured onto ice (200 mL) and extracted with diethyl ether ( 2 x 100 mL). The diethyl ether layers were combined, extracted with water (1 x 100 mL), 1 N HCl (2 x 100 mL), and  
20 saturated sodium bicarbonate (3 x 100 mL). Hexanes (20 mL) were added and the solution was extracted with brine (1 x 100 mL). The solution was dried over sodium sulfate and solvent was removed in vacuo. The crude ester was purified by  
25 flash chromatography over silica gel (with ethyl acetate as eluant) to afford the ester that was crystallized from diethyl ether/hexanes (830 mg, 44%): mp 136.2-138.0 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.98 (dd, 2H, J = 8.4, 2.0 Hz), 7.90 (d, 1H, J =  
30 2.0 Hz), 7.29 (s, 1H), 7.22 - 7.38 (m, 5H), 7.02 (d, 1H, J = 8.4 Hz), 5.75 (q <sub>H-F</sub>, 1H, J=7.2 Hz), 4.25 - 4.40 (m, 2H), 4.21 (s, 2H), 1.34 (t, 3H, J = 7.0 Hz). FABLRMS m/z 391(M + H<sup>+</sup>).

35    Step 2. Preparation of 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The acid was converted from the ester (Step 1) via a method similar to that described in Example 1, step 2: mp 159.0-164.0 °C  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 8.04-8.16 (m, 3H), 7.87 (s, 1H), 7.05-7.30 (m, 5H), 5.86 (q  $\text{H-F}$ , 1H,  $J=7.2\text{ Hz}$ ), 4.31 (s, 2H). FABLRMS  $m/z$  363 ( $\text{M} + \text{H}^+$ ). Anal. Calc'd for  $\text{C}_{19}\text{H}_{13}\text{F}_3\text{O}_4 \cdot 0.29\text{ H}_2\text{O}$ : C, 62.08; H, 3.73. Found: C, 62.04; H, 4.03.

### EXAMPLE 60




#### 6-(2,2-Dimethylpropylcarbonyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

15

The title compound was prepared by a similar procedure to that described in Example 59: mp 198-200 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300\text{ MHz}$ ) 7.98-8.06 (m, 2H), 7.88 (s, 1H), 7.07 (d, 1H,  $J = 8.9\text{ Hz}$ ), 5.86 (q  $\text{H-F}$ , 1H,  $J = 7.2\text{ Hz}$ ), 2.88 (s, 2H), 1.05 (s, 9H). FABHRMS  $m/z$  343.1175 ( $\text{M} + \text{H}^+$ ,  $\text{C}_{17}\text{H}_{18}\text{F}_3\text{O}_4$  requires 343.1157). Anal. Calc'd for  $\text{C}_{17}\text{H}_{17}\text{F}_3\text{O}_4$ : C, 59.65; H, 5.01. Found: C, 59.70; H, 4.97.

### EXAMPLE 61

25



**6,8-Dichloro-7-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

Step 1: Preparation of ethyl 7-methoxy-2-trifluoromethyl-benzopyran-2H-3-carboxylate.

4-Methoxysalicylaldehyde (2.38 g, 15.64 mmol),  
K<sub>2</sub>CO<sub>3</sub> (2.16 g, 15.64 mmol) and ethyl 4,4,4-trifluorocrotonate (2.8 mL, 3.16 g, 18.77 mmol) were dissolved in DMF (10 mL). The reaction was stirred at room temperature for 24 hours, diluted with water and extracted with Et<sub>2</sub>O. The combined Et<sub>2</sub>O phases were washed with water, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding an oil. Trituration with hexanes induced crystallization. Collection of the solid by vacuum filtration yielded the ester as a light brown crystalline solid (1.80 g, 38%): mp 78-80 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) δ 7.69 (s, 1H), 7.14 (d, 1H, J = 8.1 Hz), 6.59-6.50 (m, 2H), 5.68 (q, 1H, J = 7.1 Hz), 4.39-4.24 (m, 2H), 3.82 (s, 3H), 1.34 (t, 3H, J = 7.3 Hz). FABLRMS m/z 303 (M+H). FABHRMS m/z 303.0849 (M+H Calc'd 303.0844). Anal. Calc'd for C<sub>14</sub>H<sub>13</sub>F<sub>3</sub>O<sub>4</sub>: C, 55.63; H, 4.34. Found: C, 55.47; H, 4.31.

Step 2. Preparation of ethyl 6,8-dichloro-7-methoxy-2-trifluoromethyl-benzopyran-2H-3-carboxylate.


Chlorine gas (excess) was added to a stirred solution of the ester (Step 1) (1.35 g, 4.47 mmol) in HOAc (30 mL) until the yellow color persisted. After 20 minutes, the reaction was sparged with nitrogen causing the reaction to become straw colored. Zinc (0.86 g, 13.40 mmol) was added to this solution with vigorous stirring. After 45 minutes, additional zinc (0.86 g, 13.40 mmol) was added and the reaction was stirred overnight. The

crude mixture was diluted with EtOH and filtered through diatomaceous earth. The filtrate was concentrated *in vacuo* yielding a crystalline mass. This solid was dissolved in EtOAc, washed with 2N HCl, brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* yielding an oil. The oil was dissolved in a minimum of isooctane, inducing crystallization. Vacuum filtration of the suspension yielded tan needles (1.078 g) which were recrystallized from isooctane yielding the dichloro ester as tan crystals (0.71 g, 43%) of suitable purity to use in the next step: mp 113.3-115.1 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.88 (s, 1H), 7.63 (s, 1H), 6.02 (q, 1H, *J* = 6.8 Hz), 4.38-4.22 (m, 2H), 3.93 (s, 3H), 1.31 (t, 3H, *J* = 7.1 Hz). <sup>19</sup>F NMR (acetone-d<sub>6</sub>/282 MHz) -80.00 (d, *J* = 7.2 Hz).

Step 3: Preparation of 6,8-dichloro-7-methoxy-2-trifluoromethyl-benzopyran-2H-3-carboxylic acid.

To a stirred solution of the dichloro ester from Step 2 (0.686 g, 1.848 mmol) in THF (10 mL) and EtOH (3 mL) was added NaOH (0.81 mL of 2.5 M aqueous solution, 2.03 mmol) in one portion. After stirring overnight the reaction was partially concentrated, diluted with H<sub>2</sub>O and washed with diethyl ether. The resulting aqueous phase was sparged with nitrogen and acidified with 2N HCl solution causing the solution to become turbid. Filtration of this suspension yielded the title compound as a white powder (0.559 g, 88 %): mp 195.6-199.1 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.90 (s, 1H), 7.64 (s, 1H), 6.01 (q, 1H, *J* = 6.8 Hz), 3.94 (s, 3H). <sup>19</sup>F NMR (CDCl<sub>3</sub>/282 MHz) -79.63 (d, *J* = 7.1 Hz). FAB/MS *m/z* 349 (M+Li). EIMS *m/z* 341.9681 (M+, Calc'd 341.9673). Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>4</sub>: C, 42.01; H, 2.06. Found: C, 41.76; H, 2.14.

## EXAMPLE 62

5     **2-Trifluoromethyl-2H-naphtho[1,2-b]pyran-3-carboxylic acid**


Step 1. Preparation of ethyl 2-trifluoromethyl-3H-naphthopyran-carboxylate.

A mixture of 2-hydroxy-1-naphthaldehyde (8.6 g, 0.050  
10 mol) and ethyl 4,4,4-trifluorocrotonate (9.2 g, 0.055 mol)  
dissolved in anhydrous dimethylformamide (DMF) and treated  
with anhydrous K<sub>2</sub>CO<sub>3</sub> (13.8 g, 0.100 mol). The solution was  
maintained at room temperature for 50 hours and diluted with  
water. The solution was extracted with ethyl acetate, and  
15 the combined extracts were washed with brine, dried over  
anhydrous MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to  
afford 4.8 g of an oil. The oil was purified by HPLC,  
eluting with hexanes:ethyl acetate (30:1). The appropriate  
fractions were concentrated to afford 1.6 g (10%) of the  
20 naphthopyran ester as a yellow solid.

Step 2. Preparation of 2-trifluoromethyl-3H-naphthopyran-carboxylic acid. A solution of the ester from Step 1 (0.8  
g, 2.5 mmol) was dissolved in 40 mL of ethanol and 10 mL of  
25 tetrahydrofuran, treated with sodium hydroxide (2.5 N, 10  
mL, 25 mmol) and stirred at room temperature for 16 hours.  
The reaction mixture was acidified with 1.0 N HCl, whereupon  
a solid formed that was isolated by filtration. The solid  
was washed with 20 mL of water to afford 0.7 g (95%) of the  
30 title compound as a yellow solid: mp 245.9-248.6 °C. <sup>1</sup>H NMR  
(acetone-d<sub>6</sub>/300 MHz) 8.57 (s, 1H), 8.28 (d, 1H, J = 8.7 Hz),  
8.03 (d, 1H, J = 9.0 Hz), 7.93 (d, 1H, J = 8.7), 7.67 (m,  
1H), 7.50 (m, 1H), 7.28 (d, 1H, J = 9.0), 5.96 (q H-F, 1H, J

= 7.2 Hz). FABHRMS  $m/z$  295.0561 (M+H, Calc'd 295.0582).  
Anal. Calc'd for  $C_{15}H_9O_3F_3 + 3.31\% H_2O$ : C, 59.21; H, 3.35.  
Found: C, 59.17; H, 3.07.


5

**EXAMPLE 63****2-Trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid**

10


2-Hydroxy-naph-1-aldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 244.7-249.8 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 8.61 (s, 1H), 8.09 (d, 1H,  $J = 8.3$  Hz), 7.90 (d, 1H,  $J = 8.9$  Hz), 7.82 (d, 1H,  $J = 8.3$  Hz), 7.63 (t, 1H,  $J = 8.1$  Hz), 7.47 (t, 1H,  $J = 8.1$  Hz), 7.23 (d, 1H,  $J = 9.1$  Hz), 5.84 (q, 1H,  $J = 6.8$  Hz).  $^{19}F$  NMR ( $CDCl_3$ /282 MHz) -79.56 (d,  $J = 7.3$  Hz). FABLRMS  $m/z$  295 (M+H). FABHRMS  $m/z$  295.0560 (M+H, Calc'd 295.0582).  
Anal. Calc'd for  $C_{15}H_9F_3O_3$ : C, 61.23; H, 3.08. Found: C, 60.85; H, 3.12.

20

**EXAMPLE 64**

**2-Trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid**

3-Hydroxynaphthalene-2-carboxylic acid was converted to 3-hydroxynaphthalene-2-carboxaldehyde by a similar procedure to that described in Example 24, Steps 1 & 2. The 3-hydroxynaphthalene-2-carboxaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp decompose >300 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.99 (s, 1H), 7.90 (s, 1H), 7.84 (d, 1H, *J* = 8.2 Hz), 7.74 (d, 1H, *J* = 8.2 Hz), 7.50 (t, 1H, *J* = 8.2 Hz), 7.39 (t, 1H, *J* = 8.2 Hz), 7.34 (s, 1H), 5.77 (q, 1H, *J* = 6.6 Hz). EIHMS *m/z* 294.0474 (M<sup>+</sup>, Calc'd 294.0504).

**EXAMPLE 65****6-Chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid****Step 1: Synthesis of 5-chloro-thiosalicylaldehyde.**

Tetramethylethylenediamine (TMEDA) (10.44 mL, 8.035 g, 69.15 mmol) was added via syringe to n-BuLi (43.22 mL of 1.6 M in hexanes, 69.15 mmol) and the solution was chilled to 0 °C. A solution of 4-chlorothiophenol (5.00 g, 34.57 mmol) in cyclohexane (25 mL) was added with stirring over 1 hour. The resulting tan slurry was stirred overnight at room temperature, chilled to 0 °C, and DMF (2.94 mL, 2.78 g, 38.03 mmol) was added via syringe over 2 minutes. The resulting gummy slurry was stirred at room temperature for 30 hours and became a powdery suspension. A mixture of 2 N HCl and ice was added to the reaction mixture until the pH

became acidic (pH = 1). During this addition, the mixture warmed and became first red and then pale yellow. This mixture was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over  $\text{MgSO}_4$ ,  
5 filtered and concentrated in vacuo yielding a clear red-brown oil. This oil was triturated with hexanes yielding a red-brown semisolid. This semisolid was purified by plug flash chromatography over silica gel, eluting with 1:1, hexanes:dichloromethane to afford 5-chloro-  
10 thiosalicylaldehyde (0.858 g, 14%) as an intensely yellow solid suitable for use without further purification.

Step 2: Preparation of ethyl 6-chloro-2-trifluoromethyl-benzo-1-thiopyran-2-H-3-carboxylate.

15 5-Chloro-thiosalicylaldehyde (Step 1) (0.84 g, 4.86 mmol) was added to DMF (3 mL) and ethyl 4,4,4-trifluorocrotonate (1.10 mL, 1.22 g). With stirring,  $\text{K}_2\text{CO}_3$  (0.67 g, 4.86 mmol) was added causing the reaction to become a deep red. After stirring overnight at room temperature,  
20 the reaction was diluted with diethyl ether and washed with water, saturated  $\text{NaHCO}_3$  solution, aqueous  $\text{KHSO}_4$  solution (0.25 M), brine, dried over  $\text{MgSO}_4$ , filtered and concentrated in vacuo yielding an oil. The oil was purified by flash chromatography (5:1; hexanes: ethyl acetate) yielding upon  
25 concentration ethyl 6-chloro-2-trifluoromethyl-benzo-1-thiopyran-2-H-3-carboxylate as a bright orange solid (0.492 g, 31%): mp 94.6-97.4 °C  $^1\text{H}$  NMR (acetone  $d_6$ /300 MHz)  $\delta$  8.01 (s, 1H), 7.71 (d, 1H,  $J = 2.2$  Hz), 7.50 (d, 1H,  $J = 8.5$  Hz), 7.44 (d of d, 1H,  $J = 2.3, 8.3$  Hz), 5.07 (q, 1H,  $J =$   
30 8.5 Hz), 4.42-4.23 (m, 2H), 1.35 (t, 3H,  $J = 7.1$  Hz).  
FABLRMS  $m/z$  329 (M+Li).


Step 3: Preparation of 6-chloro-2-trifluoromethyl-benzo-1-thiopyran-2-H-3-carboxylic acid.

35 To a stirred solution of the ester from Step 2 (0.413 g, 1.280 mmol) in THF: EtOH :  $\text{H}_2\text{O}$  (7 : 2 : 1, 10 mL) was



added NaOH solution (0.56 mL of 2.5 N solution, 1.408 mmol) with stirring. After stirring overnight, the reaction was partially concentrated *in vacuo* to remove the organic solvents, diluted with H<sub>2</sub>O and washed with several portions of diethyl ether. Acidification of the stirred aqueous phase with concentrated HCl caused precipitation of a flocculent yellow precipitate. Vacuum filtration of the suspension yielded 6-chloro-2-trifluoromethyl-benzo-1-thiopyran-2H-3-carboxylic acid as a yellow powder (0.25 g, 66 %): mp 188.8-198.7 °C <sup>1</sup>H NMR (acetone d<sub>6</sub>/300 MHz) δ 8.02 (s, 1H), 7.71 (d, 1H, *J* = 2.22 Hz), 7.50 (d, 1H, *J* = 8.5 Hz), 7.44 (d of d, 1H, *J* = 2.2, 8.5 Hz), 5.05 (q, 1H, *J* = 8.6 Hz). <sup>19</sup>F NMR (Acetone d<sub>6</sub>/282 MHz) δ -75.22 (d, *J* = 8.7 Hz). FABLRMS *m/z* 301 (M+Li); ESLRMS (neg. ion) *m/z* 293 (M-H).

#### EXAMPLE 66



#### (S)-6-Chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid


To a solution of 6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (Example 1, Step 2) (12.00 g, 43.07 mmol) and (S)(-)- $\alpha$ -methylbenzylamine (2.61 g, 21.54 mmol) in methyl-*tert*-butyl ether (30 mL) was slowly added *n*-heptane (200 mL) until the mixture became cloudy. The mixture was heated (steam bath) to boiling and set aside for 24 h during which time crystals formed. Filtration of the suspension yielded a crystalline product (5.5 g) which was recrystallized from methyl-*tert*-butyl ether (30 mL) and *n*-heptane (200 mL) yielding upon filtration a white solid (3.1

g). This solid was dissolved in EtOAc (100 mL) and washed with 1 N hydrochloric acid (50 mL) and brine (2 x 50 mL), dried over  $\text{MgSO}_4$  and concentrated *in vacuo* yielding a white solid. Recrystallization of this solid from methyl-t-butyl ether/n-heptane yielded the title compound as the highly enriched isomer, a white solid (2.7 g, 45%): mp 126.7-128.9 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 7.78 (s, 1H), 7.3-7.1 (m, 3H), 6.94 (d, 1H,  $J = 8.7$  Hz), 5.66 (q, 1H,  $J = 6.9$  Hz). Anal. Calc'd for  $\text{C}_{11}\text{H}_6\text{O}_3\text{F}_3\text{Cl}$ : C, 47.42; H, 2.17; N, 0.0. Found: C, 47.53; H, 2.14; N, 0.0. This compound was determined to have an optical purity of greater than 90% ee.

Procedure for determining optical purity.

To a solution of the free acid (title compound) (0.005 g, 0.017 mmol) in ethyl acetate (1.5 mL) in a test tube was added (trimethylsilyl)diazomethane (30  $\mu\text{L}$  of 2.0 N solution in hexanes, 60 mmol). The resulting yellow solution was warmed until the solution began to gently boil and then was allowed to cool to room temperature and stand for 0.08 hours. With vigorous mixing, the solution was quenched with aqueous 1 N HCl (1.5 mL). The layers were separated and a sample of the ethyl acetate fraction (0.3 mL) was transferred to a vial, concentrated under a stream of nitrogen, was diluted with hexane (total of 1 mL) and a sample (10  $\mu\text{L}$ ) analyzed by chiral chromatography. The HPLC utilized a Daicel ChiralPak AD column eluting with 10% isopropanol-hexane at 0.5 mL/min using a UV detector set at 254 nm.


**EXAMPLE 67**



(S)-6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

To a solution of 6-trifluoromethoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 16) (17.72 g, 54.00 mmol) and (-)-cinchonidine (7.95 g, 27.04 mmol) in methyl-tert-butyl ether (100 mL) heated on a steam-bath was added n-heptane (200 mL). The mixture was heated on the steam bath to boiling and allowed to cool for 4 h during which time crystals formed. Filtration of the suspension yielded a crystalline solid (18.7 g). This solid was dissolved in 2-butanone (30 mL) followed by the addition of n-heptane (500 mL). After standing for 16 hours, the resulting suspension was filtered yielded a white solid (10.3 g). This solid was dissolved in ethyl acetate (150 mL), washed with 1 N hydrochloric acid (100 mL) and brine (2 x 50 mL), dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo* yielding a viscous yellow oil (5.2 g, 59%):  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.16 (s, 1H), 6.77 (d, 1H,  $J = 2.7$  Hz), 6.94 (d, 1H,  $J = 8.7$  Hz), 6.64 (m, 1H), 6.39 (d, 1H,  $J = 8.7$  Hz) 5.13 (q, 1H,  $J = 7.2$  Hz). Anal. Calc'd for  $\text{C}_{12}\text{H}_6\text{O}_4\text{F}_6$ : C, 43.92; H, 1.84; N, 0.0. Found: C, 43.79; H, 1.83; N, 0.0. This compound was determined to have an optical purity of greater than 90% ee. Chiral purity was determined as describe in Example 66.


EXAMPLE 68



(S)-6-Chloro-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

To a solution of 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (Example 8) (11.4 g, 34.1 mmol) and (S)-(-)-2-amino-3-phenyl-1-propanol (2.57 g, 17.00 mmol) was added n-heptane (200 mL) and the mixture set aside for 16 h. The resulting suspension was filtered yielding a solid (3.8 g). This solid was recrystallized from 2-butanone (20 mL) and n-heptane (200 mL) yielding upon filtration a white solid (3.0 g). This solid was dissolved in ethyl acetate (100 mL) and washed with 1 N hydrochloric acid (50 mL) and brine (2 x 50 mL), dried over MgSO<sub>4</sub> and concentrated *in vacuo* yielding a white solid. This solid was recrystallized from n-heptane yielding the title compound of high optical purity as a crystalline solid (1.7 g, 30%): mp 175.4-176.9 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.86 (s, 1H), 7.52 (s, 1H), 7.12 (s, 1H), 5.83 (q, 1H, *J* = 7.1 Hz), 1.48 (s, 9H). Anal. Calc'd for C<sub>15</sub>H<sub>14</sub>O<sub>3</sub>F<sub>3</sub>Cl: C, 53.83; H, 4.22; N, 0.0; Cl, 10.59. Found: C, 53.78; H, 4.20; N, 0.0; Cl, 10.65. This compound was determined to have an optical purity of greater than 90% ee. Chiral purity was determined as describe in Example 66.

### EXAMPLE 69




#### 6-[[[(2-Furanylmethyl)amino]sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

The title compound was prepared by a similar procedure to that described in Example 49: mp 170-173 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.78 (s, 1H), 7.66-7.76 (m, 2H), 7.18-7.22 (m, 1H), 7.00-7.08 (m, 1H), 6.12-6.18 (m, 1H), 6.02-6.06

(m, 1H), 5.85 (q, 1H,  $J = 7.0$  Hz), 4.13 (s, 2H). EIHRMS  $m/z$  403.0332 ( $M^+$ , Calc'd 403.0337).


### EXAMPLE 70



**6-[(Phenylmethyl)sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared analogous to the procedure described in Example 56: mp 172-176 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300$  MHz) 7.73 (s, 1H), 7.43-7.56 (m, 2H), 7.21-7.33 (m, 3H), 7.20-7.21 (m, 3H), 5.88 (q, 1H,  $J = 7.0$  Hz), 4.83 (s, 2H). EIHRMS  $m/z$  398.0399 ( $M^+$ , Calc'd 398.0436).

### EXAMPLE 71




**6-[[[(Phenylethyl)amino]sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared analogous to the procedure described in Example 49: mp 187-190 °C.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}/300$  MHz) 7.82 (s, 1H), 7.74-7.90 (m,

2H), 7.08-7.29 (m, 6H), 5.89 (q, 1H,  $J = 6.8$ ), 3.12 (t, 2H,  $J = 7.3$  Hz), 2.72 (t,  $J = 7.3$  Hz). EIHRMS  $m/z$  427.0675 ( $M^+$ , Calc'd 427.0701)


### EXAMPLE 72



#### 7-Chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

4-Chlorosalicylic acid was converted to 3-chlorosalicylaldehyde by a procedure similar to that described in Example 24, Steps 1 & 2. The 3-chlorosalicylaldehyde was converted to the title compound by a procedure similar to Example 1: mp 175.2-177.6 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.90 (s, 1H), 7.51 (d, 1H,  $J = 7.8$  Hz), 7.12 (m, 2H), 5.86 (q  $\text{H-F}$ , 1H,  $J = 7.2$  Hz). FABHRMS  $m/z$  285.0114 ( $M+\text{Li}$ , Calc'd 285.0118). Anal. Calc'd for  $\text{C}_{11}\text{H}_6\text{ClF}_3\text{O}_3$ : C, 47.42; H, 2.17; Cl, 12.72. Found: C, 47.54; H, 2.37; Cl, 12.85.

### EXAMPLE 73



#### 6-Chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

Step 1. Preparation of 3-iodo-5-chlorosalicylaldehyde

N-Iodosuccinimide (144.0 g, 0.641 mole) was added to a solution of 5-chlorosalicylaldehyde (100 g, 0.638 mole) in dimethylformamide (400 mL). The reaction mixture was stirred for two days at room temperature. Additional N-iodosuccinimide (20 g, 0.089 mole) was added and the stirring was continued for an additional two days. The reaction mixture was diluted with ethyl acetate (1 liter), washed with hydrochloric acid (300 mL, 0.1 N), water (300 mL), sodium thiosulfate (300 mL, 5%), and brine (300 mL). It was dried over  $\text{MgSO}_4$ , and was concentrated to dryness to afford the desired aldehyde as a pale yellow solid (162 g, 90%): mp 84.8-86.7 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 11.67 (s, 1H), 9.71 (s, 1H), 7.92 (d, 1H,  $J = 2.5$  Hz), 7.54 (d, 1H,  $J = 2.6$  Hz). FABLRMS  $m/z$  281.0 (M-H). ESHRMS  $m/z$  280.8851 (M-H, Calc'd. 280.88630).


Step 2. Preparation of ethyl 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate

5-Chloro-3-iodosalicylaldehyde (20 g, 70.8 mmol), ethyl 4,4,4-trifluorocrotonate (17.85 g, 106 mmol), and triethylamine (14.33 g, 142 mmol) were dissolved in DMSO (200 mL). The reaction mixture was stirred at 90 °C for three days. The reaction mixture was poured into ethyl acetate (800 mL). It was extracted with 10% HCl (2 x 200 mL), saturated aqueous  $\text{NaHCO}_3$  (2 x 200 mL), and water (2 x 200 mL). The ethyl acetate phase was dried over  $\text{MgSO}_4$ , filtered and evaporated to yield a brown solid. It was then run through a plug of silica with ethyl acetate-hexane (1:20). The solvent was evaporated to give a yellow solid, that was recrystallized in hexane to afford the ester as a white solid (19.61 g, 64%): mp 92.1-93.9 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 7.71 (d, 1H,  $J = 2.2$  Hz), 7.56 (s, 1H), 7.20 (d, 1H,  $J = 2.2$  Hz), 5.81 (q, 1H,  $J = 6.7$  Hz), 4.37-4.29 (m, 2H), 1.35 (t, 3H,  $J = 7.2$  Hz). FABLRMS  $m/z$  431.9 (M-H). EIHRMS  $m/z$  431.9269 (M-H, Calc'd. 431.9237).

Step 3. Preparation of 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester (Step 2) was converted to the acid by a procedure similar to the method described in Example 1, Step 2: mp 220-223 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.77 (d, 1H, *J* = 2.2 Hz), 7.71 (s, 1H), 7.41 (d, 1H, *J* = 2.2 Hz), 5.87 (q, 1H, *J* = 7.0 Hz). EIHRMS *m/z* 403.8893 (M-H, Calc'd. 403.8924). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>ClF<sub>3</sub>IO<sub>3</sub>: C, 32.66; H, 1.25. Found: C, 33.13; H, 1.29.

**EXAMPLE 74**



**8-Bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of ethyl 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate

A mixture of 3-bromo-5-chlorosalicylaldehyde (1.9 g, 4.2 mmol), potassium carbonate (0.58 g, 4.2 mmol), and ethyl 4,4,4-trifluorocrotonate (0.79 g, 4.7 mmol) was stirred in *N,N*-dimethylformamide (5 mL) at 95° C for 18 h. Water (100 mL) was added and the mixture was extracted with ether (3 x 50 mL). The combined organic extracts were washed with sodium hydroxide (10 mL) and water (2 x 50 mL). After drying over MgSO<sub>4</sub> and concentrating, the mixture filtered through of a pad of silica eluting with ethyl acetate-hexanes (1:4). The eluant was concentrated and a light yellow solid was crystallized from cold hexane (0.43 g, 26%): mp 101.0-102.2 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.65 (d, H, *J* = 2.4 Hz), 7.61 (d, H, *J* = 2.4 Hz),




6.03 (q H-F, 1H,  $J = 6.9$  Hz), 4.34 (m, 2H), 1.33 (t, 3H,  $J = 7.5$  Hz). ESHRMS  $m/z$  384.9435 (M-H, Calc'd 384.9454). Anal. Calc'd for  $C_{13}H_9BrClF_3O_3$ : C, 40.50; H, 2.35. Found: C, 40.61; H, 2.40.

Step 2. Preparation of 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

Ethyl 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate (0.3 g), ethanol (15 mL), tetrahydrofuran (10 mL), and sodium hydroxide solution (10 mL, 2.5 N) were stirred at room temperature for 16 h. Hydrochloric acid (1 N) was added until the mixture was acidic to pH paper. The addition of water (50 mL) caused the formation of a precipitate which was collected by filtration yielding the title compound as a white solid (0.2 g, 72%): mp 227.8-228.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.90 (s, 1H), 7.65 (dd, 2H,  $J = 2.4$  and  $J = 28.8$  Hz), 6.00 (q H-F, 1H,  $J = 7.2$  Hz). FABHRMS  $m/z$  356.9134 (M+H, Calc'd 356.9141). Anal. Calc'd for  $C_{11}H_5BrClF_3O_3$ : C, 36.96; H, 1.41. Found: C, 37.05; H, 1.33.

**EXAMPLE 75**



**6-Formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of ethyl 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


A 50 mL round bottom flask was charged with 5-formylsalicylaldehyde (3.21 g, 21.39 mmol), ethyl 4,4,4-trifluorocrotonate (3.50 mL, 3.96 g, 23.53 mmol),

dimethylformamide (15 mL) and potassium carbonate (2.95 g, 21.39 mmol) and heated to 60 °C for 12 hours. Additional ethyl 4,4,4-trifluorocrotonate (3.50 mL, 3.96 g, 23.53 mmol) was added and the reaction heated for 16 hours at 75 °C. After cooling to room temperature, the reaction was partitioned between H<sub>2</sub>O and diethyl ether. The organic phase was washed with saturated NaHCO<sub>3</sub> solution, KHSO<sub>4</sub> solution (0.25 M), brine, treated with decolorizing carbon (warmed gently). The resulting black suspension was dried over MgSO<sub>4</sub>, vacuum filtered through diatomaceous earth, and concentrated *in vacuo* yielding an orange crystalline mass. This material was recrystallized from hot hexanes yielding the ester (1.51 g, 24 %) as orange crystals: mp 84.3-86.2 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 9.96 (s, 1H), 8.06 (d, 1H, *J* = 2Hz), 8.02 (s, 1H), 7.99 (dd, 1H, *J* = 8.5, 2.0Hz), 7.24 (d, 1H, *J* = 8.5 Hz), 5.99 (q, 1H, *J* = 7.1 Hz), 4.43-4.25 (m, 2H), 1.34 (t, 3H, *J* = 7.3 Hz). FABLRMS *m/z* 301 (M+H). EIHRMS *m/z* 300.0605 (M+, Calc'd 300.0609). Anal. Calc'd for C<sub>14</sub>H<sub>11</sub>F<sub>3</sub>O<sub>4</sub>: C, 56.01; H, 3.69. Found: C, 56.11; H, 3.73.

Step 2. Preparation of 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester (Step 1) was converted to the acid via a method similar to that described in Example 1, Step 2: mp 211.3-215.7 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 9.97 (s, 1H), 8.07 (d, 1H, *J* = 2.0Hz), 8.03 (s, 1H), 8.00 (dd, 1H, *J* = 8.3, 2.0 Hz), 7.25 (d, 1H, *J* = 8.5 Hz), 5.98 (q, 1H, *J* = 6.9 Hz). FABLRMS *m/z* 273 (M+H). EIHRMS *m/z* 272.0266 (M+, Calc'd 272.0296). Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>F<sub>3</sub>O<sub>4</sub>: C, 52.95; H, 2.59. Found: C, 52.62; H, 2.58.

**EXAMPLE 76**



**6-Chloro-8-Formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of 4-chloro-2,6-bis(hydroxymethyl)phenol.

Potassium hydroxide (84.82 g, 1.30 mole) was dissolved in H<sub>2</sub>O (200 mL) in a two liter 3-neck round bottom flask fitted with thermocouple, mechanical stirrer, and stopper. With stirring, 4-chlorophenol (128.56 g, 1.0 mole) was added with cooling (ice bath) resulting in the temperature rising to 26 °C. Formalin (230 mL of 37% aqueous solution, 2.83 mole) was added portion-wise maintaining the temperature below 25 °C. The reaction was warmed to 35 °C for 48 hours. To this solution was added aqueous acetic acid (80.0 mL, 84.1 g, 1.40 mole in 800 mL H<sub>2</sub>O) causing the solution to become turbid. Vacuum filtration of the suspension yielded a tan solid. The solid was stirred with acetone (100 mL) and the insoluble product collected by vacuum filtration. The solution was diluted with hexanes yielding several crops of the diol as fine tan needles (35.0 g, 19%). mp 160.6-163.3 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>, NaOD, D<sub>2</sub>O/300 MHz) 6.69 (s, 2H), 4.48 (s, 4H), 7.88 (d, 1H, J = 2.6 Hz), 7.75 (d, 1H, J = 2.6 Hz), 6.08 (q, 1H, J = 6.9 Hz). ESLRMS m/z 206 (M+NH<sub>4</sub><sup>+</sup>). ESHRMS m/z 187.0131 (M-H, Calc'd 187.0162).

Step 2. Preparation of 5-chloro-3-formyl-salicylaldehyde.

To a stirred suspension of diol (Step 1) (33.0 g, 0.18 mole) in chloroform (1.5 L) in a 2 L round bottom flask was added manganese dioxide (139 g, 1.60 mole) and the resulting suspension heated to a gentle reflux for 10 hours. The reaction was allowed to cool to room temperature, was filtered through diatomaceous earth, concentrated in vacuo, presorbed on silica gel and purified by flash chromatography

(hexane/ethyl acetate) yielding the as a mustard colored powder dialdehyde (22.42 g, 67 %): mp 120.7-122.8 °C. This solid was of suitable purity to use in the next step without further purification.

Step 3. Preparation of ethyl 6-chloro-8-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


A stirred solution of the dialdehyde (Step 2) (1.13 g, 6.14 mmol), dimethyl sulfoxide (6 mL), ethyl 4,4,4-trifluorocrotonate (1.37 mL, 1.55g, 9.21 mmol) and triethylamine (1.71 mL, 1.24g, 12.28 mmol) in a round bottom flask fitted with condenser was heated to 80 °C for 8 h. Upon cooling to room temperature the reaction was diluted with diethyl ether (100 mL) and the resulting mixture washed with aqueous sodium bicarbonate solution (3 X 75 mL), 1 N HCl solution (3 X 70 mL), and brine (1 X 75 mL), dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding a tan powder. This powder was taken up in hot hexane-ethyl acetate and filtered to remove insoluble matter. Upon cooling of the filtrate, crystallization followed by vacuum filtration yielded the desired ester as tan crystals (0.726 g, 35%): mp 118.1-119.7 °C. This material was of suitable purity to use without further purification.

Step 4. Preparation of 6-chloro-8-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

To a stirred solution of the ester (Step 3) (0.284 g, 0.849 mmol) in THF:EtOH:H<sub>2</sub>O (7:2:1, 5 mL) was added aqueous NaOH solution (0.41 mL of 2.5 M, 1.02 mmol). After stirring 40 hours, the reaction was partially concentrated in vacuo to remove the organic solvents, diluted with H<sub>2</sub>O, washed with diethyl ether, sparged with nitrogen to remove trace diethyl ether, and acidified with concentrated HCl yielding a suspension. Vacuum filtration of the suspension yielded the title compound as a pale yellow powder (0.160 g, 23 %). mp 243.3-252.4 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 10.39 (s,

1H), 7.98 (s, 1H), 7.88 (d, 1H,  $J = 2.6$  Hz), 7.75 (d, 1H,  $J = 2.6$  Hz), 6.08 (q, 1H,  $J = 6.9$  Hz). FABLRMS  $m/z$  307 (M+H). ESHRMS  $m/z$  304.9839 (M-H, Calc'd 304.9828). Anal. Calc'd for  $C_{12}H_6Cl_1F_3O_4$ : C, 47.01; H, 1.97. Found: C, 46.64; H, 1.86.


### EXAMPLE 77



#### 6-Bromo-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

7-(1,1-Dimethylethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 12) (0.6 g, 2 mmol), chloroform (50 mL), iron filings (0.01 g, 0.2 mmol), and bromine (0.48 g, 3.00 mmol) were stirred at reflux for 16 h. The mixture was allowed to cool and was washed with brine (2 x 50 mL). After drying over  $MgSO_4$ , the mixture was filtered, concentrated in vacuo, and the residue crystallized from ether-hexanes yielding the title compound as a white solid (0.5 g, 66%): mp 198.6-199.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.85 (s, 1H), 7.72 (s, 1H), 7.13 (s, 1H), 5.83 (q, 1H,  $J = 7.2$  Hz), 1.5 (s, 9H). Anal. Calc'd for  $C_{15}H_{14}O_3F_3Br$ : C, 47.52; H, 3.72; N, 21.07. Found: C, 47.42; H, 3.68; N, 21.15.


### EXAMPLE 78



**5,6-Dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5,6-Dichlorosalicylaldehyde was prepared by the procedure described in Cragoe, E.J.; Schultz, E.M., U.S. Patent 3 794 734, 1974. This salicylaldehyde was converted to the title compound by a similar procedure to that described in Example 1: mp 211.5-213.5 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 8.09 (s, 1H), 7.63 (d, 1H,  $J = 8.9$  Hz), 7.12 (d, 1H,  $J = 8.9$  Hz), 5.94 (q, 1H,  $J = 7.0$  Hz). ESLRMS  $m/z$  311 (M-H). EIHRMS  $m/z$  311.9583 (M+, Calc'd 311.9568). Anal. Calc'd for  $\text{C}_{11}\text{H}_5\text{Cl}_2\text{F}_3\text{O}_3$ : C, 42.20; H, 1.61. Found: C, 42.33; H, 1.67.

**EXAMPLE 79**



**6-Cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of ethyl 6-[(hydroxyimino)methyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

A 50 mL round bottom flask was charged with hydroxylamine HCl (0.255 g, 3.67 mmol), ethyl 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 75, Step 1) (1.00 g, 3.34 mmol), sodium acetate (0.301 g, 3.67 mmol), ethanol (10 mL), and  $\text{H}_2\text{O}$  (2 mL). The reaction was stirred at room temperature for 18 hours, then diluted with

H<sub>2</sub>O and diethyl ether. The layers were separated and the organic phase washed with H<sub>2</sub>O, brine, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo yielding an orange semi-crystalline mass. Recrystallization of this solid from hot ethyl acetate and isooctane yielded the oxime (0.578 g, 55%): mp 113.0-116.2 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 10.46 (s, ca.1 exch.), 8.11 (s, 2H), 7.92 (s, 1H), 7.72 (d, 1H, J = 2Hz), 7.68 (dd, 1H, J = 8.5, 2.0Hz), 7.07 (d, 1H, J = 8.5 Hz), 5.89 (q, 1H, J = 7.1 Hz), 4.43-4.22 (m, 2H), 1.34 9t, 3H, J = 7.3 Hz). FABLRMS m/z 316 (M+H). EIHRMS m/z 315.0719 (M+, Calc'd 315.0733). Anal. Calc'd for C<sub>14</sub>H<sub>12</sub>F<sub>3</sub>N<sub>1</sub>O<sub>4</sub>: C, 53.34; H, 3.84; N 4.44. Found: C, 53.85; H, 3.90; N, 4.19.

Step 2. Preparation of ethyl 6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

To a stirred solution of oxime (Step 1) (0.264 g, 0.840 mmol) in dioxane (4.5 mL) in a 25 mL pear-shaped flask was added trifluoroacetic anhydride (0.130 mL, 0.194 g, 0.924 mmol) and triethylamine (0.140 mL, 0.102 g, 1.008 mmol). The reaction was stirred at room temperature for 12 hours, then heated to 85 °C for 4 hours. After cooling to room temperature, aqueous HCl (50 mL, 1 N HCL) was added, and the resulting mixture extracted with ethyl acetate. The ethyl acetate phase was washed with chilled aqueous HCl (1 N), brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated in vacuo yielding a pale yellow oil. This oil was resubmitted to similar reaction conditions. After dissolution of the pale yellow oil in dioxane (4.5 mL), trifluoroacetic anhydride (0.130 mL, 0.194 g, 0.924 mmol) and triethylamine (0.140 mL, 0.102 g, 1.008 mmol) were then added. After stirring 3 hours at room temperature, more triethylamine 0.50 mL, 0.36 g, 3.6 mmol) was added and then heated to 85 °C for 3 hours. After cooling to room temperature, aqueous HCl (50 mL, 1 N HCL) was added, and the resulting mixture extracted with ethyl acetate. The ethyl acetate phase was washed with


chilled aqueous HCl (1 N), brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, concentrated in vacuo yielding a pale yellow oil. Addition of hexanes induced crystallization followed by vacuum filtration yielded the title compound (0.101g, 40 %) as a yellow powder: mp 101.6-106.1 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.97 (d, 1H,  $J$  = 2.2 Hz), 7.95 (s, 1H), 7.82 (dd, 1H,  $J$  = 8.5, 2.0Hz), 7.24 (d, 1H,  $J$  = 8.5 Hz), 6.01 (q, 1H,  $J$  = 7.1 Hz), 4.38-4.24 (m, 2H), 1.34 (t, 3H,  $J$  = 7.3 Hz). FABLRMS  $m/z$  298 (M+H). EIHRMS  $m/z$  297.0575 (M+, Calc'd 297.0613).

Step 3. Preparation of 6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

To a stirred solution of the ester (Step 2) (0.077 g, 0.259 mmol) in THF-EtOH- $\text{H}_2\text{O}$  (7:2:1, 2 mL) in a 5 mL pear-shaped flask was added aqueous NaOH (0.13 mL, 2.5 N solution) in one portion. After stirring for 6 hours at room temperature the solution was partially concentrated in vacuo to remove most of the THF and EtOH. The resulting solution was diluted with  $\text{H}_2\text{O}$  and washed with diethyl ether. The resulting aqueous phase was sparged with nitrogen to remove trace diethyl ether and was acidified with concentrated HCl yielding a sticky suspension. The suspension was extracted with diethyl ether and the ether was dried over  $\text{MgSO}_4$ , filtered and concentrated in vacuo yielding a pale yellow oil. This oil was crystallized from methylene chloride-hexanes yielding the title compound (0.041 g, 59 %) as a tan powder: mp 185.1-186.1 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.99-7.94 (m, 2H), 7.83 (dd, 1H,  $J$  = 8.5, 2.0 Hz), 7.25 (d, 1H,  $J$  = 8.5 Hz), 5.99 (q, 1H,  $J$  = 7.0 Hz). FABLRMS  $m/z$  270 (M+H). EIHRMS  $m/z$  269.0316 (M+, Calc'd 269.0300).

**EXAMPLE 80**






**6-Hydroxymethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

To a chilled (ice bath), stirred solution of 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 75, Step 2) (0.133 g, 0.489 mmol) in THF (1 mL) and ethanol (1 mL) in a 10 mL round bottom flask was added NaBH<sub>4</sub> (0.020 g, 0.528 mmol) in two portions. The reaction was allowed to warm to room temperature and more NaBH<sub>4</sub> (0.050 g, 1.322 mmol) was added. The total reaction time was 3 hours. The reaction was quenched with aqueous HCl (1 N solution) and was extracted with chloroform. The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* yielding a foam. This crude product was purified by flash chromatography (silica gel 60, eluant 1:1, hexane-ethyl Acetate with 2 % acetic acid). The product collected from the chromatography was recrystallized from hexanes and ethyl acetate, and collected by vacuum filtration yielding the title compound (0.042 g, 31 %) as a very pale yellow powder: mp 177.5-180.8 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.89 (s, 1H), 7.44 (s, 1H), 7.41 (d, 1H, *J* = 8.3 Hz), 6.99 (d, 1H, *J* = 8.3 Hz), 5.80 (q, 1H, *J* = 7.3 Hz), 4.59 (s, 2H). FAB/MS *m/z* 275 (M+H). EIMS *m/z* 274.0417 (M+, Calc'd 274.0453). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>4</sub>: C, 52.57; H, 3.31. Found: C, 52.43; H, 3.34.

**EXAMPLE 81**



6-(Difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

Step 1. Preparation of ethyl 6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


Ethyl 6-formyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate (Example 75, Step 1) (1.672 g, 5.569 mmol) in methylene chloride (1.5 mL) was added to methylene chloride (1.5 mL) and diethylaminosulfur trifluoride (DAST) (0.74 mL, 0.898 g, 5.569 mmol) over 0.07 hours via syringe. After stirring for 20 hours the reaction was poured into aqueous HCl (2.0 N) and the mixture was extracted with diethyl ether. The ethereal phase was washed with dilute aqueous HCl (2.0 N), saturated NaHCO<sub>3</sub> solution, brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* yielding a clear colorless oil. This oil was purified by flash chromatography (Silica gel 60, Eluant (5:1; Hexanes : Ethyl Acetate) yielding ethyl 6-difluoromethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate (0.96 g, 54 %) as an oil which solidified upon standing. This product was of sufficient purity to be used in the next step without further purification: <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.97 (s, 1H), 7.74 (s, 1H), 7.65 (d, 1H, *J* = 8.5 Hz), 7.18 (d, 1H, *J* = 8.5 Hz), 6.90 (t, 1H, *J* = 56.0 Hz), 5.94 (q, 1H, *J* = 7.0 Hz), 4.40-4.25 (m, 2H), 1.34 (t, 3H, *J* = 7.0 Hz).

Step 2. Preparation of 6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

Aqueous NaOH (1.31 mL, 3.277 mmol, 2.5 M solution) was added in one portion to the ester (Step 1) (0.880 g, 2.731 mmol) in THF:EtOH:H<sub>2</sub>O (7:2:1, 10 mL). The resulting solution was stirred for 60 hours. The reaction mixture was partially concentrated *in vacuo* to remove the organic solvents and was diluted with H<sub>2</sub>O. The resulting aqueous solution was washed with diethyl ether, sparged with nitrogen to remove trace ether, and acidified with concentrated HCl. The resulting oily suspension was

extracted with diethyl ether. The combined organic phases were dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo* yielding the title compound (0.483 g, 60%) as an oil which solidified as a white crystalline mass: mp 134.7-136.2 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.97 (s, 1H), 7.73 (s, 1H), 7.67 (dd, 1H,  $J$  = 8.5, 1.0 Hz), 7.17 (d, 1H,  $J$  = 8.5 Hz), 6.89 (t, 1H,  $J$  = 56.2 Hz), 5.90 (q, 1H,  $J$  = 7.1 Hz). FAB-ESLRMS  $m/z$  293 (M-H). EIHRMS  $m/z$  293.0235 (M-H, Calc'd 293.0237). Anal. Calc'd for  $\text{C}_{12}\text{H}_7\text{F}_5\text{O}_3$ : C, 49.00; H, 2.40. Found: C, 48.78; H, 2.21.

## EXAMPLE 82



### 2,6-Bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

#### Step 1. Preparation of Ethyl 2,6-bis(trifluoromethyl)-4-oxo-4H-1-benzopyran-3-carboxylate.

To a stirred solution of ethyl 4,4,4-trifluoroacetoacetate (3.22 mL, 4.06 g, 22.07 mmol) in toluene (100 mL) was added portion-wise sodium hydride (0.971 g, of 60 % oil dispersion reagent, 22.07 mmol) causing gas evolution. After gas evolution has subsided, 2-fluoro-5-(trifluoromethyl)benzoyl chloride (5.00 g, 22.07 mmol) was added. The reaction was stirred at room temperature for 24 hours, then heated to 105 °C for 24 hours. After cooling to room temperature, the reaction was diluted with diethyl ether and the resulting solution was washed with  $\text{H}_2\text{O}$  and brine, dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo* yielding a slightly sticky white solid. This solid was triturated with hexanes yielding the desired ester (3.05 g, 39 %) as a white powder: mp 116-120.1 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 8.52 (d, 2H,  $J$  = 1.6 Hz), 8.03

(dd, 1H,  $J = 8.9, 2.2$  Hz), 7.71 (d, 1H,  $J = 8.9$  Hz), 4.48 (q, 2H,  $J = 7.3$  Hz), 1.39 (t, 3H,  $J = 7.3$  Hz). FABLRMS  $m/z$  355 (M+H). Anal. Calc'd for  $C_{14}H_8F_6O_4$ : C, 47.45; H, 2.28.

Found: C, 47.59; H, 2.43.

Step 2. Preparation of ethyl 2,6-bis(trifluoromethyl)-4-oxo-dihydrobenzopyran-3-carboxylate.

A 250 mL round bottom flask was charged with ethyl 2,6-bis(trifluoromethyl)-benzopyran-4-one-3-carboxylate (Step 1) (2.307 g, 6.513 mmol) and THF (20 mL) yielding a pale yellow solution. Ethanol (20 mL) was added and the reaction chilled in an ice-salt bath. While maintaining the reaction temperature at below 9 °C,  $NaBH_4$  (0.246 g, 6.513 mmol) was added in two portions and the mixture stirred 1 h. The crude reaction mixture was poured into a vigorously stirred mixture of ice (200 mL) and concentrated HCl (12 N, 5 mL) yielding a precipitate. Vacuum filtration of the resulting suspension yielded the desired keto ester (2.204 g, 87%) as faint pink powder of suitable purity to use in the next step without further purification: mp 71.8-76.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 12.71 (br s, 1H exch), 8.01 (d, 1H,  $J = 2.0$  Hz), 8.01 (d, 1H,  $J = 2.0$  Hz), 7.88 (dd, 1H,  $J = 8.7, 1.8$  Hz), 7.31 (d, 1H,  $J = 8.7$  Hz), 5.98 (q, 1H,  $J = 6.6$  Hz), 4.51-4.28 (m, 2H), 1.35 (t, 3H,  $J = 7.0$  Hz). FABLRMS  $m/z$  355 (M-H). ESHRMS  $m/z$  355.0394 (M-H, Calc'd 355.0405). Anal. Calc'd for  $C_{14}H_{10}F_6O_4$ : C, 47.21; H, 2.83. Found: C, 47.31; H, 2.97.

Step 3. Preparation of ethyl 2,6-bis(trifluoromethyl)-4-trifluoromethanesulfonato-2H-1-benzopyran-3-carboxylate.

A 50 mL 3-neck Morton flask fitted with addition funnel, 2 stoppers was charged with 2,6-di-tert-butylpyridine (1.576 g, 1.50 mmol), methylene chloride (12 mL), and then via syringe was added trifluoromethanesulfonic anhydride (1.08 mL, 1.80 g, 1.25 mmol). To this solution was added dropwise a solution the keto ester (Step 2) (1.822

g, 5.115 mmol) in methylene chloride (10 mL) over 0.33 h and the reaction stirred for 48 h. The resulting off-white suspension was transferred to a 100 mL round bottom flask and was concentrated *in vacuo*. The residue was suspended in diethyl ether (50 mL) and vacuum filtered to remove salts. The filtrate was further diluted with diethyl ether (50 mL) and was washed with ice cold HCl solution (2 N), brine, and dried over  $\text{Na}_2\text{CO}_3$ , filtered and concentrated *in vacuo* yielding the desired triflate (1.64 g, 66%) as a tan clumpy powder of suitable purity to use in the next step without further purification.

Step 4. Preparation of ethyl 2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


A 25 mL pear flask was charged with LiCl (0.136 g, 3.219 mmol), affixed to a high vacuum line and heated with a heat gun removing superficial water. The flask was allowed to cool to room temperature, and tetrakis(triphenylphosphine)palladium(0) (0.124 g, 0.107 mmol) and THF (2 mL) were added. A reflux condenser was affixed to the flask and the apparatus was purged with nitrogen. A solution of the triflate (Step 3) (0.524 g, 1.073 mmol) in THF (2 mL) and tri-n-butyltin hydride (0.32 mL, 0.34 g, 1.18 mmol) were added sequentially via syringe. The resulting light orange solution was heated to 50 °C with stirring for 1 h, 60 °C for one hour, and 65 °C for one hour. The reaction was allowed to cool to room temperature and was poured into 2 N HCl, stirred, and extracted with hexanes. The hexane phase was dried over  $\text{MgSO}_4$ , filtered and concentrated yielding a light brown oil. The oil was dissolved in hexane and was washed with aqueous ammonium fluoride solution. The resulting hexane phase was dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo* yielding a dull yellow oily solid which solidified as a flaky powder (0.443 g). This solid was purified by flash silica chromatography (eluant: hexanes-methylene chloride, 4:1) yielding ethyl 2,6-di-trifluoromethyl-2H-1-benzopyran-3-

carboxylate(0.069 g, 19 %) as a white crystalline solid of suitable purity to proceed with the next step.

Step 5. Preparation of 2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

To a stirred solution of the ester (Step 4) (0.065 g, 0.191 mmol) in THF-EtOH-H<sub>2</sub>O (7:2:1, 1 mL) was added NaOH solution (0.084 mL, 0.210 mmol) in one portion at room temperature and allowed to stir overnight. The reaction was partially concentrated in vacuo yielding a pale yellow clear syrup. The syrup was diluted with water (5 mL) and brine (1 mL) and was washed with diethyl ether (3 X 5 mL). The resulting aqueous phase was sparged with nitrogen to remove trace ether. With stirring, concentrated HCl was added to the aqueous phase causing the formation of a very fine white precipitate. This suspension was extracted with diethyl ether and the ether dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated by slow evaporation at atmospheric pressure. The resulting product was recrystallized from hexanes and ethyl acetate yielding the title compound (0.038 g, 64 %) as a fine tan powder: mp 143.5-145.2 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 11.97-11.67 (br s, 1H), 8.03 (s, 1H), 7.92 (s, 1H), 7.77 (d, 1H, *J* = 8.5 Hz), 7.26 (d, 1H, *J* = 8.7 Hz), 5.96 (q, 1H, *J* = 7.0 Hz). FAB/MS *m/z* 311 (M-H). ESHRMS *m/z* 311.0107 (M-H, Calc'd 311.0143).


**EXAMPLE 83**



**5,6,7-Trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

3,4,5-Trichlorophenol was converted to 3-ethoxysalicylaldehyde via a procedure similar to that described in Example 11, Step 1. The 4,5,6-trichlorosalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 236.2-239.3 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 8.05 (s, 1H), 7.40 (s, 1H), 5.99 (q, 1H,  $J = 7.0$  Hz). ESLRMS  $m/z$  345 (M-H). ESHRMS  $m/z$  344.9113 (M-H, Calc'd 344.9100). Anal. Calc'd for  $\text{C}_{11}\text{H}_4\text{Cl}_3\text{F}_3\text{O}_3 + 0.89$  wt %  $\text{H}_2\text{O}$ : C, 37.68; H, 1.25; Cl, 30.33. Found: C, 37.48; H, 1.25; Cl, 30.33.


#### EXAMPLE 84



#### 6,7,8-Trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

2,3,4-Trichlorophenol was converted to 3-ethoxysalicylaldehyde via a procedure similar to that described in Example 11, Step 1. The 3,4,5-trichlorosalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 222.0-225.3 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.94 (s, 1H), 7.78 (s, 1H), 6.07 (q, 1H,  $J = 7.0$  Hz). ESLRMS  $m/z$  345 (M-H). EIHRMS  $m/z$  344.9117 (M-H, Calc'd 344.9100). Anal. Calc'd for  $\text{C}_{11}\text{H}_4\text{Cl}_3\text{F}_3\text{O}_3 + 1.56$  wt %  $\text{H}_2\text{O}$ : C, 37.43; H, 1.32; Cl, 30.13. Found: C, 37.79; H, 0.93; Cl, 29.55.


#### EXAMPLE 85



**7-Ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

3-Ethylphenol was converted to the title compound by a procedure similar to that described in Example 2. : mp 167.0-168.6 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 7.84 (s, 1H), 7.15 (d, 1H,  $J = 7.5$  Hz), 6.84 (m, 2H), 5.66 (q, 1H,  $J = 6.8$  Hz), 2.63 (q, 2H,  $J = 7.7$  Hz,  $J = 7.7$  Hz), 1.24 (t, 3H,  $J = 7.7$  Hz). Anal. Calc'd for  $\text{C}_{13}\text{H}_{11}\text{F}_3\text{O}_3$ : C, 57.36; H, 4.07. Found: C, 57.25; H, 4.10.

**EXAMPLE 86**



**6-(Methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

**Step 1. Preparation of ethyl 6-(methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.**

Ethyl 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 2, Step 2) (1.014 g, 3.18 mmol) in methylene chloride was chilled to -50 °C (dry ice acetone). With stirring, meta-chloroperoxybenzoic acid (0.91 g of 60 % reagent, 3.18 mmol) was added and reaction allowed to proceed for 3 hours. Aqueous  $\text{NaHSO}_3$  solution (40 mL 0.25 M) was poured into the reaction. More methylene chloride was added and the layers mixed, then separated. The organic




phase was washed with aqueous NaHSO<sub>3</sub> solution, aqueous saturated NaHCO<sub>3</sub> solution, brine, dried over MgSO<sub>4</sub>, filtered and concentrated yielding an oil. The oil was diluted with isooctane (2 mL) and concentrated yielding an oil which upon  
5 standing crystallized. Hexanes was added, the solution was heated, and methylene chloride added until partial dissolution occurred. After cooling and standing overnight the suspension was vacuum filtered yielding the sulfoxide substituted ethyl ester (0.753 g, 71%) as white needles: mp  
10 92.2-98.4 °C. This ester was of sufficient purity to be used without further purification.

Step 2. Preparation of 6-(methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

15 To a stirred solution of the ester (Step 1) (0.683 g, 2.043 mmol) in THF:EtOH:H<sub>2</sub>O (7:2:1, 4 mL) was added aqueous NaOH solution (0.98 mL of 2.5 M, 2.45 mmol). After stirring 12 hours, the reaction was partially concentrated in vacuo to remove the organic solvents. The residue was diluted  
20 with H<sub>2</sub>O, washed with diethyl ether, sparged with nitrogen to remove trace diethyl ether, and acidified with concentrated HCl yielding a oily suspension. The suspension was extracted with diethyl ether, and the resulting organic phase dried over MgSO<sub>4</sub>, filtered, and diluted with hexanes.  
25 Upon concentration in vacuo the title acid was obtained as a sticky white powder (0.425 g, 68 %): mp 148.3-151.0 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.99 (s, 1H), 7.82 (s, 1H), 7.78-7.68 (m, 1H), 7.24 (d, 1H, *J* = 8.3 Hz), 5.92 (q, 1H, *J* = 7.1 Hz), 2.73 (s, 3H). FABLRMS *m/z* 307 (M+H). ESHRMS *m/z*  
30 305.0098 (M-H, Calc'd 305.0095). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>O<sub>4</sub>S<sub>1</sub>: C, 47.06; H, 2.96; S, 10.47. Found: C, 46.69; H, 2.86; S, 10.45.

**EXAMPLE 87**




**5,8-Dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5

2,5-Dichlorophenol was converted to 3,6-dichlorosalicylaldehyde via a procedure similar to that described in Example 2, Step 1. The 3,6-dichlorosalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 11, Steps 2 & 3: mp 205.7-207.1 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.02 (s, 1H), 7.53 (d, 1H, J = 8.7 Hz), 7.22 (d, 1H, J = 8.7 Hz), 6.04 (q, 1H, J = 7.1 Hz). FABLRMS m/z 311 (M-H). ESHRMS m/z 310.9506 (M-H, Calc'd 310.9490). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub> + 0.63 wt % H<sub>2</sub>O: C, 41.94; H, 1.67. Found: C, 41.54; H, 1.27.

**EXAMPLE 88**



20

**6-(Pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

25 Step 1. Preparation of ethyl 6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


Potassium pentafluoropropionate (0.476 g, 2.35 mmol) was dissolved in toluene (6 mL) and DMF (6 mL). The vessel was fitted with a distilling head, and CuI (0.471 g, 2.474 mmol) was added with stirring. The reaction was heated to

30

120 °C, removing the toluene by distillation. Ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 72, Step 3) (0.469 g, 1.178 mmol) was added and the reaction was heated to 150 °C for 2 hours. The reaction was allowed to cool to room temperature and was partitioned between diethyl ether and H<sub>2</sub>O. The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo. The resulting residue was purified by flash chromatography (silica gel 60, eluant: hexanes- ethyl acetate, 8:1) yielding, upon concentration of the solution, the desired ester (0.096 g, 21%) as a tan solid mass of suitable purity to use without further purification: <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.04 (s, 1H), 7.91 (d, 1H, J = 2.2 Hz), 7.74 (dd, 1H, J = 8.7, 2.2 Hz), 6.00 (q, 1H, J = 7.1 Hz), 4.42-4.24 (m, 2H), 1.34 (t, 3H, J = 7.3 Hz).

Step 2. Preparation of 6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.


To a stirred solution of the ethyl ester (Step 1) (0.090 g, 0.231 mmol) in THF:EtOH:H<sub>2</sub>O(7:2:1) (4 mL) was added aqueous NaOH solution (0.11 mL, 2.5 M). After stirring 16 hours, the reaction was partially concentrated in vacuo to remove the organic solvents, diluted with H<sub>2</sub>O, and washed with diethyl ether. The resulting aqueous phase was acidified with concentrated HCl, extracted with diethyl ether, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding an oil. The oil was purified by flash chromatography (silica, hexanes-ethyl acetate, 3:1 with 5 % acetic acid). This procedure yielded the title acid (0.020 g, 24 %) as a white powder: mp 162.3-164.7 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.05 (s, 1H), 7.90 (s, 1H), 7.74 (d, 1H, J = 8.7 Hz), 7.29 (d, 1H, J = 8.7 Hz), 5.97 (q, 1H, J = 6.8 Hz). FABLRMS m/z 361 (M-H). ESHRMS m/z 361.0111 (M-H, Calc'd 361.0094).

**EXAMPLE 89**

5    **6-(1,1-Dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**


4-tert-Butylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp  
10    170.6-173.2 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.89 (s, 1H), 7.5-7.4 (m, 2H), 6.93 (d, 1H, J = 8.4 Hz), 5.76 (q, 1H, J = 7.2 Hz), 1.3 (s, 9H). Anal. Calc'd for C<sub>15</sub>H<sub>13</sub>O<sub>3</sub>F<sub>3</sub>: C, 60.00; H, 5.04. Found: C, 59.93; H, 5.12.

15

**EXAMPLE 90**


20    **5-(Hydroxymethyl)-8-methyl-2-(trifluoromethyl)-2H-pyrano[2,3-c]pyridine-3-carboxylic acid**

3-Hydroxymethyl-5-methyl-4-formylpyridine was converted to the title compound by a procedure similar to that described in Example 1: mp 76.1-80.1 °C. <sup>1</sup>H NMR  
25    (acetone-d<sub>6</sub>/300 MHz) 8.15 (s, 2H), 5.93 (q, 1H, J = 7.2 Hz), 1.3 (s, 9H) 5.30 (br s, 1H), 4.79 (br s, 1H), 2.41 (s, 3H). ESHRMS m/z 288.0485 (M+H, Calc'd 288.0483).

**EXAMPLE 91**

5           **2-(Trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid**

4-(Trifluoromethoxy)phenol was converted to 5-(trifluoromethoxy)salicylaldehyde via a procedure similar to that described in Example 2, Step 1. The 5-(trifluoromethoxy)salicylaldehyde was converted to the title compound by a similar procedure to that described in Example 11, Steps 2 & 3: mp 139.1-143.2 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.95 (s, 1H), 7.88 (d, 2H, *J* = 2.4 Hz), 7.71-7.75 (m, 1H), 6.93 (d, 1H, *J* = 8.7 Hz), 5.91 (q, 1H, *J* = 6.9 Hz). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>O<sub>3</sub>F<sub>3</sub>S: C, 41.87; H, 1.76. Found: C, 41.94; H, 1.84.

**EXAMPLE 92**

20           **6-(Trifluoromethyl)-6H-1,3-dioxolo[4,5-g][1]benzopyran-7-carboxylic acid**


25

4-*tert*-Butylphenol was converted to the title compound by a procedure similar to that described in Example 2: mp 245.8-247.8 °C <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.77 (s, 1H), 6.95 (s, 1H), 6.12 (s, 1H), 6.05 (d, 2H, *J* = 0.90 Hz), 5.91

(q, 1H,  $J = 7.2$  Hz). Anal. Calc'd for  $C_{12}H_7O_3F_3$ : C, 50.01; H, 2.45. Found: C, 50.02; H, 2.50.

### EXAMPLE 93

5




#### 8-Ethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

10

2-Ethoxyphenol was converted to 3-ethoxysalicylaldehyde via a procedure similar to that described in Example 11, Step 1. The 3-ethoxysalicylaldehyde was converted to the title compound by a procedure similar to that described in

15 Example 1: mp 159.4-160.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.86 (s, 1H), 6.97-7.14 (m, 3H), 5.83 (q  $H-F$ , 1H,  $J = 7.2$  Hz), 4.12 (q, 2H,  $J = 7.2$  Hz), 1.38 (t, 3H,  $J = 7.2$  Hz). FABHRMS  $m/z$  289.0656 ( $M+H$ , Calc'd 289.0686). Anal. Calc'd for  $C_{13}H_{11}F_3O_4$ : C, 54.17; H, 3.85. Found: C, 54.06; H, 20 3.83.

### EXAMPLE 94




25

#### 6-Chloro-2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

4-Chloro-3-(trifluoromethyl)phenol was converted to the title compound by a procedure similar to that described in Example 11: mp 180.9-182.4 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.96 (s, 1H), 7.84 (s, 1H), 7.47 (s, 1H), 5.96 (q, 1H, *J* = 6.8 Hz), 2.50 (s, 3H). FABLRMS *m/z* 345 (M-H). FABHRMS *m/z* 344.9767 (M-H, Calc'd 344.9753). Anal. Calc'd for C<sub>12</sub>H<sub>5</sub>ClF<sub>3</sub>O<sub>3</sub>: C, 41.58; H, 1.45; Cl, 10.23. Found: C, 41.57; H, 1.50; Cl, 10.33.


10

**EXAMPLE 95**

15 **5-Methoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

6-Methoxysalicylaldehyde was converted to the title compound by a similar procedure to that described in Example 11, Steps 2 & 3: mp 204.5-206.7 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 8.08 (s, 1H), 7.38 (dd, 1H, *J* = 8.5 Hz 8.3 Hz), 6.74 (d, 1H, *J* = 8.5 Hz), 6.65 (d, 1H, *J* = 8.3 Hz), 5.80 (q, 1H, *J* = 7.2 Hz), 3.94 (s, 3H). FABLRMS *m/z* 273 (M-H). EIHRMS *m/z* 274.0444 (M+, Calc'd 274.0453). Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>F<sub>3</sub>O<sub>4</sub>: C, 52.57; H, 3.31. Found: C, 52.47; H, 3.34.

25

**EXAMPLE 96**

6-Benzoyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic  
acid

5 Step 1. Preparation of ethyl 6-benzoyl-2-(trifluoromethyl)-  
2H-1-benzopyran-3-carboxylate.

Ethyl 2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate  
(Example 10, Step 1) (1.59 g, 5.8 mmol) was dissolved in 1,2-  
dichloroethane (3 mL) and added to a 0 °C suspension of  
aluminum chloride (2.59 g, 19.4 mmol) in 1,2-dichloroethane  
10 (3 mL). A solution of benzoyl chloride (1.01 g, 7.2 mmol) in  
1,2-dichloroethane (3 mL) was added and the reaction was  
heated to 80 °C and stirred for 4 hours. The solution was  
poured onto 3 N HCl and ice and extracted with ethyl  
acetate. The ethyl acetate layers were combined, washed  
15 with 3N HCl, saturated sodium bicarbonate, brine, dried over  
MgSO<sub>4</sub> and concentrated in vacuo. The crude ester was  
purified by flash chromatography over silica gel (with 1:9  
ethyl acetate/hexane as eluant) to afford the ester as a  
white crystalline solid (0.26 g, 12%): mp 114.7-116.1 °C. <sup>1</sup>H  
20 NMR (CDCl<sub>3</sub>/300MHz) 7.82 (dd, 1H, J = 8.5 Hz 2.0 Hz), 7.76  
(m, 4H), 7.61 (m, 1H), 7.50 (m, 2H), 7.09 (d, 1H, J = 8.7  
Hz), 5.79 (q, 1H, J = 6.8 Hz), 4.34 (m, 2H), 1.36 (t, 3H, J  
= 7.2 Hz).


25 Step 2. Preparation of 6-benzoyl-2-trifluoromethyl-2H-1-  
benzopyran-3-carboxylic acid.

The ester from Step 1 (0.24 g, 0.64 mmol) was dissolved  
in THF (2 mL) and ethanol (2 mL), treated with 2.5 N sodium  
hydroxide (1.5 mL, 3.8 mmol), and stirred at room  
30 temperature for 4.3 hours. The reaction mixture was  
concentrated in vacuo, acidified with 3N HCl yielding a  
solid. The solid was collected by filtration and was  
recrystallized from ethanol-water to yield a white solid  
(0.14 g, 64%): mp 269.8-270.8 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300  
35 MHz) 8.04 (s, 1H), 7.99 (d, 1H, J = 2.0 Hz), 7.88 (dd, 1H, J  
= 8.5 Hz 2.0 Hz), 7.79 (m, 2H), 7.68 (m, 1H), 7.57 (m, 1H),



7.23 (d, 1H,  $J = 8.6$  Hz), 5.98 (q, 1H,  $J = 7.0$  Hz). FABLRMS  $m/z$  347 (M-H). ESHRMS  $m/z$  347.0560 (M-H, Calc'd 347.0531). Anal. Calc'd for  $C_{18}H_{11}F_3O_4$ : C, 62.08; H, 3.18. Found: C, 61.48; H, 3.22.

5


**EXAMPLE 97**

10     **6-(4-Chlorobenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared analogous to the procedure described in Example 96: mp

15     268.3-269.4 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 8.03 (s, 1H), 7.99 (d, 1H,  $J = 2.0$  Hz), 7.89 (dd, 1H,  $J = 8.5$  Hz, 2.0 Hz), 7.81 (d, 2H,  $J = 8.5$  Hz), 7.62 (d, 2H,  $J = 8.5$  Hz), 7.23 (d, 1H,  $J = 8.5$  Hz), 5.98 (q, 1H,  $J = 7.1$  Hz). FABLRMS  $m/z$  381 (M-H). ESHRMS  $m/z$  381.0135 (M-H, Calc'd 381.0141). Anal.

20     Calc'd for  $C_{18}H_{10}ClF_3O_4$ : C, 56.49; H, 2.63; Cl, 9.26. Found: C, 56.35; H, 2.66; Cl, 9.34.


**EXAMPLE 98**

25

**6-(4-Hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared analogous to the procedure described in Example 96: mp 234.0-239.5 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 8.03 (s, 1H), 7.92 (d, 1H, *J* = 2.0 Hz), 7.83 (dd, 1H, *J* = 8.5 Hz 2.0 Hz), 7.74 (d, 2H, *J* = 8.7 Hz), 7.20 (d, 1H, *J* = 8.5 Hz), 7.00 (d, 1H, *J* = 8.7 Hz), 5.94 (q, 1H, *J* = 7.1 Hz). ESHRMS *m/z* 363.0471 (M-H, Calc'd 363.0480).


### EXAMPLE 99



#### 6-Phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

4-Phenoxyphenol was converted to 5-phenoxysalicylaldehyde by a similar procedure to that described in Example 2, Step 1. 5-Phenoxysalicylaldehyde was converted into the title compound by a similar procedure to that described in Example 11, Steps 2 & 3: mp 184.9-186.4 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.39 (m, 2H), 7.20 (d, 1H, *J* = 2.0 Hz), 7.08 (m, 3H), 7.02 (m, 2H), 5.98 (q, 1H, *J* = 7.2 Hz). FABLRMS *m/z* 335 (M-H). FABHRMS *m/z* 337.0663 (M+H, Calc'd 337.0687). Anal. Calc'd for C<sub>17</sub>H<sub>11</sub>F<sub>3</sub>O<sub>4</sub>: C, 60.72; H, 3.30. Found: C, 60.62; H, 3.29.

### EXAMPLE 100



**8-Chloro-6-(4-chlorophenoxy)-2-trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5    Step 1. Preparation of 5-phenoxy-salicylaldehyde.

        Ethyl magnesium bromide (67.5 mL of an approximately 3.0 M solution in diethyl ether, 202.5 mmol) was added to toluene (50 mL). A solution of 4-phenoxyphenol (25.00 g, 134.26 mmol) in diethyl ether (35 mL) was added resulting in  
10    the evolution of gas. The reaction was heated to 80 °C causing distillation of the diethyl ether. Toluene (300 mL), HMPA (23.4 mL, 24.059 g, 134.26 mmol), and paraformaldehyde (10.07 g, 335.65 mmol) were added and the reaction was heated to 85 °C for 4 hours. The reaction was  
15    cooled to room temperature and was acidified with 2N HCl. The resulting layers were separated and the organic phase collected. The organic phase was washed with brine. The combined aqueous phases were extracted with methylene chloride. The organic phases were combined, dried over  
20    MgSO<sub>4</sub>, filtered and concentrated *in vacuo* yielding a yellow oil. The oil was purified by silica flash chromatography (hexanes-ethyl acetate, 95:5). Concentration *in vacuo* of the desired fractions provided the salicylaldehyde as a pale yellow powder (12.0 g, 42%) of suitable purity to use in  
25    subsequent steps.

Step 2. Preparation of 3-chloro-5-(4-chlorophenoxy)salicylaldehyde.

        To a stirred solution of the salicylaldehyde (Step 1) (  
30    0.981 g, 4.58 mmol) in acetic acid (20 mL) was added chlorine gas via a tube until the yellow color of chlorine persisted. After stirring for four hours at room temperature the reaction was sparged with nitrogen and diluted with water (50 mL). The resulting oily suspension  
35    was extracted with methylene chloride. The methylene chloride phase was washed with sodium bisulfite solution, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*


providing the dichlorinated salicylaldehyde as a yellow oil (0.66 g, 51%) of suitable purity for use in subsequent steps without further purification.

5    Step 3. Preparation of ethyl 8-chloro-6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

A mixture of the dichlorinated salicylaldehyde (Step 2) (0.66 g, 2.3 mmol), triethylamine (0.49 g, 4.8 mmol), ethyl 4,4,4-trifluorocrotonate (0.59 g, 3.5 mmol) in dimethyl  
10    sulfoxide (5 mL) was heated to 85 °C for 3.5 hours. The reaction was allowed to cool to room temperature and was diluted with ethyl acetate (50 mL). The resulting mixture was washed with 3 N HCl (50 mL), aqueous potassium carbonate solution (10 weight %, 2 X 30 mL), and brine. The organic  
15    phase was dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding a brown oil. This oil was purified by flash silica chromatography (hexanes-ethyl acetate, 9:1) providing the substituted 2H-1-benzopyran (0.39 g, 39%) of suitable purity to use in subsequent steps without further  
20    purification.

Step 4. Preparation of 8-chloro-6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

To a solution of the substituted 2H-1-benzopyran ethyl  
25    ester (Step 3) (0.37 g, 0.85 mmol) in ethanol-THF (4 mL, 1:1) was added sodium hydroxide solution (2 mL of 2.5 N, 5 mmol). After stirring for six hours the mixture was concentrated in vacuo. Acidification of the mixture with 3 N HCl yielded a solid which was collected by vacuum filtration. This solid  
30    was recrystallized from ethanol-water yielding the title compound as yellow crystals (0.134 g, 38%): mp 227.8-228.9 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.93 (s, 1H), 7.42 (d, 2H, *J* = 8.9 Hz), 7.24 (s, 2H), 7.12 (d, 2H, *J* = 8.9 Hz), 5.97 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 403 (M-H). FABHRMS *m/z*  
35    405.9790 (M+H, Calc'd 405.9801). Anal. Calc'd for C<sub>17</sub>H<sub>9</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>4</sub> + 2.33% H<sub>2</sub>O: C, 49.22; H, 2.45. Found: C, 49.19; H, 2.27.


**EXAMPLE 101**

5

**2-(Trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid**

4-(4-Trifluoromethylphenyl)phenol was converted to 5-(4-trifluoromethylphenyl)salicylaldehyde via a procedure similar to that described in Example 2, Step 1. The 5-(4-trifluoromethylphenyl)salicylaldehyde was converted to the title compound by a similar procedure to that described in Example 11, Steps 2 & 3: mp 153.5-154.4 °C. <sup>1</sup>H NMR (acetone-  
10 *d*<sub>6</sub>/300 MHz) 7.91 (s, 1H), 7.71 (d, 2H, *J* = 8.9 Hz), 7.33 (s, 1H, *J* = 2.8 Hz), 7.15 (m, 4H), 5.86 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 403 (M-H). ESHRMS *m/z* 403.0399 (M-H, Calc'd 403.0405). Anal. Calc'd for C<sub>18</sub>H<sub>10</sub>F<sub>6</sub>O<sub>4</sub>: C, 53.48; H, 2.49. Found: C, 53.52; H, 2.55.

20

**EXAMPLE 102**


25

**8-(1-Methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

4-(4-Methoxyphenyl)phenol was converted to the title compound by a procedure similar to that  
30 described in Example 2: mp 210.5-211.5 °C. <sup>1</sup>H NMR

(acetone- $d_6$ /300 MHz) 7.86 (s, 1H), 7.35 (d, 1H,  $J$  = 7.7 Hz), 7.28 (s, 1H,  $J$  = 7.5 Hz), 7.04 (t, 1H,  $J$  = 7.7 Hz), 5.85 (q, 1H,  $J$  = 7.2 Hz), 3.33 (sept, 1H,  $J$  = 7.1 Hz), 1.25 (d, 6H,  $J$  = 7.1 Hz). Anal. Calc'd for  $C_{14}H_{13}F_3O_3$ : C, 58.74; H, 4.58. Found: C, 58.65; H, 4.60.


### EXAMPLE 103



#### 6-Chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid

8-(1-Methylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 6) was converted to the title compound by a procedure similar to that described in Example 9. mp 185.4-189.2 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.87 (s, 1H), 7.38 (d, 1H,  $J$  = 2.4 Hz), 7.34 (d, 1H,  $J$  = 2.4 Hz), 5.90 (q, 1H,  $J$  = 7.3 Hz), 3.31 (m, 1H), 1.24 (d, 6H,  $J$  = 6.8 Hz). Anal. Calc'd for  $C_{15}H_{14}ClF_3O_3$ : C, 52.43; H, 3.77; Cl, 11.05. Found: C, 52.58; H, 3.79; Cl, 10.96.

### EXAMPLE 104




**6-(4-Chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared  
 5 6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 99) as the starting material by a procedure similar to that described in Example 9: mp 140.5-142.5 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.39 (d, 2H, *J* = 9.1 Hz), 7.25 (d, 1H, *J* = 2.6 Hz) 7.01-7.15 (m, 4H), 6.85 (q, 1H, *J* = 7.2 Hz). FABLRMS *m/z* 370 (M<sup>+</sup>). ESHRMS *m/z* 369.0130 (M-H, Calc'd 369.0141). Anal. Calc'd for C<sub>17</sub>H<sub>11</sub>ClF<sub>3</sub>O<sub>4</sub> + 0.96% H<sub>2</sub>O: C, 54.55; H, 2.80. Found: C, 54.38; H, 2.80.

**3-(Trifluoromethyl)-2-Hydroxydi**

compound by a procedure similar to that described in Example 99: mp 253.5-254.5 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 8.23 (d, 1H, *J* = 7.5 Hz), 7.50 (m, 1H), 7.25 (d, 1H, *J* = 7.5 Hz). FABLRMS *m/z* 333.0375 (M<sup>+</sup>). Anal. Calc'd 333.0375. Found: C, 52.71; H, 2.71.

**EXAMPLE 105**



**8-Chloro-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid**

20

The benzopyran-3-carboxylic acid was prepared using 6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 101) as the starting material by a similar procedure to that described in Example 99: mp 223.7-226.0 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.74 (d, 2H, *J* = 8.5 Hz), 7.35 (m, 2H), 7.25 (d, 2H, *J* = 8.5 Hz), 6.00 (q, 1H, *J* = 7.0 Hz). FABLRMS *m/z* 437 (M<sup>+</sup>). ESHRMS *m/z* 437.0000 (M-H, Calc'd 437.0015). Anal. Calc'd for C<sub>18</sub>H<sub>9</sub>ClF<sub>5</sub>O<sub>4</sub>: C, 49.28; H, 2.07; Cl, 8.08. Found: C, 49.42; H, 2.12; Cl, 8.17.


**6-Chloro-8-cyano-2H-1-benzopyran-3-carboxylic acid**

**Step 1. Preparation of 6-Chloro-8-cyano-2H-1-benzopyran-3-carboxylic acid**

Hydroxylamine hydrochloride (1.50 g, 15.0 mmol) and sodium acetate (1.50 g, 15.0 mmol) were dissolved in 50 mL of water. A solution of 6-chloro-8-cyano-2H-1-benzopyran-3-carboxylic acid (1.50 g, 5.0 mmol) in 50 mL of water was added to the mixture. The mixture was stirred at room temperature for 24 hours. The mixture was then extracted with ethyl acetate (2 x 50 mL). The organic layers were combined and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to give the product.

ethyl acetate, washed with water, brine, dried over  $\text{MgSO}_4$ , concentrated in vacuo, and recrystallized from diethyl ether/hexane to give a white powder (0.10 g, 21%): mp 238.1-239.7 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.97 (s, 1H), 7.92 (d, 1H,  $J$  = 2.4 Hz), 7.89 (d, 1H,  $J$  = 2.4 Hz), 6.14 (q, 1H,  $J$  = 6.6 Hz). FABLRMS  $m/z$  302 (M-H). ESHRMS  $m/z$  301.9819 (M-H, Calc'd 301.9832). Anal. Calc'd for  $\text{C}_{12}\text{H}_5\text{ClF}_3\text{NO}_3$ : C, 47.47; H, 1.66; N, 4.61. Found: C, 47.41; H, 1.70; N, 4.55.

10

**EXAMPLE 108**

15


**6-Chloro-8-[(hydroxyimino)methyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared from the ethyl ester (Example 107, Step 2) by a method similar to the procedure described in Example 1, Step 2: mp 246.9-247.9 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 10.90 (brs, 1H), 8.35 (s, 1H), 7.92 (s, 1H), 7.78 (d, 1H,  $J$  = 2.6 Hz), 7.61 (d, 1H,  $J$  = 2.6 Hz), 5.98 (q, 1H,  $J$  = 7.0 Hz). FABLRMS  $m/z$  320 (M-H). ESHRMS  $m/z$  319.9959 (M-H, Calc'd 319.9937). Anal. Calc'd for  $\text{C}_{12}\text{H}_4\text{ClF}_3\text{NO}_4$ : C, 44.81; H, 2.19; N, 4.35. Found: C, 44.92; H, 2.25; N, 4.26.

25

**EXAMPLE 109**






**6-Chloro-8-(hydroxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 80 using the carboxylic acid (Example 76, step 4) as the starting material: mp 174.6-178.9 °C. <sup>1</sup>H NMR (acetone-  
10 *d*<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.57 (d, 1H, *J* = 2.6 Hz), 7.47 (d, 1H, *J* = 2.6 Hz), 5.87 (q, 1H, *J* = 7.0 Hz), 4.70 (s, 2H). FABLRMS *m/z* 309 (M+H). ESHRMS *m/z* 306.9981 (M-H, Calc'd 306.9985). Anal. Calc'd for C<sub>12</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>3</sub> (3.81 wt.% H<sub>2</sub>O): C, 47.37; H, 3.08. Found: C, 47.33; H, 2.82.

15

**EXAMPLE 110**



**8-(1H-Benzimidazol-2-yl)-6-chloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

20

Step 1. Preparation of ethyl 8-(1H-Benzimidazol-2-yl)-6-chloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

A solution of the aldehyde (Example 76, Step 3) (0.33 g, 0.99 mmol) and 1,2-phenylenediamine (0.11 g, 1.02 mmol) in nitrobenzene (20 mL) was heated to 150 °C for 1.8 hours. The reaction mixture was extracted with ethyl acetate, washed  
5 with brine, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* and purified by flash chromatography over silica gel (with 1:9 ethyl acetate/hexane as eluant) to give the ester as a brown solid (0.18 g, 43%) which was used in the next step without further purification.


10

Step 2. Preparation of 8-(1H-imidazol-2-yl)-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 1 (0.18 g 1.5 mmol) was dissolved in THF (5 mL) and ethanol (5 mL), treated with 2.5 N sodium  
15 hydroxide (2.6 mL, 6.5 mmol), and stirred at room temperature for 1.7 hours. The reaction mixture was concentrated *in vacuo*, acidified with 3 N HCl, filtered and recrystallized from ethanol-water to give a tan solid (0.09 g, 52%): mp >300 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300MHz) 8.59 (d, 1H,  
20  $J = 2.6$  Hz), 8.03 (s, 1H), 7.73 (d, 1H,  $J = 2.6$  Hz), 7.67 (brs, 2H), 7.28 (m, 2H), 6.13 (q, 1H,  $J = 6.8$  Hz). FAB/MS  $m/z$  395 (M-H( $^{37}\text{Cl}$ )). ESI/MS  $m/z$  393.0262 (M-H, Calc'd 393.0254). Anal. Calc'd for  $\text{C}_{16}\text{H}_{10}\text{ClF}_3\text{N}_2\text{O}_3$  (2.88 wt %  $\text{H}_2\text{O}$ ): C, 53.19; H, 2.80; N, 6.89. Found: C, 53.22; H, 2.90; N, 6.80.

25

**EXAMPLE 111**



30 **7-(1,1-Dimethylethyl)-2-(pentafluoroethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of ethyl 3-hydroxy-4,4,5,5,5-pentafluoropentanoate.

A solution of ethyl 4,4,5,5,5-pentafluoro-3-oxopentanoate (41.32 g, 0.18 mole) in diethyl ether (70 mL) was  
5 cooled to 0 °C and treated with NaBH<sub>4</sub> (7.09 g, 0.19 mole). The reaction was allowed to warm to room temperature and stirred for 2 hours before quenching with 1 N HCl (200 mL). The layers were separated and the aqueous layer was  
10 extracted with diethyl ether. The combined organic layers were washed with 1 N HCl, brine, dried over MgSO<sub>4</sub>, and concentrated in vacuo to give the hydroxy ester as a clear oil (46.40 g) which was used in the next step without further purification.

Step 2. Preparation of ethyl 4,4,5,5,5-pentafluoro-2-pentenoate.

The hydroxy ester from Step 1 (46.40 g, 0.18 mole) was stirred at 120 °C with P<sub>2</sub>O<sub>5</sub> (25.59 g, 0.09 mole) for 2.6  
20 hours then vacuum distilled (95 torr, 45-64 °C) to give the ester as a clear oil (13.70 g, 35%): <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 6.78 (m, 1H), 6.57 (dt, 1H, *J* = 15.9 Hz 2.0 Hz), 4.30 (q, 2H, *J* = 7.3 Hz), 1.34 (t, 3H, *J* = 7.1 Hz).

Step 3. Preparation of ethyl 7-(1,1-Dimethylethyl)-2-(pentafluoroethyl)-2H-1-benzopyran-3-carboxylate.

A mixture of 4-tert-butylsalicylaldehyde Example 8, step 1 (1.15 g, 6.4 mmol) and the ethyl ester from Step 2 (1.59 g, 7.3 mmol) was dissolved in anhydrous DMF (4 mL).  
25 With stirring, K<sub>2</sub>CO<sub>3</sub> (1.10 g, 9.0 mmol) was added causing the reaction to become deep red. The reaction was stirred at room temperature for 100 hours, acidified with 3 N HCl,  
30 diluted with ethyl acetate and washed with saturated NaHCO<sub>3</sub> solution, brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding a brown oil. This oil was purified by  
35 flash chromatography over silica gel, eluting with 10% ethyl acetate/hexanes to afford a yellow oil (1.72 g, 70%): <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.76 (s, 1H), 7.14 (d, 1H, *J* = 8.1 Hz), 7.04


(dd, 1H,  $J = 8.1$  Hz  $1.8$  Hz), 6.94 (s, 1H), 5.92 (dd, 1H,  $J = 22.4$  Hz  $3.0$  Hz), 4.32 (m, 2H), 1.35 (t, 3H,  $J = 7.2$  Hz), 1.30 (s, 9H).

5    Step 4. Preparation of 7-(1,1-Dimethylethyl)-2-(pentafluoroethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 3 (1.58 g 4.20 mmol) was dissolved in THF (3 mL) and ethanol (3 mL), treated with 2.5 N sodium hydroxide (2 mL, 5 mmol), and stirred at room temperature  
10    for 23.3 hours. The reaction mixture was concentrated in vacuo, acidified with 3 N HCl yielding a suspension. The solid was collected by filtration and was recrystallized from ethanol-water to yield a yellow solid (0.76 g, 52%): mp 171.0-173.5 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.93 (s, 1H),  
15    7.39 (d, 1H,  $J = 8.1$  Hz), 7.18 (dd, 1H,  $J = 8.1$  Hz  $1.8$  Hz), 7.02 (s, 1H), 6.01 (dd, 1H,  $J = 23.1$  Hz  $3.2$  Hz), 1.32 (s, 9 H). FABLRMS  $m/z$  351 (M+H). EIHRMS  $m/z$  350.0945 (M+, Calc'd 350.0941). Anal. Calc'd for  $\text{C}_{16}\text{H}_{13}\text{F}_5\text{O}_3$ : C, 54.86; H, 4.32. Found: C, 54.88; H, 4.32.

20

**EXAMPLE 112**



25    **6-Chloro-8-(methoxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

Step 1. Preparation of ethyl 6-chloro-8-(hydroxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

30    A suspension of the aldehyde (Example 75, Step 1) (4.78 g, 14.3 mmol) was cooled to 0 °C and treated with  $\text{NaBH}_4$  (0.33 g, 4.8 mmol). The solution was stirred for 10 minutes then

quenched with 3N HCl, extracted with ethyl acetate, washed with saturated NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, and concentrated in vacuo to give a brown solid which was filtered through a plug of silica gel to give the alcohol as a brown solid (3.60 g, 75%). <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.66 (s, 1H), 7.41 (d, 1H, J = 2.4 Hz), 7.17 (d, 1H, J = 2.4 Hz), 5.75 (q, 1H, J = 6.8 Hz), 4.71 (s, 2H), 4.33 (m, 2H), 1.85 (brs, 1H), 1.36 (t, 3H, J = 7.1). This solid was used in the next step without further purification.

Step 2. Preparation of ethyl 6-chloro-8-(methoxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

The alcohol from Step 1 (0.44 g, 1.3 mmol), silver triflate (0.36 g, 1.4 mmol) and 2,6-di-tert-butylpyridine (0.37 g, 1.9 mmol) were dissolved in methylene chloride (3 mL) cooled to 0 °C and treated with methyl iodide (0.40 g, 2.8 mmol). The reaction was allowed to warm and stirred at room temperature for 4.6 hours. The reaction was filtered through diatomaceous earth and the filtrate was washed with 3N HCl, saturated NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, and concentrated in vacuo yielding a brown oil. This oil was purified by flash chromatography over silica gel, eluting with 10% ethyl acetate-hexanes to afford the substituted 2H-1-benzopyran (0.19 g, 41%) as a white oily solid suitable for use without further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.63 (s, 1H), 7.39 (d, 1H, J = 2.6 Hz), 7.13 (d, 1H, J = 2.6 Hz), 5.72 (q, 1H, J = 6.8 Hz), 4.44 (m, 2H), 4.30 (m, 2H), 3.41 (s, 3H), 1.85 (brs, 1H), 1.33 (t, 3H, J = 7.1).


Step 3. Preparation of 6-chloro-8-(methoxymethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 2 was hydrolyzed via a procedure similar to that described in Example 1, Step 2. mp 166.7-168.0 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.50 (d, 1H, J = 2.6), 7.46 (d, 1H, J = 2.4 Hz), 5.92 (q, 1H, J = 7.1 Hz), 4.49 (s, 2H), 3.42 (s, 3H). FABLRMS m/z 321 (M-H).

ESHRMS  $m/z$  321.0141 (M-H, Calc'd 321.0141). Anal. Calc'd for  $C_{13}H_{10}ClF_3O_4$ : C, 48.39; H, 3.12. Found: C, 48.45; H, 3.11.

### EXAMPLE 113

5



#### 6-Chloro-8-(benzyloxymethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid


10

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 112: mp 133.8-135.4 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.90 (s, 1H), 7.54 (d, 1H,  $J$  = 2.6), 7.51 (d, 1H,  $J$  = 2.4 Hz), 7.42 (m, 5H), 5.91 (q, 1H,  $J$  = 7.1 Hz), 4.68 (s, 2H), 4.63 (s, 2H). FABLRMS  $m/z$  399 (M+H). ESHRMS  $m/z$  397.0454 (M-H, Calc'd 397.0461). Anal. Calc'd for  $C_{19}H_{11}ClF_3O_4$ : C, 57.23; H, 3.54; Cl, 8.89. Found: C, 57.34; H, 3.63; Cl, 8.77.

15

20

### EXAMPLE 114



#### 6-Chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

25

Step 1. Preparation of ethenyl-6-chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

In a 100 mL round bottomed flask under N<sub>2</sub>, ethyl 8-bromo-6-chloro-2-trifluoromethyl-2H-benzopyran-3-carboxylate (Example 74, Step 1) (2.21 g, 5.73 mmol) was dissolved in toluene (30 mL of anhydrous reagent).

- 5 Tetrakis(triphenylphosphine)palladium(0) (0.132 g, 0.115 mmol) was added, followed by tributylethylenylstannane (2.0 g, 6.31 mmol). The resulting solution was heated to reflux for 5 hours. The reaction mixture was allowed to cool to room temperature, was poured into 50 mL of 20% ammonium  
10 fluoride solution and stirred for one hour. Diethyl ether (100 mL) was added and the mixture was washed with water (2 x 50 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered, and evaporated to yield a yellow oil. The crude material was purified by flash chromatography (0.5% ethyl acetate in  
15 hexanes) to afford the ester as a yellow solid (0.86 g, 45%): mp 75.9-77.2 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.64 (s, 1H), 7.45 (d, 1H, J = 2.5 Hz), 7.12 (d, 1H, J = 2.6 Hz), 6.92 (dd, 1H, J = 17.7 Hz, 11.3 Hz), 5.81 (d, 1H, J = 17.7 Hz), 5.76 (q, 1H, J = 6.8 Hz), 5.41 (d, 2H, J = 11.1 Hz), 4.36-  
20 4.29 (m, 2H), 1.36 (t, 3H, J = 7.3 Hz). FABLRMS m/z 350.1 (M+NH<sub>4</sub><sup>+</sup>). ESHRMS m/z 350.0796 (M+NH<sub>4</sub><sup>+</sup>, Calc'd. 350.0771). Anal. Calc'd. for C<sub>15</sub>H<sub>12</sub>ClF<sub>3</sub>O<sub>3</sub> + 4.07% H<sub>2</sub>O: C, 51.95; H, 3.94. Found: C, 51.67; H, 3.69.


25 Step 2. Preparation of 6-chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

- The ester (Step 1) (0.350 g, 1.05 mmol) was dissolved in a solution of THF:ethanol:water(7:2:1; 10 mL), was treated with sodium hydroxide (0.46 mL, 1.05 mmol of a 2.5 N  
30 solution), and stirred at room temperature for 18 hours. The solvent was removed in vacuo and the residue was dissolved in water (10 mL). Diethyl ether (10 mL) was added and the mixture acidified with concentrated HCl. The layers were separated, and the aqueous phase was extracted with  
35 diethyl ether (2 x 10 mL). The organic phases were combined, dried over MgSO<sub>4</sub>, filtered, and evaporated to yield a yellow solid, which was recrystallized in diethyl ether-hexane to afford the title compound as a yellow solid

- (0.288 g, 90%): mp 183.2-185.8 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.77 (s, 1H), 7.49 (d, 1H, *J* = 2.2 Hz), 7.16 (d, 1H, *J* = 2.4 Hz), 6.93 (dd, 1H, *J* = 11.3, 17.7 Hz), 5.82 (d, 1H, *J* = 17.7 Hz), 5.74 (q, 1H, *J* = 6.9 Hz), 5.43 (d, 1H, *J* = 11.1 Hz).
- 5 FABLRMS *m/z* 303 (M-H). ESHRMS *m/z* 303.0014 (M-H, Calc'd. 303.003582). Anal. Calc'd. for C<sub>13</sub>H<sub>6</sub>ClF<sub>3</sub>O<sub>3</sub> + 1.58 % H<sub>2</sub>O: C, 50.44; H, 2.78. Found: C, 50.42; H, 2.65.

### EXAMPLE 115

10




#### 6-Chloro-8-ethynyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

15

- The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 114: mp 186.2-189.0 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.87 (s, 1H), 7.60 (d, 1H, *J* = 2.4 Hz), 7.51 (d, 1H, *J* = 2.4 Hz), 5.95 (q, 1H, *J* = 7.0 Hz), 4.02 (s, 1H). FABLRMS *m/z* 301 (M-H). ESHRMS *m/z* 300.9875 (M-H, Calc'd 300.9879). Anal. Calc'd. for C<sub>13</sub>H<sub>6</sub>ClF<sub>3</sub>O<sub>3</sub>: C, 51.59; H, 2.00; Cl, 11.71. Found: C, 51.26; H, 2.06; Cl, 11.40.

25

### EXAMPLE 116






**6-Chloro-8-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 114: mp 257.5-258.8 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.91 (s, 1H), 7.79 (d, 1H, *J* = 2.4 Hz), 7.74-7.72 (m, 1H), 7.62-7.61 (m, 1H), 7.51 (d, 1H, *J* = 2.4 Hz), 7.19-7.16 (m, 1H), 6.04 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 359 (M-H). ESHRMS *m/z* 358.9747 (M-H, Calc'd. 358.9756). Anal. Calc'd. for C<sub>15</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>3</sub>S: C, 49.94; H, 2.24; Cl, 9.83; S, 8.89. Found: C, 50.26; H, 2.45; Cl, 9.72; S, 9.00.

**EXAMPLE 117**




**6-Chloro-8-(2-furanyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 114: mp 171.5-173.3 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.93 (s, 1H), 7.82 (d, 1H, *J* = 2.6 Hz), 7.72-7.71 (m, 1H), 7.50 (d, 1H, *J* = 2.6 Hz), 7.16 (d, 1H, *J* = 2.4 Hz), 6.65-6.63 (m, 1H), 6.11 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 343 (M-H). ESHRMS *m/z* 342.9995 (M-H, Calc'd. 342.9985). Anal. Calc'd. for C<sub>15</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>4</sub> + 1.31 % H<sub>2</sub>O: C, 51.59; H, 2.46; Cl, 10.15. Found: C, 51.57; H, 2.33; Cl, 10.14.

**EXAMPLE 118**

181



**6-Chloro-8-(5-chloro-1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5


Step 1. Preparation of ethyl 6-chloro-8-(5-chloro-1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

Ethyl 6-chloro-8-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate (Example 73, Step 2) (1.50 g, 3.47 mmol), tetrakis-(triphenylphosphine)palladium(0) (0.2 g, 0.174 mmol), copper(I)iodide (0.066 g, 0.347 mmol), and triethylamine (1.05 g, 10.4 mmol) were dissolved in toluene (50 mL). 5-Chloro-1-pentyne (0.53 g, 5.20 mmol) was added via syringe and the mixture stirred for 18 hours at room temperature. The reaction was diluted with diethyl ether (50 mL), extracted with 0.5 N HCl (2 x 25 mL), and water (2 x 25 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered, and evaporated to yield an orange oil. The crude material was purified by flash chromatography in 2% ethyl acetate in hexane. Recrystallization from hexane afforded the ester as a white solid (0.96 g, 68%): mp 84.8-85.9 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.61 (s, 1H), 7.33 (d, 1H, J = 2.6 Hz), 7.14 (d, 1H, J = 2.6 Hz), 5.79 (q, 1H, J = 6.7 Hz), 4.37-4.29 (m, 2H), 3.75 (t, 2H, J = 6.7 Hz), 2.67 (t, 2H, J = 6.7 Hz), 2.11-2.03 (m, 2H), 1.35 (t, 3H, J = 7.2 Hz). FAB/MS m/z 424.1 (M+NH<sub>4</sub><sup>+</sup>). ESI/MS m/z 424.0694 (M+NH<sub>4</sub><sup>+</sup>, Calc'd. 424.0694). Anal. Calc'd. for C<sub>18</sub>H<sub>15</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 53.09; H, 3.71; Cl, 17.41. Found: C, 53.02; H, 3.90; Cl, 17.63.

Step 2. Preparation of 6-chloro-8-(5-chloro-1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

The ester (Step 1) (0.500 g, 1.23 mmol) was dissolved in THF-ethanol-water(7:2:1; 10 mL). It was treated with sodium hydroxide (0.49 mL, 1.23 mmol of a 2.5 N solution), and stirred at room temperature for 18 hours. The solvent was evaporated and the residue was dissolved in water (10 mL). Diethyl ether (10 mL) was added and the mixture acidified with concentrated HCl. The organic layer was separated, and the aqueous phase was extracted with diethyl ether (2 x 10 mL). The combined extracts were dried over MgSO<sub>4</sub>, filtered, and evaporated to yield a yellow solid, which was recrystallized in diethyl ether-hexane to afford the title compound as a yellow solid (0.371 g, 80%): mp 154.4-156.4 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.88 (s, 1H), 7.53 (d, 1H, J = 2.4 Hz), 7.44 (d, 1H, J = 2.4 Hz), 5.94 (q, 1H, J = 7.1 Hz), 3.83 (t, 2H, J = 6.5 Hz), 2.68 (t, 2H, J = 6.8 Hz), 2.12-2.04 (m, 2H). ESLRMS m/z 377 (M-H). ESHRMS m/z 376.9930 (M-H, Calc'd. 376.9959). Anal. Calc'd. for C<sub>16</sub>H<sub>11</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub> + 1.18 % H<sub>2</sub>O: C, 50.08; H, 3.02; Cl, 18.48. Found: C, 50.11; H, 2.73; Cl, 18.28.

### EXAMPLE 119




### 25 6-Chloro-8-(1-pentynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 118: mp 168.1-171.2 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.75 (s, 1H), 7.37 (d, 1H, J = 2.6 Hz), 7.15 (d, 1H, J = 2.4 Hz), 5.77 (1, 1H,

$J = 6.7$  Hz), 2.44 (t, 2H,  $J = 6.9$  Hz), 1.68-1.61 (m, 2H), 1.07 (t, 3H,  $J = 7.25$  Hz). FABLRMS  $m/z$  345 (M+H). ESHRMS  $m/z$  343.0373 (M-H, Calc'd, 343.0349). Anal. Calc'd. for  $C_{16}H_{12}ClF_3O_3 + 0.69\% H_2O$ : C, 55.36; H, 3.56. Found: C, 55.21; H, 3.62.


### EXAMPLE 120



#### 6-Chloro-8-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 118: mp 190.1-192.1 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 7.92 (s, 1H), 7.61-7.57 (m, 4H), 7.47-7.44 (m, 3H), 6.01 (q, 1H,  $J = 7.0$  Hz). ESHRMS  $m/z$  377 (M-H). ESHRMS  $m/z$  377.0167 (M-H, Calc'd, 377.0192). Anal. Calc'd. for  $C_{19}H_{10}ClF_3O_3$ : C, 60.26; H, 2.66; Cl, 9.36. Found: C, 60.09; H, 2.73; Cl, 9.09.


### EXAMPLE 121



**6-Chloro-8-(3,3-dimethyl-1-butyryl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a  
5 procedure similar to the method described in Example 118: mp  
218.3-222.4 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.87 (s, 1H),  
7.51 (d, 1H, J = 2.4 Hz), 7.38 (d, 1H, J = 2.6 Hz), 5.92 (q,  
1H, J = 6.9 Hz), 1.32 (s, 9H). FABLRMS m/z 359 (M+H).  
ESHRMS m/z 357.0490 (M-H, Calc'd. 357.0505). Anal. Calc'd.  
10 for C<sub>17</sub>H<sub>14</sub>ClF<sub>3</sub>O<sub>3</sub>: C, 56.92; H, 3.93; Cl, 9.88. Found: C,  
56.63; H, 3.94; Cl, 10.03.

**EXAMPLE 122**




15

**6-Chloro-8-[(4-chlorophenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

20 The 2H-1-benzopyran-3-carboxylic acid was prepared by a  
procedure similar to the method described in Example 118: mp  
210.4-211.4 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.75 (s, 1H), 7.48-  
7.43 (m, 3H), 7.36 (s, 1H), 7.33 (s, 1H), 7.22 (d, 1H, J =  
2.6 Hz), 5.82 (q, 1H, J = 6.6 Hz). FABLRMS m/z 411 (M-H).  
25 ESHRMS m/z 410.9802 (M-H, Calc'd. 410.980259). Anal.  
Calc'd. for C<sub>20</sub>H<sub>12</sub>Cl<sub>2</sub>F<sub>3</sub>O<sub>3</sub>: C, 55.23; H, 2.20; Cl, 17.16.  
Found: C, 55.22; H, 2.07; Cl, 17.39.


**EXAMPLE 123**



**6-Chloro-8-[(4-methoxyphenyl)ethynyl]-2-(trifluoromethyl)-  
2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 118: mp 217.7-218.7 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.75 (s, 1H), 7.51-7.47 (m, 3H), 7.18 (d, 1H, *J* = 2.4 Hz), 6.91-6.88 (m, 2H), 5.82 (t, 1H, *J* = 6.7 Hz). ESI/MS *m/z* 407 (M-H). ESI/MS *m/z* 407.0293 (M-H, Calc'd 407.0298). Anal. Calc'd for C<sub>20</sub>H<sub>12</sub>ClF<sub>3</sub>O<sub>4</sub>: C, 58.77; H, 2.96; Cl, 8.67. Found: C, 58.68; H, 2.85; Cl, 9.15.

**EXAMPLE 124**




**6-(Phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 118 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 24, Step 3) as the starting material:

mp 240.1-241.3 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.70-7.69 (m, 1H), 7.61-7.53 (m, 3H), 7.44-7.41 (m, 3H), 7.10 (d, 1H, *J* = 7.1 Hz). ESHRMS *m/z* 343.0550 (M-H, Calc'd. 343.0582). Anal. Calc'd. for C<sub>19</sub>H<sub>11</sub>F<sub>3</sub>O<sub>3</sub>: C, 66.29; H, 3.22. Found: C, 66.26; H, 3.29.

### EXAMPLE 125



#### 6-Chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

##### Step 1. Preparation of ethyl 6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

Ethyl 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 73, Step 2) (1.3 g, 3.02 mmol), potassium carbonate (1.25 g, 9.06 mmol), 4-chlorophenylboronic acid (0.52 g, 3.33 mmol), and tetrakis(triphenylphosphine)palladium(0) (0.174 g, 0.151 mmol) were added to toluene (30 mL) and the resulting solution was heated to reflux for 18 hours. After cooling to room temperature the reaction mixture was poured into ethyl acetate (50 mL). It was washed with 1 N HCl (2 x 25 mL), saturated aqueous sodium bicarbonate (2 x 25 mL), and water (2 x 25 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo to yield a brown oil. The crude material was purified by flash chromatography using 1% ethyl acetate in hexane yielding a white solid. Recrystallization from hexane afforded the ester as a white solid (0.79 g, 64%): mp 114.2-115.9 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300


- MHz) 7.69 (s, 1H), 7.41 (s, 4H), 7.30 (d, 1H,  $J = 2.4$  Hz), 7.22 (d, 1H,  $J = 2.6$  Hz), 5.70 (q, 1H,  $J = 6.9$  Hz), 4.37-4.29 (m, 2H), 1.35 (t, 3H,  $J = 7.1$  Hz). ES LRMS  $m/z$  434 ( $M+NH_4^+$ ). FAB HRMS  $m/z$  434.0574 ( $M+NH_4^+$ , Calc'd. 434.0538).
- 5 Anal. Calc'd. for  $C_{19}H_{13}Cl_2F_3O_3$ : C, 54.70; H, 3.14; Cl, 17.00. Found: C, 54.79; H, 3.18; Cl, 16.65.

Step 2. Preparation of 6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

- 10 The ester from Step 1 (0.500 g, 1.20 mmol) was dissolved in a solution of THF:ethanol:water (7:2:1; 10 mL), treated with sodium hydroxide (0.48 mL, 1.20 mmol of a 2.5 N solution), and stirred at room temperature for 18 hours. The solvent was removed in vacuo and the residue was
- 15 dissolved in water (10 mL). Diethyl ether (10 mL) was added and the mixture acidified with concentrated HCl. The organic layer was separated, and the aqueous phase was extracted with diethyl ether (2 x 10 mL). The combined
- 20 extracts were dried over  $MgSO_4$ , filtered, and evaporated to yield a white solid, which was recrystallized in diethyl ether-hexane to afford the title compound as a white solid (0.40 g, 86%): mp 205.5-207.3 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 7.81 (s, 1H), 7.42 (s, 4H), 7.34 (d, 1H,  $J = 2.4$  Hz), 7.25 (s, 1H), 5.69 (q, 1H,  $J = 6.8$  Hz). FAB LRMS  $m/z$  387 (M-H).
- 25 ESHRMS  $m/z$  386.9788 (M-H, Calc'd. 386.980259). Anal. Calc'd. for  $C_{17}H_9Cl_2F_3O_3$ : C, 52.47; H, 2.33; Cl, 18.22. Found: C, 52.38; H, 2.47; Cl, 18.20.

**EXAMPLE 126**

30





**6-Chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5     Step 1. Preparation of ethyl 6-chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.


          In a 100 mL round bottomed flask under nitrogen, ethyl 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 73, Step 2) (1.00 g, 2.31 mmol) and 3-methoxyphenylboronic acid (0.369 g, 2.43 mmol) were  
10   dissolved in 1-propanol (50 mL). The mixture was stirred at room temperature for 0.5 hours, allowing for the solids to dissolve. The resulting solution was treated with palladium (II) acetate (0.016 g, 0.0693 mmol), triphenylphosphine (0.055 g, 0.208 mmol), sodium carbonate (0.294 g, 2.77  
15   mmol), and deionized water (10 mL). The reaction mixture was heated to reflux for 3 hours. After cooling to room temperature the mixture was extracted with ethyl acetate (1 X 150 mL, 2 x 25 mL). The combined organic phases were washed with saturated aqueous NaHCO<sub>3</sub> (50 mL) and brine (2 x  
20   50 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo to yield a yellow oil. The crude material was purified by flash chromatography in 0.5% ethyl acetate in hexane yielding a white solid. The solid was recrystallized from hexane yielding the desired ester as a white solid  
25   (0.60 g, 63%): mp 93.7-95.1 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.69 (s, 1H), 7.35-7.32 (m, 2H), 7.22 (d, 1H, J = 2.6 Hz), 7.05-7.03 (m, 2H), 6.96-6.93 (m, 1H), 5.72 (q, 1H, J = 6.7 Hz), 4.34-4.31 (m, 2H), 1.35 (t, 3H, J = 7.1 Hz). FABLRMS m/z 413 (M+H). ESHRMS m/z 413.0765 (M+H, Calc'd. 413.076747).  
30   Anal. Calc'd. for C<sub>20</sub>H<sub>18</sub>ClF<sub>3</sub>O<sub>4</sub>: C, 58.19; H, 3.91; Cl, 8.59. Found: C, 58.33; H, 4.10; Cl, 8.61.

35     Step 2. Preparation of 6-chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.

          The ester from Step 1 (0.300 g, 0.727 mmol) was dissolved in THF-ethanol-water (7:2:1, 10 mL). It was treated with sodium hydroxide (0.29 mL of a 2.5 N solution, 0.727 mmol), and stirred at room temperature for 18 hours.

The solvent was evaporated and the residue was dissolved in water (10 mL). Ether (10 mL) was added, followed by a few drops of concentrated HCl. The ether layer was separated, and the aqueous phase was extracted with ether (2 x 10 mL).  
5 The ether extracts were combined, dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo to yield a white solid, which was recrystallized in diethyl ether-hexane to afford the title compound as a white solid (0.23 g, 81%): mp 173.1-177.4 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ /300 MHz) 7.81 (s, 1H), 7.39-7.37 (m, 2H), 7.05-7.04 (m, 2H), 6.97-6.94 (m, 1H), 5.71 (q, 1H,  $J = 6.7$  Hz), 3.85 (s, 3H). ESHRMS  $m/z$  383.0278 (M-H, Calc'd. 383.029796). Anal. Calc'd. for  $\text{C}_{18}\text{H}_{12}\text{ClF}_3\text{O}_4$ : C, 56.20; H, 3.14; Cl, 9.21. Found: C, 55.90; H, 3.11; Cl, 9.48.

15

**EXAMPLE 127**

20 **6-Chloro-8-[(4-methylthio)phenyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 126: mp 211.4-212.5 °C  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 7.94 (s, 1H), 7.57 (d, 1H,  $J = 2.6$  Hz), 7.53-7.50 (m, 2H), 7.45 (d, 1H,  $J = 2.6$  Hz), 7.39-7.36 (m, 2H), 5.87 (q, 1H,  $J = 7.1$  Hz), 2.55 (s, 3H). ESHRMS  $m/z$  399.0051 (M-H, Calc'd. 399.0069). Anal. Calc'd. for  $\text{C}_{18}\text{H}_{12}\text{ClF}_3\text{O}_3\text{S}$ : C, 53.94; H, 3.02; Cl, 8.84; S, 8.00. Found: C, 53.86; H, 2.82; Cl, 8.91; S, 8.21.


30

**EXAMPLE 128**



The sulfone from Step 1 (0.300 g, 0.651 mmol) was dissolved in a solution of THF:ethanol:water (7:2:1; 10 mL). It was treated with sodium hydroxide (0.26 mL, 0.651 mmol of a 2.5 N solution), and stirred at room temperature for 18 hours. The solvent was removed in vacuo and the residue was dissolved in water (10 mL). Diethyl ether (10 mL) was added and the mixture acidified with concentrated HCl. The organic layer was separated, and the aqueous phase was extracted with diethyl ether (2 x 10 mL). The combined organic extracts were dried over MgSO<sub>4</sub>, filtered, and evaporated to yield a white solid. Recrystallization of this solid in ether-hexane afforded the title compound as a white solid (0.20 g, 73%): mp 286.5-287.8 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.07 (d, 2H, J = 6.7 Hz), 7.97 (s, 1H), 7.84 (d, 2H, J = 6.7 Hz), 7.67 (d, 1H, J = 2.6 Hz), 7.55 (d, 1H, J = 2.6 Hz), 5.92 (q, 1H, J = 7.1 Hz), 3.20 (s, 1H). ESHRMS m/z 430.9947 (M-H, Calc'd. 430.996782). Anal. Calc'd. for C<sub>18</sub>H<sub>12</sub>ClF<sub>3</sub>O<sub>3</sub>S: C, 49.95; H, 2.80; Cl, 8.19. Found: C, 50.04; H, 2.80; Cl, 8.25.

### EXAMPLE 129



### 25      6-Chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid


#### Step 1. Preparation of ethyl 6-chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

30      A mixture of ethyl 6-chloro-8-bromo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 74, Step 1) (2.0 g, 5.2 mmol), tetrakis(triphenylphosphine)

palladium(0) (2.15 g, 1.7 mmol), triphenylphosphine (0.013 g, 0.05 mmol), and tributylphenyltin (1.9 mL, 5.7 mmol) in toluene (60 mL) was heated to 110 °C for 3 days. The reaction mixture was allowed to cool to room temperature and  
5 filtered through a plug of silica gel eluting with 25% ethyl acetate in hexanes. The filtrate was concentrated *in vacuo* and then purified by flash chromatography (silica gel, ethyl acetate-hexanes, 1:9). The fractions containing desired product were combined and concentrated *in vacuo*. To remove  
10 the remaining tin impurities the mixture was taken up in THF (10 mL) and aqueous ammonium fluoride solution (10 wt %, 20 mL) and stirred at room temperature for 2 hours. The solution was extracted with ethyl acetate. The extracts were combined, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to afford the ester as an oil (1.30 g, 65%). <sup>1</sup>H  
15 NMR(CDCl<sub>3</sub>/300 MHz) 7.67 (s, 1H), 7.47-7.36 (m, 5H), 7.31 (d, 1H, *J* = 2.6 Hz), 7.18 (d, 1H, *J* = 2.4 Hz), 5.69 (q, 1H, *J* = 6.8 Hz), 4.30 (m, 2H), 1.33 (t, 3H, *J* = 7.1 Hz). <sup>19</sup>F NMR (CDCl<sub>3</sub>/282 MHz) d -78.27 (d, *J* = 7.2 Hz). FABLRMS *m/z* 383 (M+H). ESHRMS *m/z* 400.0937 (M+NH<sub>4</sub><sup>+</sup>, Calc'd 400.0927)  
20

Step 2. Preparation of 6-chloro-8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

A solution of the ester from step 1 (1.0 g, 2.6 mmol)  
25 was dissolved in THF (5 mL) and methanol (5 mL) was treated with a 2.5 N NaOH solution (4.0 mL, 10.4 mmol). The resulting mixture was stirred at room temperature for 18 hours. The solvent was removed *in vacuo*, and the residue taken up in ethyl acetate and acidified with 3 N HCl. The  
30 solution was extracted with ethyl acetate. The extracts were combined, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* yielding a yellow solid. Recrystallization from ethyl acetate-hexanes afforded the title compound as a pale yellow solid (0.42 g, 46%): mp 196.3-197.7 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300  
35 MHz) d 7.65 (s, 1H), 7.40-7.23 (m, 6H), 7.15 (s, 1H), 5.63 (q, 1H, *J* = 6.5 Hz), 3.35 (broad s, 1H). <sup>19</sup>F NMR (CDCl<sub>3</sub>/282 MHz) d -78.71 (d, *J* = 5.8 Hz). FABLRMS *m/z* 355 (M+H). ESHRMS *m/z* 353.0198 (M-H, Calc'd 353.0192).


**EXAMPLE 130**

5

**6-Bromo-8-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

4-Bromo-2-fluorophenol was converted to the title  
10 compound by a procedure similar to that described in Example  
2: mp 206-208 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.78 (s, 1H), 7.36-  
7.48 (m, 2H), 5.87 (q, 1H, *J* = 6.8 Hz). EIHRMS *m/z* 339.9349  
(Calc'd 339.9358). Anal. Calc'd for C<sub>11</sub>H<sub>5</sub>BrF<sub>2</sub>O<sub>3</sub>: C 38.74, H  
1.48; Found C 38.97, H, 1.60.

15


**EXAMPLE 131****6-(4-Fluorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a  
procedure similar to the method described in Example 125  
25 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylate (Example 24, Step 3) as the starting material:  
mp 207-210 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.87 (s, 1H), 7.54-  
7.64 (m, 4H), 7.10-7.20 (m, 2H), 7.03 (d, 1H, *J* = 9.4 Hz),  
5.77 (q, 1H, *J* = 7.0 Hz). EIHRMS *m/z* 338.0573 (Calc'd

338.0566) Anal. Calc'd for  $C_{11}H_6F_3IO_3 + 1.25\% H_2O$ : C, 59.62; H, 3.08. Found C, 59.61; H, 3.09.

### EXAMPLE 132

5



#### 6-Phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid


10

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 125 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 24, Step 3) as the starting material:

15 mp 197-198 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.87 (s, 1H), 7.28-7.64 (m, 7H), 7.03 (d, 1H,  $J = 6.8$  Hz), 5.76 (q, 1H,  $J = 7.0$  Hz). EIHRMS  $m/z$  320.0604 ( $M^+$ , Calc'd 320.0660). Anal. Calc'd for  $C_{17}H_{11}F_3O_3$ : C, 63.75; H 3.46. Found C, 63.56; H, 3.46.

20

### EXAMPLE 133




#### 8-Chloro-6-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

2-Chloro-4-fluorophenol was converted to the title compound by a procedure similar to that described in Example 2:  
25 2: mp 240-241 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 7.77 (s, 1H), 7.26  
30

(dd, 1H,  $J = 8.3, 2.9$ ), 7.14 (dd, 1H,  $J = 8.1, 2.9$ ), 5.87 (q, 1H,  $J = 6.8$  Hz). EIHRMS  $m/z$  295.9836 (Calc'd 295.9863). Anal. Calc'd for  $C_{11}H_5ClF_4O_3$ : C, 44.54; H, 1.70. Found C, 44.70; H, 1.73.


5

**EXMAPLE 134**

10 **6,8-Diiodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 1: mp 243-244 °C.  $^1H$  NMR ( $CD_3OD/300$  MHz) 8.07 (d, 1H,  $J = 2.0$  Hz), 7.71 (s, 1H), 7.70 (d, 1H,  $J = 2.0$  Hz), 5.89 (q, 1H,  $J = 6.8$  Hz). ESHRMS  $m/z$  494.8174 (Calc'd for M-H 494.8202) Anal. Calc'd for  $C_{11}H_5F_3I_2O_3$ : C, 26.64; H, 1.02. Found C, 26.75; H, 1.06.

20

**EXAMPLE 135**

25 **6-(5-Chloro-2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 125 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-


30



carboxylate (Example 72, Step 3) as the starting material:  
mp 205-206 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.83 (s, 1H), 7.50-  
7.58 (m, 2H), 7.14 (d, 1H, *J* = 4.0 Hz), 7.00 (d, 1H, *J* =  
8.86 Hz), 6.93 (d, 1H, *J* = 4.0 Hz), 5.77 (q, 1H, *J* = 7.0  
5 Hz). EIHRMS *m/z* 359.9810 (M<sup>+</sup>, Calc'd 359.9835). Anal.  
Calc'd for C<sub>15</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>3</sub>S: C, 49.94; H 2.24. Found C, 50.14; H,  
2.29.

### EXAMPLE 136

10




#### 6-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

15

The 2H-1-benzopyran-3-carboxylic acid was prepared by a  
procedure similar to the method described in Example 125  
using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-  
carboxylate (Example 24, Step 3) as the starting material:  
20 mp 209-212 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.83 (s, 1H), 7.58-  
7.62 (m, 2H), 7.30-7.38 (m, 2H), 6.80-7.09 (m, 2H), 5.76 (q,  
1H, *J* = 7.0 Hz) FABHRMS *m/z* 325.0153 (Calc'd for M-H  
325.0146)

25

### EXAMPLE 137




#### 6-(4-Chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

30

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 125 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 24, Step 3) as the starting material:  
mp 212-213 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.89 (s, 1H), 7.56-7.66 (m, 4H), 7.40-7.48 (m, 2H), 7.04-7.10 (m, 1H), 5.77 (q, 1H, *J* = 7.0 Hz). ESHRMS *m/z* 353.0190 (Calc'd for M-H 353.0192). Anal. Calc'd for C<sub>17</sub>H<sub>10</sub>ClF<sub>3</sub>O<sub>3</sub>: C, 57.56; H, 2.84.  
Found C, 57.41; H, 2.82.

### EXAMPLE 138




15

#### 6-(4-Bromophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 126: using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 24, Step 3) as the starting material:  
mp 215-216 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.89 (s, 1H), 7.06-7.71 (m, 6H), 7.04-7.06 (m, 1H), 5.78 (q, 1H, *J* = 6.8 Hz). ESHRMS *m/z* 396.9681 (Calc'd for M-H 396.9687).

25

### EXAMPLE 139




**6-(Ethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

5        The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 118 using ethyl 6-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 24, Step 3) as the starting material: mp 198-200 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.80 (s, 1H), 7.47 (dd, 1H, *J* = 8.5, 2.0 Hz), 7.41 (d, 1H, *J* = 2.0 Hz), 6.97 (d, 1H, *J* = 8.5 Hz), 5.71 (q, 1H, *J* = 6.8 Hz), 3.06 (s, 1H). ESHRMS *m/z* 267.0271 (Calc'd for M-H 267.0269) Anal. Calc'd for C<sub>13</sub>H<sub>7</sub>F<sub>3</sub>O<sub>3</sub> + 1.06% H<sub>2</sub>O: C, 57.60; H, 2.72. Found C, 57.59; H, 2.62.

15

**EXAMPLE 140**




20        **6-Methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

      4-Methylsalicylaldehyde was converted to the title compound by a procedure similar to that described in Example 1: mp 191.8-193.0 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.80 (s, 1H), 7.72-7.73 (m, 2H), 6.90 (d, 1H, *J* = 8.4 Hz), 5.91 (q, 1H, *J* = 7.2 Hz). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>O<sub>3</sub>F<sub>3</sub>: C, 55.82; H, 3.51. Found: C, 55.89; H, 3.49.

30

**EXAMPLE 141**




**6-Chloro-8-(4-methoxyphenyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid**

5

The 2H-1-benzopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 126: mp 194.0-196.0 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.81 (s, 1H), 7.44 (s, 1H), 7.41 (s, 1H), 7.34 (d, 1H, *J* = 2.4 Hz), 7.21 (d, 1H, *J* = 2.4 Hz), 6.99 (s, 1H), 6.96 (s, 1H), 5.69 (q, 1H, *J* = 6.7 Hz), 3.86 (s, 3H). FABLRMS *m/z* 402.2 (M+NH<sub>4</sub>). ESHRMS *m/z* 383.0267 (M-H, Calc'd. 383.029796). Anal. Calc'd. for C<sub>18</sub>H<sub>12</sub>ClF<sub>3</sub>O<sub>4</sub>: C, 56.20; H, 3.14; Cl, 9.21. Found: C, 56.08; H, 3.11; Cl, 9.13.

15

**EXAMPLE 142**



**6-Chloro-2-(trifluoromethyl)-4-ethenyl-2H-1-benzopyran-3-carboxylic acid**

Step 1: Preparation of ethyl 3-(5-chloro-2-hydroxyphenyl)-3-oxo-propionate

A solution of lithium hexamethyldisilazide (800 mL of 1.0 M solution in THF, 800.0 mmol) was chilled to -78 °C under a nitrogen atmosphere. A solution of 5-chloro-2-hydroxyacetophenone (45.493 g, 266.67 mmol) in THF (130 mL)

was added dropwise to the stirred solution over 0.5 hour. The reaction was held at -78 °C for 1 hour, warmed to -10 °C for 2 hours, warmed to 0 °C for 1 hour, then cooled to -78 °C. Diethyl carbonate (35.54 mL, 34.65 g, 29.34 mmol) was added via syringe in one portion. The temperature was maintained at -78 °C for 0.5 hour, warmed to room temperature over 0.5 hour, and stirred for 3 hours. The crude reaction mixture was carefully poured over a mixture of rapidly stirred ice (1200 mL)/conc HCl (222 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate. The combined organic phase was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo yielding an oil that began to crystallize. Hexanes (150 mL) was added and crystallization proceeded. The crystalline product was collected by vacuum filtration to afford the title compound (29.04 g, 45%) as tan crystalline needles: mp 71.8-73.1 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.63 (d, 1H, *J* = 2.4 Hz), 7.45 (dd, 1H, *J* = 8.9, 2.6), 6.98 (d, 1H, *J* = 8.9 Hz), 4.25 (q, 2 H, *J* = 7.3 Hz), 3.98 (s, 2H), 1.29 (t, 3H, 7.3 Hz). FABLRMS *m/z* 249 (M+Li). EIHRMS *m/z* 242.0346 (M+, Calc'd 242.0346). Anal. Calc'd for C<sub>11</sub>H<sub>11</sub>ClO<sub>4</sub>: C, 54.45; H, 4.57. Found: C, 54.48; H, 4.62.

Step 2. Preparation of ethyl 2-(trifluoromethyl)-6-chloro-4-oxo-4H-1-benzopyran-3-carboxylate.

The keto-ester (Step 1) (19.2 g, 79.1 mmol), was added to trifluoroacetic anhydride (67.2 mL, 49.9 g, 475.8 mmol), potassium carbonate (44 g, 318 mmol) and toluene (400 mL). This suspension was stirred at room temperature for 36 hours, then heated to reflux for 4 hours. After cooling to room temperature, the suspension was poured over rapidly stirred (mechanical stirrer) ice (300 mL) and aqueous HCl (12 N, 50 mL). The resulting organic phase was separated from the clear mixture, was washed with water ( 5 X 500 mL), brine (1 X 500 mL), dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding tan solid which was dried under high vacuum. This sample was partially dissolved in

heptane (100 mL) and ethyl acetate (12 mL) with heating on a steam bath, was filtered to remove insoluble material. The filtrate was allowed to cool to room temperature yielding the desired 4-oxo-4H-1-benzopyran as a fluffy tan solid  
5 (14.17 g, 56%): mp 106.7-108.6 °C. This material was of suitable purity to use in the next step without further purification.

10 Step 3. Preparation of ethyl 2-(trifluoromethyl)-4-oxo-dihydro-1-benzopyran-3-carboxylate.

A stirred, chilled (0 °C) solution of the ketone (Step 2) (6.92 g, 21.58 mmol) in tetrahydrofuran (40 mL) and ethanol (50 mL) was treated portion-wise with sodium borohydride (NaBH<sub>4</sub>, 0.41 g, 10.79 mmol). After 3 h  
15 additional sodium borohydride (0.30 g, 7.93 mmol) was added portion-wise over 1 hour. The reaction was poured into rapidly stirred cold aqueous HCl (15 mL of 12 N HCl diluted to 300 mL). During the addition a precipitate formed, that was collected by vacuum filtration and dried under high  
20 vacuum yielding the desired substituted 4-oxo-dihydro-1-benzopyran as a white powder (6.92 g, 99%): mp 80.2-84.9 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 12.60 (br s, 1H), 7.69 (d, 1H, J = 2.6 Hz), 7.34 (dd, 1H, J = 2.6, 8.7 Hz), 6.93 (d, 1H, J = 8.7 Hz), 5.59 (q, 1H, 6.6 Hz), 4.46-4.23 (m, 2H), 1.35 (t,  
25 3H, J = 7.0 Hz). FABLRMS m/z 329 (M+Li). EIHRMS m/z 322.0213 (M+, Calc'd 322.0220). Anal. Calc'd for C<sub>13</sub>H<sub>9</sub>ClF<sub>3</sub>O<sub>4</sub> with 3.57% water: C, 46.67; H, 3.41. Found: C, 46.62; H, 3.14.

30 Step 4. Preparation of ethyl 6-chloro-4-(trifluoromethanesulfonyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

A 50 mL Morton flask fitted with septa and addition funnel was charged with 2,6-di-tert-butylpyridine (1.782 g,  
35 8.679 mmol), methylene chloride (15 mL), and trifluoromethanesulfonic anhydride (1.22 mL, 2.04 g, 7.23 mmol) followed by the dropwise addition of the chroman-4-one

- (Step 3) (2.145 g, 5.786 mmol) in methylene chloride (12 mL) over 0.33 hour. After stirring for 16 h at room temperature, the reaction was concentrated in vacuo and diluted with diethyl ether (50 mL) yielding a suspension.
- 5 The suspension was vacuum filtered and the filtrate washed with cold 2 N HCl and brine, dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo yielding the desired triflate as a light yellow powder (1.45 g, 55%) of suitable purity to use without further purification: mp 79.2-80.4 °C. <sup>1</sup>H NMR
- 10 (CDCl<sub>3</sub>/300 MHz) 7.40 (s, 1H), 7.37 (d, 1H, *J* = 2.4 Hz), 7.02-6.99 (m, 1H), 5.92 (q, 1H, *J* = 6.6 Hz), 4.47-4.32 (m, 2H), 1.39 (t, 3H, *J* = 7.2 Hz).

Step 5. Preparation of ethyl 6-chloro-4-ethenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

- Ethyl 6-chloro-4-trifluoromethanesulfoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Step 4) (1.50 g, 3.30 mmol) was dissolved in anhydrous THF (40 mL) in a 100 mL round bottomed flask under nitrogen.
- 20 Tetrakis(triphenylphosphine)palladium(0) (0.267 g, 0.231 mmol) and lithium chloride (0.140 g, 3.3 mmol) were added, followed by tributylethenylstannane (1.15 g, 3.6 mmol). The resulting solution was heated to reflux for 18 hours. GCMS analysis indicated the starting material had been consumed.
- 25 The reaction mixture was allowed to cool to room temperature and was poured into 20% ammonium fluoride solution (50 mL). After stirring for one hour, diethyl ether (100 mL) was added and the mixture was washed with water (2 x 50 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered, and
- 30 concentrated in vacuo yielding a brown oil. The crude material was purified by flash column chromatography (hexane) to afford the ester as a yellow oil, which crystallized upon standing (0.760 g, 69%): mp 51.9-53.2 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.46 (d, 1H, *J* = 2.4 Hz), 7.28-7.14
- 35 (m, 2H), 6.96 (d, 1H, *J* = 8.7 Hz), 5.77-5.71 (m, 2H), 5.38 (dd, *J* = 1.2, 17.9 Hz), 4.32-4.26 (m, 2H), 1.33 (t, 2H, *J* = 7.1 Hz). FABLRMS *m/z* 333.2 (M+H). ESHRMS *m/z* 333.0510 (M+H, Calc'd. 333.050532. Anal. Calc'd for C<sub>15</sub>H<sub>12</sub>ClF<sub>3</sub>O<sub>3</sub> (1.14


wt % H<sub>2</sub>O): C, 53.53; H, 3.72; Cl, 10.53. Found: C, 53.46; H, 3.42; Cl, 10.70.

Step 6. Preparation of 6-chloro-4-ethenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 5 (0.300 g, 0.902 mmol) was dissolved in a THF-EtOH-H<sub>2</sub>O mixture (10 mL, 7:2:1) and treated with sodium hydroxide (0.360 mL, 0.902 mmol of a 2.5 N solution). This solution was stirred at room temperature for 18 hours. The solvent was evaporated and the residue was dissolved in water (10 mL). Diethyl ether (10 mL) was added and the mixture acidified by the addition of concentrated HCl. The organic layer was separated, and the aqueous phase was extracted with diethyl ether (2 x 10 mL). The ether extracts were combined, dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo yielding a yellow solid, which was recrystallized in diethyl ether-hexane to afford the title compound as a white solid (0.163 g, 59%): mp 143.0-145.0 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz 7.49 (d, 1H, J = 2.6 Hz), 7.33-7.17 (m, 2H), 6.99 (d, 1H, J = 8.5 Hz), 5.82-5.72 (m, 2H), 5.42 (d, 1H, J = 17.9 Hz). ESHRMS m/z 303.00207 (M-H, Calc'd. 303.003582). Anal. Calc'd for C<sub>17</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>3</sub> (1.10 wt % H<sub>2</sub>O): C, 50.69; H, 2.74; Cl, 11.51. Found: C, 50.57; H, 2.37; Cl, 11.75.

25

**EXAMPLE 143**




30 **6-Chloro-2-(trifluoromethyl)-4-phenyl-2H-1-benzopyran-3-carboxylic acid**




The 2H-1-benzopyran-3-carboxylic acid was prepared from ethyl 6-chloro-4-(trifluoromethanesulfonyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 142, Step 4) using a procedure similar to that described in Example 142, Steps 5-6: mp 225.5-226.6 °C <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>/300 MHz). 7.46-7.39 (m, 4H), 7.20-7.13 (m, 3H), 6.52 (d, 1H, *J* = 2.42 Hz), 6.12 (q, 1H, *J* = 7.1 Hz). FABLRMS *m/z* 355.1 (M+H). ESHRMS *m/z* 353.0215 (M-H, Calc'd. 353.019232). Anal. Calc'd. for C<sub>17</sub>H<sub>10</sub>ClF<sub>3</sub>O<sub>3</sub>: C, 57.56; H, 2.84; Cl, 10.17. Found: C, 57.18; H, 2.66; Cl, 10.17.

### EXAMPLE 144



#### 6-Chloro-4-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

The 2H-1-benzopyran-3-carboxylic acid was prepared from ethyl 6-chloro-4-(trifluoromethanesulfonyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate (Example 142, Step 4) using a procedure similar to that described in Example 142, Steps 5-6: mp 200.8-206.7 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.52 (dd, 1H, *J* = 1.21, 5.04 Hz), 7.28 (dd, 1H, *J* = 2.42, 8.67 Hz), 7.15 (dd, 1H, *J* = 1.21, 3.42 Hz), 6.98-6.93 (m, 2H), 5.83 (q, 1H, *J* = 6.9 Hz). FABLRMS *m/z* 378 (M+NH<sub>4</sub>). Anal. Calc'd. for C<sub>17</sub>H<sub>8</sub>ClF<sub>3</sub>O<sub>3</sub>S: C, 49.94; H, 2.24; Cl, 9.83; S, 8.89. Found: C, 50.02; H, 1.98; Cl, 9.34; S, 8.89.

**EXAMPLE 145**

5           **6-Methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-**  
              **carboxylic acid**

Step 1. Preparation of 5-methyl-2-mercaptobenzaldehyde.

10           Tetramethylethylenediamine (TMEDA) (12.6 mL, 83.5 mmol)  
              was added via syringe to n-BuLi (33 mL of 1.6 M in hexanes,  
              82.5 mmol) and the solution was chilled to 0 °C. A solution  
              of *p*-thiocresol (4.53 g, 36.5 mmol) in cyclohexane (40 mL)  
              was added with stirring over 5 minutes. The resulting tan  
              slurry was stirred overnight at room temperature, chilled to  
15           0 °C, and DMF (4.0 mL, 3.77 g, 51.6 mmol) was added via  
              syringe over 2 minutes. The resulting gummy slurry was  
              stirred at room temperature for 1.3 hours. The reaction  
              mixture was added to 3 N HCl (150 mL). This mixture was  
              extracted with ethyl acetate. The combined organic layers  
20           were washed with brine, dried over MgSO<sub>4</sub>, filtered and  
              concentrated *in vacuo* yielding a brown oil. This oil was  
              purified by flash chromatography over silica gel, eluting  
              with 10% ethyl acetate-hexanes to afford 5-methyl-2-  
              mercaptobenzaldehyde (4.47 g, 69%) as an intensely yellow  
25           solid suitable for use without further purification.

Step 2. Preparation of ethyl 6-methyl-2-(trifluoromethyl)-  
2H-1-benzothiopyran-3-carboxylate.

30           The 5-methyl-2-mercaptobenzaldehyde (Step 1) (3.25 g,  
              21.3 mmol) was added to DMF (5 mL) and ethyl 4,4,4-  
              trifluorocrotonate (4.32 g, 25.7 mmol). With stirring, K<sub>2</sub>CO<sub>3</sub>


(3.78 g, 27.3 mmol) was added causing the reaction to become a deep red. The reaction was stirred at room temperature for 20 hours, acidified with 3N HCl, diluted with ethyl acetate and washed with water, saturated NaHCO<sub>3</sub> solution, brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* yielding an oil. The oil was crystallized from diethyl ether-petroleum ether to give ethyl 6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylate as a light yellow solid (4.47 g, 69%): mp 93.1-94.7 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.41 (s, 1H), 7.31 (d, 1H, *J* = 7.9 Hz), 7.25 (d, 1H, *J* = 7.9 Hz), 4.96 (q, 1H, *J* = 8.5 Hz), 4.33 (m, 2H), 2.34 (s, 3H), 1.35 (t, 3H, *J* = 7.0 Hz). FABLRMS *m/z* 309 (M+Li).

Step 3. Preparation of 6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid.

The ester from Step 2 (0.55 g 1.8 mmol) was dissolved in THF (1.5 mL) and ethanol (1.5 mL), treated with 2.5 N sodium hydroxide (1.5 mL, 3.8 mmol), and stirred at room temperature for 88 hours. The reaction mixture was concentrated *in vacuo*, acidified with 3 N HCl, filtered, and recrystallized from diethyl ether/petroleum ether to yield the title compound as a yellow solid (0.14 g, 28%): mp 180.8-184.2 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.95 (s, 1H), 7.42 (s, 1H), 7.31 (d, 1H, *J* = 8.1 Hz), 7.25 (d, 1H, *J* = 8.1 Hz), 4.94 (q, 1H, *J* = 8.7 Hz), 2.34 (s, 3H). FABLRMS *m/z* 281 (M+Li). EIHRMS *m/z* 274.0250 (M+, Calc'd 274.0275). Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>F<sub>3</sub>O<sub>2</sub>S: C, 52.55; H, 3.31. Found: C, 52.54; H, 3.35.

30

**EXAMPLE 146**




**6,8-Dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

5

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 145: mp 220-225 °C (dec). <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 11.5 (brs, 1H), 7.94 (s, 1H), 7.26 (s, 1H) 7.14 (s, 1H), 4.98 (q, 1H, *J* = 8.7 Hz), 2.34 (s, 3H), 2.31 (s, 3H). FABLRMS *m/z* 295 (M+Li). EIHRMS *m/z* 288.0431 (M+, Calc'd 288.0432). Anal. Calc'd for C<sub>13</sub>H<sub>11</sub>F<sub>3</sub>O<sub>2</sub>S: C, 54.16; H, 3.85. Found: C, 54.10; H, 3.91.

15

**EXAMPLE 147**




20

**6-(1,1-Dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 145: mp 183.8-184.6 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.04 (s, 1H), 7.68 (d, 1H, *J* = 2.2 Hz), 7.46 (dd, 1H, *J* = 8.3 Hz 2.2 Hz), 7.37 (d, 1H, *J* = 8.3 Hz), 4.94 (q, 1H, *J* = 8.7 Hz), 1.34 (s, 9H). FABLRMS *m/z* 334 (M+NH<sub>4</sub>). ESHRMS *m/z* 334.1087 300 MHz) 7.52(dd, 1H, *J* = 1.21, 5.04 Hz), 7.28 (dd, 1H, *J* =

2.42, 8.67 Hz), 7.15 (dd, 1H,  $J = 1.21, 3.42$  Hz), 6.98-6.93 (m, 2H), 5.83 (q, 1H,  $J = 6.9$  Hz). FABLRMS  $m/z$  378 ( $M+NH_4$ ). Anal. Calc'd. for  $C_{15}H_8ClF_3O_2S$ : C, 49.94; H, 2.24; Cl, 9.83; S, 8.89. Found: C, 50.02; H, 1.98; Cl, 9.34; S, 8.89. ( $M+NH_4$ , Calc'd 334.1089). Anal. Calc'd for  $C_{15}H_{15}F_3O_2S$ : C, 56.95; H, 4.78. Found: C, 57.03; H, 4.83.


### EXAMPLE 148



#### 7-Methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid

15 The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 145: mp 186.6-191.9 °C.  $^1H$  NMR (acetone- $d_6$ /300 MHz) 7.96 (s, 1H), 7.49 (dd, 1H,  $J = 7.6$  Hz 2.82 Hz), 7.27 (s, 1H), 7.14 (d, 1H,  $J = 7.6$  Hz), 4.96 (q, 1H,  $J = 5.3$  Hz), 2.36 (s, 20 3H). ESHRMS  $m/z$  273.0204 (M-H, Calc'd 273.0197). Anal. Calc'd for  $C_{12}H_8F_3O_2S$  (3.32 wt %  $H_2O$ ): C, 50.81; H, 3.57. Found: C, 50.79; H, 3.44.


### EXAMPLE 149



#### 6,7-Dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 145: mp 235-237 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 7.90 (s, 1H), 7.33 (s, 1H), 7.19 (s, 1H), 4.91 (q, 1H, *J* = 8.7 Hz), 2.28 (s, 3H), 2.26 (s, 3H). FABLRMS *m/z* 295 (M+Li). EIHRS *m/z* 288.0439 (M+, Calc'd 288.0432). Anal. Calc'd for C<sub>13</sub>H<sub>11</sub>F<sub>3</sub>O<sub>2</sub>S: C, 54.16; H, 3.85. Found: C, 54.13; H, 3.85.

10


**EXAMPLE 150**

15

**8-Methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**


The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 145: mp 224-225 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 11.60 (br s, 1H), 8.00 (s, 1H), 7.44 (d, 1H, *J* = 6.7 Hz), 7.31 (d, 1H, *J* = 6.8 Hz), 7.21 (m, 1H), 5.05 (q, 1H, *J* = 8.5 Hz), 2.38 (s, 3H). FABLRMS *m/z* 292 (M+NH<sub>4</sub>). ESHRMS *m/z* 292.0591 (M+NH<sub>4</sub>, Calc'd 292.0619). Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>F<sub>3</sub>O<sub>2</sub>S: C, 52.55; H, 3.31. Found: C, 52.63; H, 3.38.

25

**EXAMPLE 151**

**2-(Trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

The 2H-1-benzothiopyran-3-carboxylic acid was prepared  
5 by a procedure similar to the method described in Example  
145: mp 187-190 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.01 (s,  
1H), 7.60 (d, 1H, *J* = 7.5 Hz), 7.45 (m, 2H), 7.31 (m, 1H),  
4.98 (q, 1H, *J* = 8.7 Hz). ESHRMS *m/z* 259.0070 (M-H, Calc'd  
259.0041). Anal. Calc'd for C<sub>11</sub>H<sub>7</sub>F<sub>3</sub>O<sub>2</sub>S: C, 50.77; H, 2.71.  
10 Found: C, 50.75; H, 2.78.

**EXAMPLE 152**

15

**6-Chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

Step 1. Preparation of N,N-dimethyl-O-(4-chloro-  
20 2-formyl-5-methylphenyl)thiocarbamate.

A mixture of 5-chloro-4-methylsalicylaldehyde (12.96 g,  
76.0 mmol) and triethylamine (11.58 g, 114.4 mmol) was  
dissolved in anhydrous DMF (15 mL) treated with *N,N*-  
dimethylthiocarbamoyl chloride (11.25 g, 91.0 mmol) and  
25 stirred at room temperature for 16 hours. The reaction was  
treated with 3 N HCl (50 mL) and filtered to give an orange  
solid. The solid was dissolved in ethyl acetate washed with  
3 N HCl, water, brine, dried over anhydrous MgSO<sub>4</sub>, filtered  
and concentrated in vacuo to afford a brown solid (16.79 g)  
30 which was recrystallized from diethyl ether/hexane to give  
the O-aryl thiocarbamate as a tan solid (4.92 g, 25%): <sup>1</sup>H

NMR (acetone- $d_6$ /300 MHz) 9.96 (s, 1H), 7.80 (s, 1H), 7.19 (s, 1H), 3.46 (s, 3H), 3.42 (s, 3H), 2.43 (s, 3H).

Step 2. Preparation of N,N-dimethyl-S-(4-chloro-2-formyl-5-methylphenyl)thiocarbamate.

5 The O-aryl thiocarbamate (Step 1) (4.92 g, 19.1 mmol) was dissolved in N,N-dimethylaniline (25 mL) and immersed in and stirred at 200 °C for 1.5 hours. The reaction mixture was cooled to room temperature and poured into a mixture of 10 3 N HCl (200 mL) and ice. Filtration gave a brown semisolid which was dissolved in ethyl acetate, washed with 3 N HCl, brine, dried over anhydrous  $MgSO_4$ , filtered and concentrated in vacuo to afford the S-arylthiocarbamate as a brown oil (3.80 g, 77%) which was used in the next step without 15 further purification.

Step 3. Preparation of ethyl 6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylate.


20 The S-arylthiocarbamate (Step 2) (3.80 g, 14.7 mmol) was dissolved in THF (10 mL) and ethanol (10 mL), treated with 2.5 N sodium hydroxide (16.5 mL, 34.2 mmol), and stirred at room temperature for 0.9 hours. The reaction was diluted with diethyl ether and washed with 3 N HCl, brine, dried over  $MgSO_4$ , filtered and concentrated in vacuo to 25 yield the crude substituted 2-mercaptobenzaldehyde as a brown oil (2.82 g). This oil was added to DMF (10 mL) and ethyl 4,4,4-trifluorocrotonate (3.89 g, 23.1 mmol). With stirring,  $K_2CO_3$  (3.23 g, 23.4 mmol) was added causing the reaction to become a deep red. The reaction was stirred at 30 room temperature for 14.5 hours, acidified with 3 N HCl, extracted with ethyl acetate. The resulting organic phase was washed with brine, dried over  $MgSO_4$ , filtered and concentrated in vacuo to give a yellow solid (6.36 g) which was used in the next step without further purification.



Step 4. Preparation of 6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid

The ester from Step 3 (2.02 g, 6.0 mmol) was dissolved in THF (10 mL) and ethanol (10 mL), treated with 2.5 N sodium hydroxide (5.5 mL, 13.8 mmol), and stirred at room temperature for 4.8 hours. The reaction mixture was concentrated in vacuo, acidified with 3 N HCl yielding a suspension. The solid was collected by filtration and was recrystallized from ethanol-water to yield the title compound as a yellow solid (0.20 g, 11%): mp 240.5-241.7 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300MHz) 7.99 (s, 1H), 7.67 (s, 1H), 7.43 (s, 1H), 4.99 (q, 1H, J = 8.5 Hz), 2.39 (s, 3H). FABLRMS m/z 307 (M-H). FABHRMS m/z 306.9831 (M-H, Calc'd 306.9807). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>ClF<sub>3</sub>O<sub>2</sub>S: C, 46.69; H, 2.61; Cl, 11.48. Found: C, 46.78; H, 2.61; Cl, 11.41.


**EXAMPLE 153**



**7-Chloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 152: mp 225.7-227.3 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.02 (s, 1H), 7.63 (d, 1H, J = 8.3 Hz), 7.54 (d, 1H, J = 2.0 Hz), 7.36 (dd, 1H, J = 8.3 Hz 2.0 Hz), 5.04 (q, 1H, J = 8.5 Hz). ESHRMS m/z 292.9646 (M-H, Calc'd 292.9651).


**EXAMPLE 154**



5                   **6,7-Dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid**

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 152: mp 262.5-263.5 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.04 (s, 1H), 7.90 (s, 1H), 7.74 (s, 1H), 5.09 (q, 1H, J = 8.5 Hz). ESHRMS m/z 326.9242 (M-H, Calc'd 326.9261).

**EXAMPLE 155**




15

**2-(Trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzothiopyran-3-carboxylic acid**


20                   The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 152: mp 129.3-132.4 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>/300 MHz) 8.10 (s, 2H), 8.00 (s, 2H), 7.71 (d, 2H, J = 8.1 Hz), 7.65 (d, 2H, J = 8.1 Hz), 5.09 (q, 1H, J = 8.5 Hz). ESHRMS m/z 358.9630 (M-H, Calc'd 358.9635).

25

**EXAMPLE 156**

5           **6,8-Dichloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid**

The 2H-1-benzothiopyran-3-carboxylic acid was prepared by a procedure similar to the method described in Example 152: mp 217.9-220.3 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 12.50-11.20 (br s, 1H exch.), 8.06 (s, 1H), 7.75 (d, 1H, *J* = 2.0 Hz), 7.64 (d, 1H, *J* = 2.2 Hz), 5.23 (q, 1H, *J* = 8.5 Hz). ESLRMS *m/z* 327 (M-H). ESHRMS *m/z* 326.9272 (M-H, Calc'd 326.9261).

**EXAMPLE 157**

20

**6-Chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid**

25   Step 1. Preparation of 2-amino-5-chlorobenzaldehyde.

2-Amino-5-chlorobenzyl alcohol (4.8 g, 30 mmol) and activated manganese (IV) oxide (21 g, 240 mmol) were refluxed in chloroform (100 mL) for 1 hour. The contents were allowed to cool, filtered through diatomaceous earth and concentrated *in vacuo* to afford the 2-amino-5-chlorobenzaldehyde as a dark solid (4.14 g, 81%): mp 74-76 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 9.80 (s, 1H),

7.42 (s, 1H), 7.23 (d, 1H,  $J = 7.0$  Hz), 6.60 (d, 1H,  $J = 7.0$  Hz).

5    Step 2. Preparation of ethyl 6-chloro-1,2-  
6    dihydro-2-(trifluoromethyl)-3-  
7    quinolinecarboxylate.


8    The 2-amino-5-chlorobenzaldehyde from Step 1  
9    (15.0 g, 96 mmol), anhydrous potassium carbonate  
10    (27.6 g, 200 mmol), and ethyl 4,4,4-  
11    trifluorocrotonate (34 mL, 200 mmol) were mixed in  
12    anhydrous dimethylformamide (60 mL) and heated at  
13    100 °C for 7 hours. The contents were allowed to  
14    cool and partitioned between ethyl acetate (200  
15    mL) and water (200 mL). The aqueous layer was  
16    extracted with ethyl acetate (1 x 100 mL). The  
17    ethyl acetate extracts were combined and washed  
18    with brine (1 x 200 mL), dried over  $MgSO_4$ , and  
19    concentrated in vacuo leaving a dark oil which  
20    solidified upon standing. The solid was purified  
21    by flash chromatography (silica gel; ethyl  
22    acetate-hexanes, 1:9). Fractions containing the  
23    desired product were combined, concentrated in  
24    vacuo and the residue recrystallized from ethyl  
25    acetate-hexanes to afford the ethyl 6-chloro-1,2-  
26    dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate  
27    as a yellow solid (16.36 g, 56%): mp 132.6-134.2  
28    °C.  $^1H$  NMR ( $CDCl_3$ , 300 MHz) 7.61 (s, 1H), 7.10 (m,  
29    2H), 6.55 (d, 1H,  $J = 8.0$  Hz), 5.10 (q, 1H,  $J =$   
30    6.0 Hz), 4.55 (brs, 1H), 4.23 (m, 2H), 1.32 (t,  
31    3H,  $J = 7.0$  Hz). FABHRMS  $m/z$  306.0468 ( $M+H^+$ ,  
32    Calc'd 306.0509). Anal. Calc'd for  $C_{11}H_{11}NO_2F_3Cl$ :  
33    C, 51.08; H, 3.63; N, 4.58. Found: C, 50.81; H,  
34    3.49; N, 4.72.

35    Step 3. Preparation of 6-chloro-1,2-dihydro-2-  
36    (trifluoro-methyl)-3-quinolinecarboxylic acid.

37    The ester from Step 2 (1.7 g, 5.6 mmol) and

2.5 N sodium hydroxide (4.4 mL, 11 mmol) were mixed in tetrahydrofuran (25 mL), methanol (10 mL), and water (25 mL). After stirring overnight, contents were concentrated *in vacuo* to remove the THF and methanol. The aqueous solution remaining was extracted with diethyl ether (2 x 100 mL). The resulting aqueous layer was acidified with 2 N HCl causing the precipitation of an oil. The oil was purified by flash chromatography on silica gel, eluting with ethyl acetate-hexanes (1:1). Fractions containing the desired product were combined, and concentrated *in vacuo*. The residue was triturated with dichloromethane, and filtered to afford the 6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid as a yellow solid (0.645 g, 41 %): mp 187.8-188.8 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>, 300 MHz) 7.69 (s, 1H), 7.36 (s, 1H), 7.15 (d, 1H, *J* = 8.0 Hz), 6.83 (d, 1H, *J* = 8.0 Hz), 6.60 (brs, 1H), 5.20 (m, 1H). ESHRMS *m/z* 276.0040 (M-H, Calc'd 276.0039). Anal. Calc'd for C<sub>11</sub>H<sub>7</sub>NO<sub>2</sub>F<sub>3</sub>Cl + 2.6% H<sub>2</sub>O: C, 46.39; H, 2.98; N, 4.92. Found: C, 45.99; H, 2.54; N, 4.85.

### EXAMPLE 158




### 6,8-Dichloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that

described in Example 157: mp 223.4-225.7 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>, 300 MHz) 7.82 (s, 1H), 7.40 (m, 2H), 6.53 (brs, 1H), 5.40 (m, 1H). ESHRMS m/z 309.9657 (M-H, Calc'd 309.9649). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>NO<sub>2</sub>F<sub>3</sub>Cl<sub>2</sub>: C, 42.34; H, 1.94; N, 4.49. Found: C, 42.20; H, 1.74; N, 4.52.


### EXAMPLE 159



#### 6,7-Difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that described in Example 157: mp 186.6-188.9 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>, 300 MHz) 7.79 (s, 1H), 7.32 (m, 1H), 6.71 (m, 1H), 6.64 (brs, 1H), 5.21 (m, 1H). ESHRMS m/z 278.0262 (M-H, Calc'd 278.0240). Anal. Calc'd for C<sub>11</sub>H<sub>6</sub>NO<sub>2</sub>F<sub>3</sub> + 1.58% H<sub>2</sub>O: C, 46.58; H, 2.31; N, 4.94. Found: C, 46.20; H, 2.07; N, 4.54.

### EXAMPLE 160




#### 6-Iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

Step 1. Preparation of ethyl 6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate.

A mixture of 5-iodo-2-aminobenzaldehyde (24.0 g, 96.7 mmol), diazobicyclo[2.2.2]-undec-7-ene (32.2 g, 212.0 mmol), and ethyl 4,4,4-trifluorocrotonate (35.7 g, 212.0 mmol) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (48 mL) was heated at 60 °C for 8 hours. The solution was cooled to room temperature and the solution poured into ethyl acetate-hexanes (1:1, 500 mL). The solution was extracted with 2.5 N aqueous hydrochloric acid (2 x 200 mL), saturated aqueous ammonium chloride (2 x 200 mL), dried over sodium sulfate, filtered and concentrated in vacuo. The resulting dark yellow oil was dissolved in hexanes (100 mL) and fine yellow crystals formed upon standing. Vacuum filtration of this suspension yielded ethyl 6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate as fine yellow crystals (19.3 g, 50 % yield): mp 137-138 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 7.62 (s, 1H), 7.36-7.48 (m, 2H), 6.43 (d, *J* = 8.2 Hz), 5.36 (brs, 1H), 5.11 (q, 1H, *J* = 7.1 Hz), 4.25 -4.35 (m, 2H), 1.34 (t, 3H, *J* = 7.0 Hz). ESHRMS *m/z* 395.9716 (M-H, Calc'd 395.9708).

Step 2. Preparation of 6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

Hydrolysis of the ester (Step 1) was performed by a procedure similar to that described in Example 157, Step 3, yielding the carboxylic acid. mp 188-192 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.668 (s, 1H), 7.46 (d, 1H, *J* = 2.2 Hz), 7.39 (dd, 1H, *J* = 8.4, 2.2 Hz), 6.52 (d, 1H, *J* = 8.4 Hz), 5.01 (q, 1H, *J* = 7.5 Hz). ESHRMS *m/z* 367.9401 (M, Calc'd 367.9395).

**EXAMPLE 161**


5

**6-Bromo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid**

The 1,2-dihydro-3-quinolinecarboxylic acid  
10 was prepared by a procedure similar to that  
described in Example 160: mp 185-186 °C. <sup>1</sup>H NMR  
(CD<sub>3</sub>OD/300 MHz) 7.68 (s, 1H), 7.31 (d, 1H, *J* = 2.2  
Hz), 7.23 (dd, 1H, *J* = 8.7, 2.2 Hz), 6.64 (d, 1H,  
*J* = 8.7 Hz), 5.01 (q, 1H, *J* = 7.5 Hz). EIHMS *m/z*  
15 319.9519 (M, Calc'd 319.9534). Anal. Calc'd for  
C<sub>11</sub>H<sub>6</sub>BrF<sub>3</sub>NO<sub>2</sub>: C, 41.02; H, 2.19; N, 4.35; Found: C,  
41.27, H, 2.23, N, 4.26.

**EXAMPLE 162**

20

**1,2-Dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylic acid**

25

**Step 1. Preparation of 2-amino-5-(trifluoromethoxy)benzoic acid.**

5-(Trifluoromethoxy)isatin (15.0 g, 65 mmol)  
and potassium hydroxide pellets (4 g) were mixed  
30 in water (35 mL) and cooled to 0 °C. With  
vigorous stirring, a solution of 30% aqueous



hydrogen peroxide (11.7 g), potassium hydroxide pellets (5.8 g), and water (80 mL) was added drop-wise keeping the temperature below 10 °C. After stirring 1 hour at 0 °C, glacial acetic acid (22 mL) was added drop-wise, causing foaming and formation of a precipitate. The contents were stirred overnight and filtered to afford the 2-amino-5-trifluoromethoxybenzoic acid as an amber solid (12.5 g, 87 %). A small amount was recrystallized from ethyl acetate-hexanes to afford amber needles for an analytical sample and the remaining compound was used without further purification: mp 142.5-144.2 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 7.98 (s, 1H), 7.18 (d, 1H, *J* = 8.0 Hz) 6.62 (d, 1H, *J* = 8.0 Hz), 6.40 (brs, 2H). Anal. Calc'd for C<sub>8</sub>H<sub>6</sub>NO<sub>3</sub>F<sub>3</sub>: C, 43.45; H, 2.73; N, 6.33. Found: C, 43.40; H, 2.65; N, 6.35.

Step 2. Preparation of 2-amino-5-(trifluoromethoxy)benzyl alcohol.

The 2-amino-5-trifluoromethoxybenzoic acid (2.0 g, 9.0 mmol) in tetrahydrofuran (20 mL) was added dropwise to borane methyl sulfide complex (1.5 mL, 15.0 mmol) in tetrahydrofuran (5 mL). The reaction was refluxed overnight and allowed to cool. A solution of 30 % aqueous hydrogen peroxide (0.5 mL), 2.5 N sodium hydroxide (0.5 mL) and water (10 mL) was added drop-wise and the reaction stirred 0.5 hours. After diluting with diethyl ether (50 mL), the organic layer was washed with 0.1 M aqueous sodium meta-bisulfite (2 x 10 mL) and 2.5 N aqueous sodium hydroxide (2 x 10 mL). The organic layer was diluted further with hexanes (50 mL) and washed with brine (2 x 20 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo leaving an amber oil (1.9 g) which solidified. The solid was recrystallized from

ethyl acetate-hexanes to afford the 2-amino-5-trifluoromethoxybenzyl alcohol as a light amber solid (1.44 g, 77%): mp 75.9-77.6 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 7.00 (m, 2H), 6.65 (d, 1H, J = 8.0 Hz), 4.05 (s, 2H), 3.25 (brs, 3H). ESHRMS m/z 208.0592 (M+H<sup>+</sup>, Calc'd 208.0585). Anal. Calc'd for C<sub>8</sub>H<sub>8</sub>NO<sub>2</sub>F<sub>3</sub>: C, 46.39; H, 3.89; N, 6.76. Found: C, 46.61; H, 3.79; N, 6.71.

10 Step 3. Preparation of 2-amino-5-(trifluoromethoxy)-benzaldehyde.

The 2-amino-5-trifluoromethoxybenzyl alcohol from Step 2 (9.7 g, 47 mmol) and manganese (IV) oxide (21 g, 240 mmol) were refluxed in chloroform 15 (200 mL) for 1 hour. The contents were allowed to cool and filtered. The filtrate was concentrated in vacuo leaving an amber oil (8.2 g) which solidified. The oil was distilled (bulb to bulb apparatus) at 50 °C (0.1 mm) to afford a yellow 20 solid (7.2 g). The solid was recrystallized from hexanes to afford the desired 2-amino-5-(trifluoromethoxy)-benzaldehyde as yellow crystals (4.4 g, 46%): mp. 42-44 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 9.81 (s, 1H), 7.36 (s, 1H), 7.20 (d, 1H, J = 25 9.0 Hz), 6.64 (d, 1H, J = 9.0 Hz). EIHRMS m/z 205.0328 (M<sup>+</sup>, Calc'd 205.0350).

30 Step 4. Preparation of ethyl 1,2-dihydro-6-(trifluoro-methoxy)-2(trifluoromethyl)-3-quinolinecarboxylate.

The 2-amino-5-(trifluoromethoxy)benzaldehyde from Step 3 (5.3 g, 26 mmol), anhydrous potassium carbonate (6.9 g, 50 mmol), and ethyl 4,4,4-trifluorocrotonate (7.7 mL, 50 mmol) were mixed in 35 anhydrous dimethylformamide (50 mL) and heated at 90 °C for 6 hours. The reaction was allowed to


cool to room temperature and was partitioned between ethyl acetate (200 mL) and water (200 mL). The aqueous layer was extracted with more ethyl acetate (100 mL). The ethyl acetate extracts were  
5 combined and washed with brine (200 mL), dried over  $\text{MgSO}_4$ , and concentrated in vacuo yielding an oil (9.6 g). The oil was purified by flash chromatography on silica gel, eluting with ethyl acetate-hexanes (1:1). Fractions containing the  
10 desired product were combined, concentrated in vacuo, and the residue recrystallized from ethyl acetate-hexanes to afford the ethyl 1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylate as a yellow solid (4.05 g,  
15 32%): mp. 123-125 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz) 7.65 (s, 1H), 7.02 (m, 2H), 6.60 (m, 1H), 5.10 (m, 1H), 4.60 (brs, 1H), 4.28 (m, 2H), 1.32 (t, 3H,  $J = 7.0$  Hz). ESHRMS  $m/z$  356.0698 (M-H, Calc'd 356.0721). Anal. Calc'd for  $\text{C}_{14}\text{H}_{11}\text{NO}_3\text{F}_6$ : C, 47.34; H, 3.12; N, 3.94. Found: C, 47.37; H, 3.04; N, 3.93.

Step 5. Preparation of 1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylic acid.

25 The ethyl 1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylate from Step 4 (880 mg, 2.5 mmol) and 2.5 N aqueous sodium hydroxide (2 mL) were mixed in methanol (15 mL) and water (15 mL). The solution was heated on a  
30 steam bath for 2 hours. The reaction was allowed to cool to room temperature and was extracted with diethyl ether (50 mL). The aqueous layer was acidified (pH = 1) with 3 N HCl and extracted with ethyl acetate (2 x 50 mL). The combined ethyl  
35 acetate extracts were dried over  $\text{MgSO}_4$ , and concentrated in vacuo leaving an oil. The oil was crystallized from cold dichloromethane-hexanes to

afford the 1,2-dihydro-6-(trifluoromethoxy)-  
2(trifluoromethyl)-3-quinolinecarboxylic acid as  
yellow needles (0.727 g, 89%): mp 127.7-128.9 °C.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) 7.80 (s, 1H), 7.05 (m, 2H),  
5 6.62 (d, 1H, *J* = 8.0 Hz), 5.13 (m, 1H), 4.62 (brs,  
1H). ESHRMS *m/z* 326.0252 (M-H, Calc'd 326.0252).  
Anal. Calc'd for C<sub>12</sub>H<sub>7</sub>NO<sub>3</sub>F<sub>6</sub>: C, 44.05; H, 2.16; N,  
4.28. Found: C, 43.89; H, 2.04; N, 4.24.

10

**EXAMPLE 163**

15

**6-(Trifluoromethyl)-1,2-dihydro-2-  
(trifluoromethyl)-3-quinolinecarboxylic acid****Step 1. Preparation of N-(4-  
trifluoromethylphenyl)-2,2-dimethylpropanamide.**

A solution of dichloromethane (200 mL), 4-  
20 aminobenzotrifluoride (32.0 g, 199 mmol) and  
triethylamine (40 g, 396 mmol) was cooled to 0 °C  
under a dry nitrogen atmosphere. Trimethylacetyl  
chloride (32.9 g, 273 mmol) was added drop-wise  
over 2 hours, maintaining the temperature below 10  
25 °C. After the addition, the contents were allowed  
to warm to room temperature for 2 hours. The  
reaction was washed with water (2 X 200 mL),  
saturated ammonium chloride solution (2 X 200 mL),  
dried over sodium sulfate and filtered. The  
30 solvent was removed in vacuo to afford a white  
solid, N-(4-trifluoromethylphenyl)-2,2-  
dimethylpropanamide (48.0 g, 98%): mp 157-159 °C.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.61 (ab, 4H, *J* = 8.7, Δ*v* =

28.6 Hz), 7.47 (br s, 1H), 1.33 (s, 9H). ESHRMS  
m/z 246.1123 (M+H<sup>+</sup>, Calc'd 246.1106). Anal.  
Calc'd for C<sub>11</sub>H<sub>14</sub>F<sub>3</sub>NO: C, 58.77; H, 5.75; N, 5.71.  
Found: C, 58.28; H, 5.79; N, 5.65.

5

Step 2. Preparation of N-[2-formyl-4-  
(trifluoromethyl) phenyl]-2,2-dimethyl  
propanamide.

A 1 liter three neck round bottom flask  
10 equipped with equalizing addition funnel, magnetic  
stirrer and temperature monitoring device was  
charged with N-(4-trifluoromethylphenyl)-2,2-  
dimethyl propanamide (10.13 g, 41.4 mmol) and  
anhydrous tetrahydrofuran (150 mL). The reaction  
15 was chilled to -78 °C under nitrogen followed by  
slow addition of n-butyllithium (50 ml, 2.5 M in  
hexanes, 124 mmol) over 0.5 hours, such that the  
temperature of the reaction did not rise above -  
65°C. The contents were held at -78 °C for one  
20 hour, 0 °C for two hours, then chilled back to -78  
°C. Excess N,N-dimethylformamide (100 mL, 1.37  
mol) was added. The contents were warmed to room  
temperature and stirred for two hours. Aqueous 1 N  
HCl was added to the reaction until the pH reached  
25 1. The reaction was washed with water (2 X 200  
mL), saturated ammonium chloride solution (2 X 200  
mL), dried over sodium sulfate and filtered. The  
filtrate was concentrated in vacuo to afford a  
yellow solid. The product was purified by flash  
30 chromatography (silica gel, 10% ethyl acetate, 90%  
hexanes) to yield, upon concentration of the  
appropriate fractions, N-(2-formyl-4-  
trifluoromethylphenyl)-2,2-dimethylpropanamide as  
a solid (7.36 g, 65%): mp 69-73 °C. <sup>1</sup>H NMR  
35 (CDCl<sub>3</sub>/300 MHz) 11.5 (br s, 1H), 9.99 (s, 1H),  
8.67 (d, 1H, J = 8.8 Hz), 7.94 (d, 1H, J = 1.6

Hz), 7.83 (m, 1H), 1.37 (s, 9H). ESHRMS  $m/z$  274.1060 ( $M + H^+$ , Calc'd 274.1055). Anal. Calc'd for  $C_{13}H_{14}F_3NO_2$ : C, 57.14; H, 5.16; N, 5.13. Found: C, 57.15; H, 5.43; N, 5.01.

5


Step 3. Preparation of ethyl 6-(trifluoromethyl)-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate.

- To a suspension of N-(2-formyl-4-(trifluoromethylphenyl)-2,2-dimethyl propanamide (Step 2) (921 mg, 3.7 mol) and lithium hydride (115 mg, 14.5 mmol) in dimethyl sulfoxide (10 mL) was added ethyl 4,4,4-trifluorocrotonate (2.83 g, 16.8 mmol) and the contents warmed to 30 °C for 4 hours.
- After the addition of ethyl acetate (50 mL), the reaction was washed with water (2 X 30 mL), saturated ammonium chloride solution (2 X 30 mL), dried over sodium sulfate and filtered. The filtrate was concentrated in vacuo to afford a yellow solid. The product was purified by flash chromatography (silica gel, eluant: ethyl acetate-hexanes, 1:9) to yield, upon concentration of the appropriate fractions, ethyl 6-trifluoromethyl-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate as a yellow solid (65 mg, 5%): mp 138-139 °C.  $^1H$  NMR ( $CDCl_3$ /300 MHz) 7.67 (s, 1H), 7.26 (s, 1H), 7.04 (d, 1H,  $J = 6.6$  Hz), 6.62 (m, 1H), 5.14 (m, 1H), 4.60 (brs, 1H), 4.32 (m, 2H), 1.35 (t, 3H,  $J = 7.0$  Hz). ESHRMS  $m/z$  338.0592 ( $M-H$ , Calc'd 338.0616). Anal. Calc'd for  $C_{13}H_{11}F_3NO_2$ : C, 49.57; H, 3.27; N, 4.13; Found: C, 49.23; H, 2.81; N, 3.93.

- Step 4. Preparation of ethyl 6-trifluoromethyl-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid.

Ethyl 6-trifluoromethyl-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate from Step 3 (45 mg, 0.13 mmol) was suspended in methanol-tetrahydrofuran- water (10 mL, 7:2:1). Lithium hydroxide (24 mg, 0.52 mmol) was added, and the mixture was gently heated to reflux for two hours. The reaction was cooled to room temperature and 1 N HCl added until pH = 1. The organic solvent was removed in vacuo to afford a suspension of a crude yellow solid. Diethyl ether (20 mL) was added, and the solution was washed with water (2 X 20 mL), saturated ammonium sulfate (2 X 20 mL), dried over sodium sulfate and filtered. The filtrate was concentrated in vacuo to yield 6-trifluoromethyl-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid as a yellow solid, (0.041 g, 0.132 mmol, 99%): mp 150-156 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.78 (s, 1H), 7.48 (s, 1H), 7.40 (m, 1H), 6.81 (m, 1H), 5.17 (m, 1H). ESHRMS m/z 310.0307 (M-H, Calc'd 310.0303).

#### EXAMPLE 164



#### 6-Cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

Step 1. Preparation of ethyl 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate.

N,N-Dimethylformamide (5 mL) was degassed with nitrogen for thirty minutes in a three neck round bottom flask equipped with a condenser,

temperature monitoring, nitrogen purge and heating mantle. Ethyl 6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate (Example 158) (0.522 g, 1.32 mmol) and zinc cyanide (0.102 g, 0.792 mmol) were added to the *N,N*-dimethylformamide and stirred vigorously for ten minutes. Tetrakis(triphenylphosphine)palladium(0) (0.068 g, 0.53 mmol) was added and the contents gently warmed to 80 °C for 2 hours under a nitrogen atmosphere. Ethyl acetate (20 mL) was added, followed by extraction with aqueous 2 N ammonium hydroxide (2 X 10 mL), water (2 X 10 mL), saturated ammonium chloride (2 X 10 mL), dried over sodium sulfate and solvent removed *in vacuo* to yield a yellow solid. The product was purified by flash chromatography (silica gel, ethyl acetate-hexanes, 3:1) to yield, upon concentration of the appropriate fractions, ethyl 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate as a yellow solid (188 mg, 48%): mp 211-212 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.68 (s, 1H), 7.43 (m, 2H), 6.69 (d, 1H, *J* = 8.3 Hz), 5.22 (m, 1H), 4.98 (br s, 1H), 1.30 (m, 2H), 1.36 (t, 3H, *J* = 7.1 Hz). EIHRMS *m/z* 314.1147 (*M*+NH<sub>4</sub><sup>+</sup>, Calc'd 314.1116). Anal. Calc'd for C<sub>14</sub>H<sub>11</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: C, 56.76; H, 3.74; N, 9.46. Found: C, 56.44; H, 4.03; N, 9.29.


Step 2. Preparation of 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid.

To a suspension of ethyl 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate (140 mg, 0.45 mmol) in methanol-tetrahydrofuran-water (10 mL, 7:2:1) was added lithium hydroxide (76 mg, 0.91 mmol) and the mixture gently heated to reflux for two hours. The contents were cooled to room



temperature and 1 N aqueous hydrochloric acid added until pH = 1. The organic solvent was removed in vacuo to afford a suspension of crude yellow solid. Diethyl ether (20 mL) was added, and the solution was washed with water (2 X 20 mL), saturated ammonium sulfate (2 X 20 mL), dried over sodium sulfate and filtered. The filtrate was concentrated in vacuo to yield 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid as a yellow solid, (116 mg, 95%): mp 238-240 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.75 (s, 1H), 7.56 (m, 1H), 7.43 (m, 1H), 6.79 (d, 1H, J = 8.5 Hz) 5.19 (q, 1H, J = 7.1 Hz). EIHRMS m/z 267.0405 (M-H, Calc'd 267.0381). Anal. Calc'd for C<sub>14</sub>H<sub>11</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: C, 53.74; H, 2.63; N, 10.45. Found: C, 53.99; H, 2.89; N, 10.19.

### EXAMPLE 165



#### 6-Chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid

Step 1. Preparation of ethyl 6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylate.

Ethyl 6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylate (Example 157, Step 2) (1.28 g, 4.21 mmol), tetrabutylammonium iodide (0.36 g, 0.92 mmol) and aqueous NaOH (50%, 2 mL) were stirred vigorously in methylene chloride (40 mL). Dimethyl sulfate


(2.12 g, 16.84 mmol) was added to the dark orange mixture via syringe over 2 hours. Hexane (5 mL) was added, and the solution was washed with water (2 X 20 mL), saturated ammonium chloride solution (2 X 20 mL), dried over sodium sulfate and filtered. The filtrate was concentrated in vacuo to afford the crude ester as a yellow solid. The solid was purified by flash chromatography (silica gel, 50 g; ethyl acetate-hexanes, 1:19) to yield, upon concentration of the appropriate fractions, ethyl 6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinoline-carboxylate (1.2 g, 90% yield): mp 118-120 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.71 (s, 1H), 7.30-7.26 (m, 2H), 6.77-6.74 (m, 1H), 5.12 (q, 1H, J = 6.8 Hz), 4.44-4.22 (m, 2H), 3.18 (s, 3H), 1.35 (t, 3H, J = 7.0 Hz). EIHMS m/z 320.0701 (M-H, Calc'd 320.0665) Anal. Calc'd for C<sub>14</sub>H<sub>13</sub>F<sub>3</sub>NO<sub>2</sub>Cl: C, 52.60; H, 4.10; N, 4.38. Found: C, 52.57; H, 4.14; N, 4.32.

Step 2. Preparation of 6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid

Ethyl 6-chloro-1,2-dihydro-1-methyl-2-(trifluoro-methyl)-3-quinolinecarboxylate (1.21 g, 3.78 mmol) was suspended in methanol-tetrahydrofuran-water (20 mL, 7:2:1). Lithium hydroxide (0.262 g, 6.24 mmol) was added, and the mixture was gently heated to reflux for two hours. The reaction was cooled to room temperature and 1 N HCl added until pH = 1. The organic solvent was removed in vacuo to afford a suspension of crude yellow solid. Diethyl ether (20 mL) was added, and the resulting solution was washed with water (2 X 20 mL), saturated ammonium chloride (2 X 20 mL), dried over sodium sulfate and filtered. The filtrate was concentrated in vacuo to afford the


product as a yellow solid, 6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinoline-carboxylic acid. (1.08 g, 98% yield): mp 208-209 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.69 (d, 1H, *J* = 2.5 Hz), 7.28-7.24 (m, 2H), 6.73 (dd, 1H, *J* = 9.5, 2.5 Hz), 5.13 (q, 1H, *J* = 7.0), 3.16 (s, 3H). Anal. Calc'd for C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>NO<sub>2</sub>Cl: C, 49.42; H, 3.11; N, 4.80; Cl, 12.16. Found: C, 49.88; H, 3.29; N, 4.59; Cl, 12.42

10

**EXAMPLE 166**

15     **6-Chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(trifluoromethyl)phenyl]methyl]-3-quinolinecarboxylic acid**


20     The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that described in Example 165: mp 229-231 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.77 (s, 1H), 7.58 (d, 2H, *J* = 8.0 Hz), 7.39 (d, 2H, *J* = 8.0 Hz), 7.30 (d, 1H, *J* = 2.4), 7.13 (dd, 1H, *J* = 8.9, 2.4 Hz), 6.75 (d, 1H, *J* = 8.9 Hz), 5.27 (q, 1H, *J* = 7.0 Hz), 4.90 (ab, 2H, *J* = 16.7 Hz, Δ*v* = 95.2 Hz). EIHRMS *m/z* 434.0401 (Calc'd for M-H 434.0383) Anal. Calc'd for C<sub>19</sub>H<sub>14</sub>F<sub>6</sub>NO<sub>2</sub>Cl: C, 52.13; H, 3.22; N, 3.22; Found: C, 52.36; H, 2.91; N, 3.21.

**EXAMPLE 167**

5

**6-Chloro-1-[(4-chlorophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid**

The 1,2-dihydro-3-quinolinecarboxylic acid  
10 was prepared by a procedure similar to that  
described in Example 165: mp 250-253 °C. <sup>1</sup>H NMR  
(CD<sub>3</sub>OD/300 MHz) 7.74 (s, 1H), 7.32-7.13 (m, 6H),  
6.76 (d, 1H, *J* = 8.7 Hz), 5.22 (q, 1H, *J* = 7.0  
Hz), 4.81 (ab, 2H, *J* = 16.3 Hz, Δ*v* = 54.7 Hz).  
15 ESHRMS *m/z* 400.0105 (M-H, Calc'd 400.0119).


**EXAMPLE 168**

20

**6-Chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(methoxy)phenyl]methyl]-3-quinolinecarboxylic acid**

The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that described in Example 165: mp 196-197 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.71 (s, 1H), 7.27-7.26 (m, 1H), 7.18-7.12 (m, 3H), 6.85-6.81 (m, 3H), 5.16 (q, 1H, J = 7.1 Hz), 4.69 (ab, 2H, J = 15.3 Hz, Δv = 111.8 Hz), 3.73 (s, 3H). ESHRMS m/z 396.0625 (M-H, Calc'd 396.0614). Anal. Calc'd for C<sub>19</sub>H<sub>14</sub>F<sub>3</sub>NO<sub>2</sub>Cl: C, 52.13; H, 3.22; N, 3.22. Found: C, 52.36; H, 2.91; N, 3.21.

### EXAMPLE 169




#### 6-Chloro-1-[(4-cyanophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid

20

The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that described in Example 165: mp 258-260 °C. <sup>1</sup>H NMR (CD<sub>3</sub>OD/300 MHz) 7.78 (s, 1H), 7.66 (d, 2H, J = 8.2 Hz), 7.41 (d, 2H, J = 8.2 Hz), 7.33 (d, 1H, J = 2.7 Hz), 7.15 (dd, 1H, J = 8.7, 2.7 Hz), 6.71 (d, 1H, J = 8.7 Hz), 5.31 (q, 1H, J = 7.0 Hz), 4.94 (ab, 2H, J = 17.1, Δv = 91.8 Hz). ESHRMS m/z

391.0443 (M-H, Calc'd 391.0461). Anal. Calc'd for  $C_{19}H_{12}F_3N_2O_2Cl + 0.53 \% H_2O$ : C, 57.79; H, 3.55; N, 7.09; Found: C, 57.26; H, 3.17; N, 6.78.


5

**EXAMPLE 170**

10 **6-Chloro-1,2-dihydro-1-[(4-nitrophenyl)methyl]-2-(trifluoromethyl)-3-quinolinecarboxylic acid**

The 1,2-dihydro-3-quinolinecarboxylic acid was prepared by a procedure similar to that described in Example 165: mp 225-228 °C.  $^1H$  NMR (CD<sub>3</sub>OD-3% TFA/300 MHz) 8.14 (d, 2H,  $J = 8.8$  Hz), 7.77 (s, 1H), 7.42 (d, 2H,  $J = 8.8$  Hz), 7.29 (d, 1H,  $J = 2.4$  Hz), 7.11 (dd, 1H,  $J = 8.9, 2.4$  Hz), 6.67 (d, 1H,  $J = 8.9$  Hz), 5.27 (q, 1H,  $J = 6.8$  Hz), 4.93 (ab, 2H,  $J = 17.2$  Hz,  $\Delta v = 95.0$  Hz).

20 ESHRMS  $m/z$  411.0327 (M-H, Calc'd 411.0359).


**EXAMPLE 171**

**6-Chloro-1,2-dihydro-1-ethyl-2-(trifluoromethyl)-  
3-quinolinecarboxylic acid**

5        The 1,2-dihydro-3-quinolinecarboxylic acid  
was prepared by a procedure similar to that  
described in Example 165: mp 201-202 °C. <sup>1</sup>H NMR  
(CD<sub>3</sub>OD/300 MHz) 7.67 (s, 1H), 7.25-7.22 (m, 2H),  
6.86 (d, 1H, *J* = 8.7 Hz), 5.21 (q, 1H, *J* = 7.0  
10 Hz), 3.81-3.71 (m, 1H), 3.47-3.39 (m, 1H), 1.20  
(t, 3H, *J* = 7.2 Hz). ESHRMS *m/z* 304.0360 (M-H,  
Calc'd 304.0352).

**EXAMPLE 172**

15




**(S)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-  
quinolinecarboxylic acid**

20

To a solution of 6-chloro-1,2-dihydro-2-  
(trifluoromethyl)-3-quinolinecarboxylic acid  
(Example 157) (6.75 g, 24.3 mmol) in ethyl acetate  
(25 mL) was added (S)-(-)- $\alpha$ -methylbenzylamine  
25 (1.50 g, 12.2 mmol). To the resulting solution  
was added hexanes (50 mL) with mixing. Stirring  
was discontinued and the reaction held static at  
room temperature for 16 hours during which time  
yellow crystals formed. The crystals were  
30 collected and washed with ethyl acetate-hexanes  
(100 mL, 1:2). The resulting yellow solid (932  
mg) was dissolved in ethyl acetate (20 mL) and  
extracted with 1 N HCl (3 x 10 mL). The organic  
layer was dried over sodium sulfate and solvent

removed at reduced pressure. The (s)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid was obtained as a yellow solid (648 mg, 10% yield). mp 173-176 °C. <sup>1</sup>H NMR (acetone-d<sub>6</sub>, 300 MHz) 7.80 (s, 1H), 7.35 (d, 1H, J = 2.2 Hz), 7.18 (d, 1H, J = 8.0, J = 2.2 Hz), 6.86 (d, 1H, J = 8.0 Hz), 6.60 (brs, 1H), 5.20 (m, 1H). Anal. Calc'd. for C<sub>11</sub>H<sub>7</sub>NO<sub>2</sub>F<sub>3</sub>Cl C, 47.40 H, 2.54; N, 5.40. Found C, 47.49; H, 2.60; N, 4.98. The compound was determined to have an optical purity greater than 90% ee. Optical purity was determined by HPLC as described in Example 66.

### EXAMPLE 173



#### 6-(2,2,2-Trifluoro-1-hydroxyethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid

Step 1. Preparation of ethyl 6-(1-hydroxy-2,2,2-trifluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylate.

The aldehyde (Example 75, Step 1) (0.89 g, 3.0 mmol) was cooled to 0 °C and treated with a 0.5 M solution of trimethyl(trifluoromethyl)silane (8.4 mL, 4.2 mmol) and four drops of a 1.0M solution of tetrabutylammonium fluoride was added. The reaction was allowed to warm to room temperature and stirred for 21.1 hours. The reaction was quenched with 3 N HCl, extracted with ethyl acetate, washed with water, brine, dried over



MgSO<sub>4</sub>, and concentrated *in vacuo* to give a brown oil (1.02 g). This oil was purified by flash chromatography over silica gel, eluting with 10% ethyl acetate/hexanes to afford a brown oil (0.77 g, 58%): <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 7.72 (d, 1H, *J* = 3.4 Hz), 7.34 (m, 2H), 6.99 (d, 1H, *J* = 8.5 Hz), 5.71 (q, 1H, *J* = 6.8 Hz), 4.83 (q, 1H, *J* = 6.4 Hz), 4.33 (m, 2H), 1.35 (t, 3H, *J* = 7.1 Hz), 0.11 (s, 9H). FABLRMS *m/z* 443 (M+H).


10

Step 2. Preparation of 6-(1-hydroxy-2,2,2-trifluoroethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid.

The ester from Step 1 (0.15g 0.34 mmol) was dissolved in THF (2 mL) and ethanol (2 mL), treated with 2.5 N NaOH (1 mL, 2.5 mmol), and stirred at room temperature for 18.6 hours. The reaction mixture was concentrated *in vacuo*, acidified with 3 N HCl, extracted with ethyl acetate, washed with 3 N HCl, brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give a yellow oil which was recrystallized from ethyl acetate/hexane to yield a white solid (0.03 g, 25%): mp 114-120 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.65 (s, 1H), 7.60 (dd, 1H, *J* = 8.2 Hz 2.0 Hz), 7.11 (d, 1H, *J* = 8.3 Hz), 5.87 (q, 1H, *J* = 7.0 Hz), 5.24 (q, 1H, *J* = 7.0 Hz). FABLRMS *m/z* 341 (M-H). ESHRMS *m/z* 341.0241 (M-H, Calc'd 341.0249).

30

**EXAMPLE 174**



35

**6-Chloro-2-(trifluoromethyl)-1,2-**

**dihydro[1,8]naphthyridine-3-carboxylic acid**Step 1. Preparation of N-[5-chloropyridin-2-yl]-2,2-dimethylpropanamide.

5 To 2-amino-5-chloropyridine (10.0 g, 0.078 mol) (Aldrich) and triethylamine (12 mL, 0.086 mol) in methylene chloride (200 mL), at 0 °C, was added dropwise trimethylacetyl chloride in methylene chloride (15 mL). The reaction was allowed to  
10 warm to room temperature while stirring overnight. The resulting mixture was washed with water, brine, and was dried over MgSO<sub>4</sub> and filtered. Concentration of the filtrate *in vacuo* provided a colorless oil (19.2 g). The oil was dissolved in  
15 hexanes and cooled causing the precipitation of a solid. The solid was collected by filtration affording the amide as a white solid (14.96 g, 90%): mp 51.4 - 53.4 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 8.25-8.15(m, 2H), 8.00 (br s, 1H), 7.68-7.60 (m, 1H), 1.28 (s, 9H). Anal: Calc'd for C<sub>10</sub>H<sub>13</sub>N<sub>2</sub>OCl: C, 56.47; H, 6.16; N, 13.17 Found: C, 56.72; H, 6.34; N, 12.88.

Step 2. Preparation of N-[5-chloro-3-formylpyridin-2-yl]-2,2-dimethylpropanamide.

25 To a chilled (-78 °C), stirred solution of the amide (Step 1) (5.0 g, 0.024 mole) in tetrahydrofuran (100 mL) was added t-butyl lithium (1.7M in pentane, 32.4 mL, 0.055 mole) dropwise.  
30 Dimethylformamide (2.3 mL, 0.03 mole) was added dropwise at -78 °C over 3 hours and the mixture allowed to warm room temperature. The reaction was quenched with ice water (200 mL) and extracted with ethyl acetate. The resulting organic phase  
35 was dried over MgSO<sub>4</sub> and was concentrated *in vacuo* to a volume of 20 mL. A white solid precipitated

which was collected by filtration yielding the formylated product (3.24 g, 56%): mp 168.7-170.8 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 10.60 (br s, 1H), 9.88 (s, 1H), 8.57 (s, 1H), 8.00 (s, 1H), 1.28 (s, 9H).  
5 Anal. Calc'd for C<sub>11</sub>H<sub>13</sub>N<sub>2</sub>OCl: C, 54.89; H, 5.44; N, 11.64 Found: C, 54.87; H, 5.42; N, 11.40.

Step 3. Preparation 2-amino-5-chloro-3-formylpyridine.

10 The product of Step 2 (2.7 g, 11 mmol) and 3 N HCl (50 mL) were heated at reflux for 2 hours. The reaction was allowed to cool to room temperature and was concentrated in vacuo yielding a light yellow solid (2.1 g). The solid was  
15 partitioned between ethyl acetate and 2.5 N NaOH solution. The ethyl acetate layer was dried over MgSO<sub>4</sub> and concentrated in vacuo providing a solid (1.7 g). The solid was recrystallized from ethyl acetate to give the desired substituted pyridine  
20 as yellow needles (1.2 g, 68%): mp 176.1-177.3 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>/300 MHz) 9.80 (s, 1H), 8.21 (s, 1H), 7.75 (s, 1H), 6.75 (br s, 2H). Anal. Calc'd for C<sub>6</sub>H<sub>5</sub>N<sub>2</sub>OCl: C, 46.03; H, 3.22; N, 17.89 Found: C, 45.90; H, 3.24; N, 17.80.

25


Step 4. Preparation of ethyl 6-chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylate.

The substituted pyridine from Step 3 (1.7 g, 11 mmol), anhydrous potassium carbonate (3.0 g, 22 mmol), and ethyl 4,4,4-trifluorocrotonate (3.3 mL, 22 mmol) were mixed in anhydrous dimethylformamide (20 mL) and heated at 80 °C for 2 hours. The  
30 reaction was allowed to cool to room temperature and was partitioned between ethyl acetate (100 mL) and water (100 mL). The aqueous layer was

extracted with more ethyl acetate (100 mL). The combined organic extracts were washed with brine (100 mL), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* yielding a waxy amber solid. The solid was  
5 triturated with diethyl ether providing the ester as a yellow solid (613 mg, 18%). A small amount was recrystallized from ethyl acetate for analytical data: mp 180.1-181.9 °C.  $^1\text{H}$  NMR (CDCl<sub>3</sub>/300 MHz) 7.99 (s, 1H), 7.61 (s, 1H), 7.39  
10 (s, 1H), 6.00 (br s, 1H), 5.33-5.20 (m, 1H), 4.40-4.23 (m, 2H), 1.40-1.30 (m, 3H). Anal. Calc'd for  $\text{C}_{12}\text{H}_{10}\text{NO}_2\text{FCl}$ : C, 47.00; H, 3.29; N, 9.13 Found: C, 46.83; H, 3.03; N, 9.18.

15 Step 5. Preparation of 6-chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylic acid.

The ester from Step 4 (1.3 g, 4.4 mmol) and 2.5 N sodium hydroxide solution (3.5 mL, 9 mmol)  
20 were mixed in tetrahydrofuran (25 mL), methanol (10 mL), and water (25 mL). The mixture was heated at 50 °C for 4 hours, allowed to cool to room temperature, and was concentrated *in vacuo* to remove the tetrahydrofuran and methanol. The  
25 resulting aqueous solution was washed with diethyl ether (2 x 100 mL). The aqueous phase was acidified with 3 N HCl causing the precipitation of a yellow solid (1.1 g). The solid was triturated with ethanol-acetone and collected by  
30 vacuum filtration providing the title compound as a yellow solid (276 mg, 23%): mp 287.4-288.4 °C.  $^1\text{H}$  NMR (acetone- $d_6$ /300 MHz) 11.50 (br s, 1H), 8.03 (s, 1H), 7.83 (s, 1H), 7.75 (s, 1H), 7.28 (br s, 1H), 5.42-5.30 (m, 1H). Anal. Calc'd for  $\text{C}_{10}\text{H}_6\text{NO}_2\text{FCl}$ : C, 43.11; H, 2.17; N, 10.05 Found:  
35 C, 42.88; H, 2.03; N, 10.06.

**EXAMPLE 175**

5

**(S)-6,8-Dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid**

6,8-Dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid (Example 32) (300 g, 1.04 mol) was added to ethyl acetate (750 mL). The mixture was stirred for 5 minutes, warmed to 70 °C and held at this temperature for 5 minutes. The resulting solution was cooled to 50 °C and (s)-(-)- $\alpha$ -methylbenzylamine (58 g, 0.48 mol) was added. Heptane (1880 mL) was added and the mixture stirred for 0.5 hour, then stirring was discontinued. The reaction was allowed to cool to 22 °C and stand for 8 hours. The salt crystallized during this time and was collected by vacuum filtration. The solid was washed with ethyl acetate-heptane (1:3, 2 X 50 mL). The solid obtained was dried at 40 °C under vacuum (20 mm) for 24 hours to give the salt (35 g, 16 %).

A three-neck 2 L round bottom flask was purged with nitrogen and was charged with deionized water (750 mL) and the salt (103 g, 0.24 mole; This material was obtained using a similar procedure to that described above). To the resulting stirred suspension was added concentrated HCl (37 mL) drop-wise over 0.5 hours with good stirring below 20 °C causing the free carboxylic acid to precipitate. After stirring

for 2 hours, the suspension was vacuum filtered and the solid washed with deionized water ( 5 X 50 mL; until the washings were neutral). The solid was dried at 40 °C under vacuum (20 mm) for 12 hours yielding the title compound as a solid (74 g, 100%): mp 166.0-168.4 °C. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>/300 MHz) 7.94 (s, 1H), 7.60 (s, 2H), 6.04 (q, 1H, *J* = 6.8 Hz). ESHRMS *m/z* 310.9489 (M-H, Calc'd 310.9450). This compound was determined to have an optical purity of greater than 90% ee. The optical purity was determined by the method described in Example 66.

#### BIOLOGICAL EVALUATION

15

##### Rat Carrageenan Foot Pad Edema Test

The carrageenan foot edema test was performed with materials, reagents and procedures essentially as described by Winter, et al., (*Proc. Soc. Exp. Biol. Med.*, 111, 544 (1962)). Male Sprague-Dawley rats were selected in each group so that the average body weight was as close as possible. Rats were fasted with free access to water for over sixteen hours prior to the test. The rats were dosed orally (1 mL) with compounds suspended in vehicle containing 0.5% methylcellulose and 0.025% surfactant, or with vehicle alone. One hour later a subplantar injection of 0.1 mL of 1% solution of carrageenan/sterile 0.9% saline was administered and the volume of the injected foot was measured with a displacement plethysmometer connected to a pressure transducer with a digital indicator. Three hours after the injection of the carrageenan, the volume of the foot was again measured. The average foot swelling in a group of drug-treated animals was compared with that of a group of placebo-treated animals and the percentage inhibition of edema was determined (Otterness and Bliven, *Laboratory Models for Testing NSAIDs*, in *Non-steroidal Anti-Inflammatory Drugs*, (J. Lombardino, ed. 1985)). The %

inhibition shows the % decrease from control paw volume determined in this procedure and the data for selected compounds in this invention are summarized in Table I.

5

TABLE I.

		RAT PAW EDEMA	ANALGESIA
		% Inhibition	% Inhibition
	<u>Example</u>	<u>@ 30mg/kg body weight</u>	<u>@ 30mg/kg body weight</u>
10	1	57	58

---

Evaluation of COX-1 and COX-2 activity *in vitro*

15        The compounds of this invention exhibited inhibition *in vitro* of COX-2. The COX-2 inhibition activity of the compounds of this invention illustrated in the Examples was determined by the following methods.

20        a. Preparation of recombinant COX baculoviruses

Recombinant COX-1 and COX-2 were prepared as described by Gierse et al, [*J. Biochem.*, 305, 479-84 (1995)]. A 2.0 kb fragment containing the coding region of either human or murine COX-1 or human or murine COX-2 was cloned into a  
25    BamHI site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the baculovirus transfer vectors for COX-1 and COX-2 in a manner similar to the method of D.R. O'Reilly et al (*Baculovirus Expression Vectors: A Laboratory Manual* (1992)). Recombinant baculoviruses were  
30    isolated by transfecting 4 µg of baculovirus transfer vector DNA into SF9 insect cells ( $2 \times 10^8$ ) along with 200 ng of linearized baculovirus plasmid DNA by the calcium phosphate method. See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*,  
35    Texas Agric. Exp. Station Bull. 1555 (1987). Recombinant viruses were purified by three rounds of plaque purification and high titer ( $10^7$ - $10^8$  pfu/mL) stocks of virus were prepared. For large scale production, SF9 insect cells were

infected in 10 liter fermentors ( $0.5 \times 10^6/\text{mL}$ ) with the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were centrifuged and the cell pellet homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS). The homogenate was centrifuged at  $10,000\times G$  for 30 minutes, and the resultant supernatant was stored at  $-80^\circ\text{C}$  before being assayed for COX activity.

10

b. Assay for COX-1 and COX-2 activity

COX activity was assayed as  $\text{PGE}_2$  formed/ $\mu\text{g}$  protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme were incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme with the addition of arachidonic acid ( $10 \mu\text{M}$ ). Compounds were pre-incubated with the enzyme for 10-20 minutes prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme was stopped after ten minutes at  $37^\circ\text{C}$ /room temperature by transferring  $40 \mu\text{l}$  of reaction mix into  $160 \mu\text{l}$  ELISA buffer and  $25 \mu\text{M}$  indomethacin. The  $\text{PGE}_2$  formed was measured by standard ELISA technology (Cayman Chemical). Results are shown in Table II.

25

c. Fast assay for COX-1 and COX-2 activity

COX activity was assayed as  $\text{PGE}_2$  formed/ $\mu\text{g}$  protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme were incubated in a potassium phosphate buffer (0.05 M Potassium phosphate, pH 7.5,  $2 \mu\text{M}$  phenol,  $1 \mu\text{M}$  heme,  $300 \mu\text{M}$  epinephrine) with the addition of  $20 \mu\text{l}$  of  $100 \mu\text{M}$  arachidonic acid ( $10 \mu\text{M}$ ). Compounds were pre-incubated with the enzyme for 10 minutes at  $25^\circ\text{C}$  prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme was

30

35



stopped after two minutes at 37 °C/room temperature by transferring 40  $\mu$ l of reaction mix into 160  $\mu$ l ELISA buffer and 25  $\mu$ M indomethacin. The PGE<sub>2</sub> formed was measured by standard ELISA technology (Cayman Chemical). Results are shown in Table II.

TABLE II.

	Example	COX-2'	COX-1'	COX-2	COX-1
		IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M
10	1	0.3	45		
	2	<0.1	78	<0.1	5.0
	6	<0.1	>100		
	7	0.1	16	<0.1	1.0
15	8	<0.1	61	<0.1	21
	9	<0.1	1.4	<0.1	<0.1
	12	7	55		
	13	.3	>100		
20	14	>100	>100		
	15	>0.1	11	133.6	44
	16	<0.1	24	1.4	51
	18	12	>100		
25	21	11	3.5		
	22	>100	>100		
	23	7	>100	24	>100
	25	>100	78		
30	26	>100	20		
	27	67	>100		
	29	<0.1	>100		
	30	<0.1	1.2	16	3.8
35	31	<0.1	94		
	32	0.3	31	0.3	0.7
	33	<0.1	5.7	8.2	28
	35	2.2	8.9	1.7	11
35	38	0.2	6.2	25.7	57
	39	0.2	45	1.3	>100
	40	<0.1	24	74	43

TABLE II. Cont.

	Exempl	COX-2'	COX-1'	COX-2	COX-1
		IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M
5	42	<0.1	2.3	<0.1	11
	43	99	85		
	44	0.3	72	21	>100
	45	0.2	47	46	>100
	46	0.2	24	74	43
10	47	1.9	31	1.7	>100
	49	24	>100	31	>100
	50	79	>100		
	52	20	>100		
	53	8	13	6	>100
15	54	19	>100		
	55	46	>100	53	>100
	56	12	>100	29	>100
	57	21	10	21	>100
	59	43	>100		
20	63	1.4	>100		
	65	<0.1	1.0		
	66	82	38	<0.1	16.9
	67	<0.1	30	<0.1	6.7
	81	<0.1	10.5	<0.1	1.6
25	82	<0.1	16	<0.1	5.6
	83	<0.1	9.6	<0.1	1.4
	84	0.1	25	<0.1	2.8
	88	<0.1	12.4	<0.1	6.4
	91	<0.1	23	0.2	36
30	96	0.2	>100	0.3	100
	97	0.2	78	0.1	25
	98	2.0	>100	1.5	19
	99	0.2	36	<0.1	23
	101	<0.1	18	<0.1	16
35	103	36	61		
	104	<0.1	24	<0.1	8.2
	105	0.3	4.5	0.2	0.1

TABLE II. Cont.

Example	COX-2		COX-1	
	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M	IC <sub>50</sub> $\mu$ M
5	106	0.2	21	<0.1
	114	<0.1	<0.1	<0.1
	115	<0.1	<0.1	<0.1
	116	<0.1	<0.1	<0.1
	120	<0.1	98	<0.1
10	125	<0.1	0.2	<0.1
	129	0.2	2.6	<0.1
	138	0.3	42.5	<0.1
	152	<0.1	74	<0.1
	154	0.5	68.5	<0.1
15	155	<0.1	1.6	<0.1
	156	<0.1	0.8	<0.1

\* fast assay

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of Formula I in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

The phrase "co-therapy" (or "combination-therapy"), in defining use of a cyclooxygenase-2 inhibitor agent and another pharmaceutical agent, is intended to embrace administration of each agent in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a substantially

simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent.

5 The phrase "therapeutically-effective" is intended to qualify the amount of each agent which will achieve the goal of improvement in disease severity and the frequency of incidence over treatment of each agent by itself, while avoiding adverse side effects typically associated with alternative therapies.

10 For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active  
15 ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier.

20 The amount of therapeutically active compounds which are administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of  
25 the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredients in the range of about 0.1 to 2000 mg, preferably in the range of about  
30 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.5 and about 20 mg/kg body weight and most preferably between about 0.1 to 10 mg/kg body weight, may be appropriate. The daily dose can be  
35 administered in one to four doses per day.

In the case of psoriasis and other skin conditions, it may be preferable to apply a topical preparation of

compounds of this invention to the affected area two to four times a day.

For inflammations of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably applied as a topical ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one

emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w.

For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the

5 compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate,

10 polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations

15 for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents

20 mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants


25 and modes of administration are well and widely known in the pharmaceutical art.

All mentioned references are incorporated by reference as if here written. The priority document, USSN 60/044,485 is also incorporated by reference.

30 Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What is claimed is:

1. A compound of Formula I'



wherein X is selected from O, S, CR<sup>a</sup>R<sup>b</sup> and NR<sup>a</sup>;  
 wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>6</sub>-alkyl,  
 (optionally substituted phenyl)-C<sub>1</sub>-C<sub>6</sub>-alkyl, acyl  
 and carboxy-C<sub>1</sub>-C<sub>6</sub>-alkyl;  
 wherein each of R<sup>b</sup> and R<sup>c</sup> is independently selected  
 from hydrido, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-  
 C<sub>6</sub>-perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-  
 alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>6</sub>-alkyl;  
 wherein R is selected from carboxyl, aminocarbonyl,  
 C<sub>1</sub>-C<sub>6</sub>-alkylsulfonylaminocarbonyl and C<sub>1</sub>-C<sub>6</sub>-  
 alkoxycarbonyl;  
 wherein R<sup>a</sup> is selected from hydrido, phenyl, thienyl  
 and C<sub>2</sub>-C<sub>6</sub>-alkenyl;  
 wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>6</sub>-perfluoroalkyl,  
 chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro,  
 cyano and cyano-C<sub>1</sub>-C<sub>6</sub>-alkyl;  
 wherein R<sup>2</sup> is one or more radicals independently  
 selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-  
 alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-  
 C<sub>1</sub>-C<sub>6</sub>-alkyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>2</sub>-C<sub>6</sub>-  
 alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-C<sub>6</sub>-  
 alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, aryloxy,  
 arylthio, arylsulfinyl, heteroaryloxy, C<sub>1</sub>-C<sub>6</sub>-  
 alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy,  
 heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, aryl-C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-  
 alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-  
 haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-



haloalkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-(haloalkyl-C<sub>1</sub>-C<sub>3</sub>-  
hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxyimino-C<sub>1</sub>-  
C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, aryl-C<sub>1</sub>-C<sub>6</sub>-  
alkylamino, heteroaryl-amino, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-  
5 alkylamino, nitro, cyano, amino, aminosulfonyl,  
C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, arylaminosulfonyl,  
heteroarylaminosulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-  
alkylaminosulfonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-  
alkylaminosulfonyl, heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-  
10 alkylsulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl,  
optionally substituted aryl, optionally  
substituted heteroaryl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,  
heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,  
heteroarylcarbonyl, arylcarbonyl, aminocarbonyl,  
15 C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, formyl, C<sub>1</sub>-C<sub>6</sub>-  
haloalkylcarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; and  
wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are  
independently selected from carbon and nitrogen  
with the proviso that at least two of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup>  
20 and A<sup>4</sup> are carbon;  
or wherein R<sup>2</sup> together with ring A forms a radical  
selected from naphthyl, quinolyl, isoquinolyl,  
quinoliziny, quinoxaliny and dibenzofuryl;  
or an isomer or pharmaceutically acceptable salt  
25 thereof.

2. A compound of Claim 1 wherein X is selected  
from O, S, CR<sup>a</sup>R<sup>b</sup> and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from  
hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, (optionally substituted  
30 phenyl)-C<sub>1</sub>-C<sub>3</sub>-alkyl, acyl and carboxy-C<sub>1</sub>-C<sub>6</sub>-alkyl;  
wherein each of R<sup>b</sup> and R<sup>b</sup> is independently selected  
from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-  
perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy,  
nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R is  
35 selected from carboxyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-  
alkylsulfonylaminocarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl;  
wherein R<sup>a</sup> is selected from hydrido, phenyl, thienyl  
and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-

perfluoroalkyl, chloro, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkoxy, nitro, cyano and cyano-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>1</sub>-C<sub>3</sub>-alkyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, aryl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, C<sub>1</sub>-C<sub>6</sub>-alkylsulfinyl, aryloxy, arylthio, arylsulfinyl, heteroaryloxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, aryl-C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-haloalkylthio, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>3</sub>-(haloalkyl-C<sub>1</sub>-C<sub>3</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, hydroxyimino-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, heteroarylamino, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, cyano, amino, aminosulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylaminosulfonyl, heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted aryl, optionally substituted heteroaryl, aryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, formyl, C<sub>1</sub>-C<sub>6</sub>-haloalkylcarbonyl and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; and wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

3. A compound of Claim 2 wherein X is selected from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl and (optionally substituted phenyl)methyl; wherein R is carboxyl; wherein R<sup>b</sup> is selected from hydrido and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R<sup>1</sup> is

selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, 5 phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy, methylenedioxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkylthio, C<sub>1</sub>-C<sub>3</sub>-alkylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, C<sub>1</sub>-C<sub>3</sub>-haloalkyl-C<sub>1</sub>-C<sub>3</sub>-hydroxyalkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyloxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-10 haloalkyl, C<sub>1</sub>-C<sub>3</sub>-haloalkoxy, C<sub>1</sub>-C<sub>3</sub>-haloalkylthio, C<sub>1</sub>-C<sub>3</sub>-hydroxyalkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, hydroxyimino-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, cyano, amino, aminosulfonyl, N-alkylaminosulfonyl, N-arylamino-15 sulfonyl, N-heteroarylamino sulfonyl, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkylsulfonyl, 5- to 8-membered heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- to 9-membered heteroaryl, 20 phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, phenylcarbonyl, 4-chlorophenylcarbonyl, 4-hydroxyphenylcarbonyl, 4-trifluoromethylphenylcarbonyl, 4-methoxyphenylcarbonyl, aminocarbonyl, formyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and 25 nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl, benzofurylphenyl, or quinolyl radical; or an isomer or pharmaceutically 30 acceptable salt thereof.

4. A compound of Claim 3 wherein X is selected from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, 35 (4-chloromethyl)benzyl, (4-methoxy)benzyl, and (4-cyano)benzyl, (4-nitro)benzyl; wherein R is carboxyl; wherein R<sup>a</sup> is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl

and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenylethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, optionally substituted phenylcarbonyl, aminocarbonyl, formyl and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl, or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

5. A compound of Claim 4 selected from compounds, and their isomers and pharmaceutically-acceptable salts, of the group consisting of 6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 7-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-ethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 35 7-isopropoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6,8-bis(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dimethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-nitro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- ethyl 6-amino-2-trifluoromethyl-2H-1-benzopyran-3-carboxylate;
- 35 6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-difluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-(N,N-diethylamino)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(phenylmethyl)amino]sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6-[(dimethylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-aminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6-(methylamino)sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(4-morpholino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(1,1-dimethylethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-[(2-methylpropyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 35 8-chloro-6-[(phenylmethyl)amino]sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 6-N,N-diethylaminosulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-(2,2-dimethylpropylcarbonyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-7-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 10 6-[[ (2-furanylmethyl) amino] sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-[(phenylmethyl) sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-[[ (phenylethyl) amino] sulfonyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6-chloro-8-iodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-formyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 5,6-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid ;
- 35 6-hydroxymethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;



- 2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5,6,7-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 6,7,8-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(methylsulfinyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 5,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-chloro-2,7-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5-methoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-benzoyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-chlorobenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-(4-chlorophenoxy)-2-trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 35 2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-(3-chloro-4-methoxyphenoxy)-2-(trifluoromethyl)-  
2H-1-benzopyran-3-carboxylic acid;
- 6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 5 8-chloro-2-(trifluoromethyl)-6-[4-  
(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-  
carboxylic acid;
- 6-chloro-8-cyano-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 10 6-chloro-8-[(hydroxyimino)methyl]-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic  
acid;
- 6-chloro-8-(hydroxymethyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 15 8-(1H-benzimidazol-2-yl)-6-chloro-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic  
acid;
- 7-(1,1-dimethylethyl)-2-(pentafluoroethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 20 6-chloro-8-(methoxymethyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 6-chloro-8-(benzyloxymethyl)-2-(trifluoromethyl)-2H-  
1-benzopyran-3-carboxylic acid;
- 6-chloro-8-ethenyl-2-(trifluoromethyl)-2H-1-  
25 benzopyran-3-carboxylic acid;
- 6-chloro-8-ethynyl-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 6-chloro-8-(2-thienyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 30 6-chloro-8-(2-furanyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 6-chloro-8-(5-chloro-1-pentynyl)-2-  
(trifluoromethyl)-2H-1-benzopyran-3-carboxylic  
acid;
- 35 6-chloro-8-(1-pentynyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;
- 6-chloro-8-(phenylethynyl)-2-(trifluoromethyl)-2H-1-  
benzopyran-3-carboxylic acid;

- 6-chloro-8-(3,3-dimethyl-1-butynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-[(4-chlorophenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(4-methoxyphenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(3-methoxyphenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-8-[(4-methylthio)phenyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(4-methylsulfonyl)phenyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-(4-fluorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 30 8-chloro-6-fluoro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6,8-diiodo-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(5-chloro-2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 35 6-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-bromophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 6-(ethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(4-methoxyphenyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-2-(trifluoromethyl)-4-ethenyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-2-(trifluoromethyl)-4-phenyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-4-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(2,2,2-trifluoro-1-hydroxyethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,8-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 25 6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,7-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 30 8-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 35 6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 7-chloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;

- 6,7-dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;  
2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid;  
5 6,8-dichloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;  
6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6,8-dichloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
10 6,7-difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
15 6-bromo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-(trifluoromethyl)-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
20 6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
25 6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(trifluoromethyl)phenyl]methyl]-3-quinolinecarboxylic acid;  
6-chloro-1-[(4-chlorophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
30 6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(methoxy)phenyl]methyl]-3-quinolinecarboxylic acid;  
6-chloro-1-[(4-cyanophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
35 6-chloro-1,2-dihydro-1-[(4-nitrophenyl)methyl]-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-1-ethyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;

- 6-chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylic acid;  
 2-trifluoromethyl-2H-naphtho[1,2-b]pyran-3-carboxylic acid;  
 5 2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;  
 2-trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid;  
 5-(hydroxymethyl)-8-methyl-2-(trifluoromethyl)-2H-pyrano[2,3-c]pyridine-3-carboxylic acid;  
 10 6-(trifluoromethyl)-6h-1,3-dioxolo[4,5-g][1]benzopyran-7-carboxylic acid; and  
 3-(trifluoromethyl)-3H-benzofuro[3,2-f][1]benzopyran-2-carboxylic acid.

15

6. A compound of Claim 2 wherein X is O; wherein R is carboxyl; wherein R' is selected from hydrido and C<sub>2</sub>-C<sub>6</sub>-alkenyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more  
 20 radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy, 5- or 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5- or 6-membered heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-  
 25 haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-heteroaryl-amino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, nitro, amino, aminosulfonyl, N-(C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N,N-di-(C<sub>1</sub>-C<sub>6</sub>-  
 30 alkyl)aminosulfonyl, N-arylaminosulfonyl, N-heteroarylaminosulfonyl, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally  
 35 substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl,

and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or  
5 pharmaceutically acceptable salt thereof.

7. A compound of Claim 6 wherein X is O;  
wherein R is carboxyl; wherein R" is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from  
10 trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl,  
15 phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl,  
20 trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino,  
25 aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl  
30 substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl,  
35 phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; and wherein one of the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> is nitrogen and the other three are

carbon; or an isomer or pharmaceutically acceptable salt thereof.

9. A compound of Claim 7 wherein X is O;
- 5 wherein R is carboxyl; wherein R" is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently selected from
- 10 hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy, methylthio, methylsulfinyl, phenyloxy, phenylthio,
- 15 phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl, hydroxy-trifluoroethyl,
- 20 methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl,
- 25 benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl
- 30 substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically
- 35 acceptable salt thereof.



10. A compound of Claim 9 selected from compounds, and their isomers and pharmaceutically-acceptable salts, of the group consisting of
- 6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 (S)-6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 (S)-6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- (S)-6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-7-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- (S)-6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 35 6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- (S)-6-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 6-hydroxymethyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(difluoromethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5 2,6-bis(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 5,6,7-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6,7,8-trichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-(methylthio)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(pentafluoroethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-7-methyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-benzoyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-(4-chlorobenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(4-hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-phenoxy-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- (S)-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- 30 6-(4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(3-chloro-4-methoxyphenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 35 6-(4-chlorophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;

- 8-chloro-2-(trifluoromethyl)-6-[4-(trifluoromethyl)phenoxy]-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-8-cyano-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(2-thienyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 10 6-chloro-8-[(4-chlorophenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-[(4-methoxyphenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 15 (S)-6-chloro-8-[(4-methoxyphenyl)ethynyl]-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-(phenylethynyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 20 6-chloro-8-(4-chlorophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-phenyl-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 25 6-(4-bromophenyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(4-methoxyphenyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid; and
- 6-(2,2,2-trifluoro-1-hydroxyethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid.
- 30

11. A compound of Claim 2 wherein X is S; wherein R is carboxyl; wherein R<sup>1</sup> is selected from
- 35 C<sub>1</sub>-C<sub>6</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy, 5- or

- 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5-  
or 6-membered heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-  
haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, N-  
phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-  
5 heteroarylamino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino,  
nitro, amino, aminosulfonyl, N-alkylaminosulfonyl,  
N-arylaminosulfonyl, N-heteroarylaminosulfonyl, N-  
(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-  
C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered  
10 heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally  
substituted phenyl, optionally substituted 5- or 6-  
membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl,  
heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl,  
and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>,  
15 A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon  
and nitrogen with the proviso that at least three of  
A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or  
pharmaceutically acceptable salt thereof.
- 20 12. A compound of Claim 11 wherein X is S;  
wherein R is carboxyl; wherein R" is selected from  
hydrido and ethenyl; wherein R<sup>1</sup> is selected from  
trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is  
one or more radicals independently selected from  
25 hydrido, chloro, bromo, fluoro, iodo, methyl, tert-  
butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-  
pentynyl, 3,3-dimethyl-1-butynyl, benzyl,  
phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl,  
4-methoxyphenyl-ethynyl, phenylethenyl, methoxy,  
30 methylthio, methylsulfinyl, phenyloxy, phenylthio,  
phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy,  
phenylmethoxy, methylenedioxy, benzyloxymethyl,  
trifluoromethyl, difluoromethyl, pentafluoroethyl,  
trifluoromethoxy, trifluoromethylthio,  
35 hydroxymethyl, hydroxy-trifluoroethyl,  
methoxymethyl, hydroxyiminomethyl, N-methylamino, N-  
phenylamino, N-(benzyl)amino, nitro, cyano, amino,  
aminosulfonyl, N-methylaminosulfonyl, N-

- phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-(furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl
- 5 substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl, benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl,
- 10 phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.
- 15 13. A compound of Claim 12 selected from compounds, and their isomers and pharmaceutically-acceptable salts, of the group consisting of
- 20 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;
- 6-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,8-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 25 6-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 6,7-dimethyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 30 8-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 35 6-chloro-7-methyl-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;
- 7-chloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;

- 6,7-dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid;  
2-(trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzopyran-3-carboxylic acid; and  
5 6,8-dichloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid.

14. A compound of Claim 2 wherein X is NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, acyl and carboxy-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R is carboxyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-perfluoroalkyl; wherein R<sup>2</sup> is one or more radicals independently selected from hydrido, halo, C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkynyl, phenyl-C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy, phenyloxy, 5- or 6-membered heteroaryloxy, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, 5- or 6-membered heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylamino, N-phenylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)amino, N-heteroarylamino, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkylamino, nitro, amino, aminosulfonyl, N-alkylaminosulfonyl, N-arylamino, N-heteroarylamino, N-(phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, N-(heteroaryl-C<sub>1</sub>-C<sub>6</sub>-alkyl)aminosulfonyl, 5- to 8-membered heterocyclylsulfonyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl, optionally substituted phenyl, optionally substituted 5- or 6-membered heteroaryl, phenyl-C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, heteroarylcarbonyl, phenylcarbonyl, aminocarbonyl, and C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an isomer or pharmaceutically acceptable salt thereof.

- 35 15. A compound of Claim 14 wherein X is NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, (4-chloromethyl)benzyl, (4-methoxy)benzyl, (4-cyano)benzyl, and (4-

- nitro)benzyl; wherein R is carboxyl; wherein R" is selected from hydrido and ethenyl; wherein R' is selected from trifluoromethyl and pentafluoroethyl; wherein R<sup>2</sup> is one or more radicals independently
- 5 selected from hydrido, chloro, bromo, fluoro, iodo, methyl, tert-butyl, ethenyl, ethynyl, 5-chloro-1-pentynyl, 1-pentynyl, 3,3-dimethyl-1-butynyl, benzyl, phenylethyl, phenyl-ethynyl, 4-chlorophenyl-ethynyl, 4-methoxyphenyl-ethynyl, phenylethenyl, methoxy,
- 10 methylthio, methylsulfinyl, phenyloxy, phenylthio, phenylsulfinyl, pyridyloxy, thienyloxy, furyloxy, phenylmethoxy, methylenedioxy, benzyloxymethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, trifluoromethoxy, trifluoromethylthio, hydroxymethyl,
- 15 hydroxy-trifluoroethyl, methoxymethyl, hydroxyiminomethyl, N-methylamino, N-phenylamino, N-(benzyl)amino, nitro, cyano, amino, aminosulfonyl, N-methylaminosulfonyl, N-phenylaminosulfonyl, N-furylaminosulfonyl, N-(benzyl)aminosulfonyl, N-
- 20 (furylmethyl)aminosulfonyl, benzylsulfonyl, phenylethylaminosulfonyl, furylsulfonyl, methylsulfonyl, phenyl, phenyl substituted with one or more radicals selected from chloro, fluoro, bromo, methoxy, methylthio and methylsulfonyl,
- 25 benzimidazolyl, thienyl, thienyl substituted with chloro, furyl, furyl substituted with chloro, benzylcarbonyl, furylcarbonyl, phenylcarbonyl, aminocarbonyl, formyl, and methylcarbonyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or an
- 30 isomer or pharmaceutically acceptable salt thereof.

16. A compound of Claim 15 selected from compounds, and their isomers and pharmaceutically-acceptable salts, of the group consisting of
- 35 6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;

- 6,8-dichloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6,7-difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
5 6-iodo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-bromo-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
1,2-dihydro-6-(trifluoromethoxy)-2-(trifluoromethyl)-  
10 3-quinolinecarboxylic acid;  
6-(trifluoromethyl)-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-cyano-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
15 6-chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(trifluoromethyl)phenyl]methyl]-3-quinolinecarboxylic acid;  
20 6-chloro-1-[(4-chlorophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-2-(trifluoromethyl)-1-[[4-(methoxy)phenyl]methyl]-3-quinolinecarboxylic acid;  
25 6-chloro-1-[(4-cyanophenyl)methyl]-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-1-[(4-nitrophenyl)methyl]-2-(trifluoromethyl)-3-quinolinecarboxylic acid;  
6-chloro-1,2-dihydro-1-ethyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid; and  
30 (S)-6-chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid.

17. A compound of Claim 2 wherein X is selected  
35 from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from hydrido, C<sub>1</sub>-C<sub>3</sub>-alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, acyl and carboxy-C<sub>1</sub>-C<sub>3</sub>-alkyl; wherein R is selected from carboxyl; wherein R<sup>1</sup> is selected from C<sub>1</sub>-C<sub>3</sub>-




perfluoroalkyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; and wherein R<sup>2</sup> together with ring A  
5 forms a naphthyl or quinolyl radical; or an isomer or pharmaceutically acceptable salt thereof.

18. A compound of Claim 17 wherein X is selected from O, S and NR<sup>a</sup>; wherein R<sup>a</sup> is selected from  
10 hydrido, methyl, ethyl, (4-trifluoromethyl)benzyl, (4-chloromethyl)benzyl, (4-methoxy)benzyl, and (4-cyano)benzyl, (4-nitro)benzyl; wherein R is carboxyl; wherein R<sup>"</sup> is selected from hydrido and ethenyl; wherein R<sup>1</sup> is selected from trifluoromethyl and  
15 pentafluoroethyl; wherein the A ring atoms A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are independently selected from carbon and nitrogen with the proviso that at least three of A<sup>1</sup>, A<sup>2</sup>, A<sup>3</sup> and A<sup>4</sup> are carbon; or wherein R<sup>2</sup> together with ring A forms a naphthyl, or quinolyl radical; or an  
20 isomer or pharmaceutically acceptable salt thereof.

19. A compound of Claim 18 selected from compounds, and their isomers and pharmaceutically-acceptable salts, of the group consisting of  
25 2-trifluoromethyl-2H-naphtho[1,2-b]pyran-3-carboxylic acid;  
2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;  
30 2-trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid;  
5-(hydroxymethyl)-8-methyl-2-(trifluoromethyl)-2H-pyrano[2,3-c]pyridine-3-carboxylic acid;  
6-(trifluoromethyl)-6h-1,3-dioxolo[4,5-  
35 g][1]benzopyran-7-carboxylic acid; and  
3-(trifluoromethyl)-3H-benzofuro[3,2-f][1]benzopyran-2-carboxylic acid.

20. A compound of Formula I



5

wherein X is selected from O or S or NR<sup>a</sup>;

wherein R<sup>a</sup> is alkyl;

wherein R is selected from carboxyl,  
aminocarbonyl, alkylsulfonylaminocarbonyl and  
10 alkoxy carbonyl;

wherein R<sup>1</sup> is selected from haloalkyl, alkyl,  
aralkyl, cycloalkyl and aryl optionally substituted  
with one or more radicals selected from alkylthio,  
nitro and alkylsulfonyl; and

15 wherein R<sup>2</sup> is one or more radicals selected from  
hydrido, halo, alkyl, aralkyl, alkoxy, aryloxy,  
heteroaryloxy, aralkyloxy, heteroaralkyloxy,  
haloalkyl, haloalkoxy, alkylamino, arylamino,  
aralkylamino, heteroaryl amino, heteroarylalkyl amino,  
20 nitro, amino, aminosulfonyl, alkylaminosulfonyl,  
arylamino sulfonyl, heteroarylaminosulfonyl,  
aralkylaminosulfonyl, heteroaralkylaminosulfonyl,  
heterocyclosulfonyl, alkylsulfonyl, optionally  
substituted aryl, optionally substituted heteroaryl,  
25 aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl,  
aminocarbonyl, and alkylcarbonyl;

or wherein R<sup>2</sup> together with ring A forms a  
naphthyl radical;

30 or an isomer or pharmaceutically acceptable salt  
thereof.

21. Compound of Claim 20 wherein X is oxygen or  
sulfur; wherein R is selected from carboxyl, lower  
alkyl, lower aralkyl and lower alkoxy carbonyl;

wherein R<sup>1</sup> is selected from lower haloalkyl, lower cycloalkyl and phenyl; and wherein R<sup>2</sup> is one or more radicals selected from hydrido, halo, lower alkyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, lower alkylamino, nitro, amino, aminosulfonyl, lower alkylaminosulfonyl, 5- or 6- membered heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl, 5- or 6- membered nitrogen containing heterocyclosulfonyl, lower alkylsulfonyl, optionally substituted phenyl, lower aralkylcarbonyl, and lower alkylcarbonyl; or wherein R<sup>2</sup> together with ring A forms a naphthyl radical; or an isomer or pharmaceutically acceptable salt thereof.

22. Compound of Claim 21 wherein X is oxygen or sulfur; wherein R is carboxyl; wherein R<sup>1</sup> is lower haloalkyl; and wherein R<sup>2</sup> is one or more radicals selected from hydrido, halo, lower alkyl, lower haloalkyl, lower haloalkoxy, lower alkylamino, amino, aminosulfonyl, lower alkylaminosulfonyl, 5- or 6- membered heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl, lower alkylsulfonyl, 6- membered nitrogen containing heterocyclosulfonyl, optionally substituted phenyl, lower aralkylcarbonyl, and lower alkylcarbonyl; or wherein R<sup>2</sup> together with ring A forms a naphthyl radical; or an isomer or pharmaceutically acceptable salt thereof.

23. Compound of Claim 22 wherein R is carboxyl; wherein R<sup>1</sup> is selected from fluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, difluoromethyl, and trifluoromethyl; and wherein R<sup>2</sup> is one or more radicals selected from hydrido, chloro, fluoro, bromo, iodo, methyl, ethyl, isopropyl, tert-butyl, butyl, isobutyl, pentyl, hexyl, methoxy, ethoxy, isopropoxy, tertbutoxy, trifluoromethyl,

5 difluoromethyl, trifluoromethoxy, amino, N,N-dimethylamino, N,N-diethylamino, N-phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-furylmethyl)aminosulfonyl, nitro, N,N-dimethylaminosulfonyl, aminosulfonyl, N-methylaminosulfonyl, N-ethylsulfonyl, 2,2-dimethylethylaminosulfonyl, N,N-dimethylaminosulfonyl, N-(2-methylpropyl)aminosulfonyl, N-morpholinosulfonyl, methylsulfonyl, benzylcarbonyl, 2,2-dimethylpropylcarbonyl, phenylacetyl and phenyl; or  
10 wherein R<sup>2</sup> together with ring A forms a naphthyl radical; or an isomer or pharmaceutically acceptable  
15 salt thereof.

24. Compound of Claim 23 wherein R is carboxyl; wherein R<sup>1</sup> is trifluoromethyl or pentafluorethyl; and wherein R<sup>2</sup> is one or more radicals selected from  
20 hydrido, chloro, fluoro, bromo, iodo, methyl, ethyl, isopropyl, tert-butyl, methoxy, trifluoromethyl, trifluoromethoxy, N-phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-furylmethyl)aminosulfonyl, N,N-dimethylaminosulfonyl,  
25 N-methylaminosulfonyl, N-(2,2-dimethylethyl)aminosulfonyl, dimethylaminosulfonyl, 2-methylpropylaminosulfonyl, N-morpholinosulfonyl, methylsulfonyl, benzylcarbonyl, and phenyl; or  
wherein R<sup>2</sup> together with ring A forms a naphthyl  
30 radical; or an isomer or pharmaceutically acceptable salt thereof.


25. A compound of Claim 24 selected from compounds, and their isomers and pharmaceutically-  
35 acceptable salts, of the group consisting of 6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 2-trifluoromethyl-3H-naphthopyran-3-carboxylic acid ;
- 10 7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-bis(dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 2-trifluoromethyl-3H-naptho[2,1-b]pyran-3-carboxylic acid;
- 6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 30 6-[[ (phenylmethyl) amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(dimethylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;

- 6-[(methylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(4-morpholino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 5 6-[(1,1-dimethylethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[(2-methylpropyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 10 8-chloro-6-[(phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 15 6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 8-chloro-5,6-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6,8-dichloro-(S)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 20 6-benzylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-[[N-(2-furylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 25 6-[[N-(2-phenylethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;
- 7-(1,1-dimethylethyl)-2-pentafluoroethyl-2H-1-benzopyran-3-carboxylic acid; and
- 30 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid.

26. A compound of Formula II



wherein X is O or S;

wherein R<sup>2</sup> is lower haloalkyl;

5        wherein R<sup>3</sup> is selected from hydrido, and halo;

      wherein R<sup>4</sup> is selected from hydrido, halo, lower  
alkyl, lower haloalkoxy, lower alkoxy, lower  
aralkylcarbonyl, lower dialkylaminosulfonyl, lower  
alkylaminosulfonyl, lower aralkylaminosulfonyl, lower  
10 heteroaralkylaminosulfonyl, and 5- or 6- membered  
nitrogen-containing heterocyclosulfonyl;


      wherein R<sup>5</sup> is selected from hydrido, lower alkyl,  
halo, lower alkoxy, and aryl; and

      wherein R<sup>6</sup> is selected from hydrido, halo, lower  
15 alkyl, lower alkoxy, and aryl;

      or an isomer or pharmaceutically acceptable salt  
thereof.

27. A compound of Formula IIa:

20



wherein R<sup>3</sup> is selected from hydrido, lower alkyl,  
lower hydroxyalkyl, lower alkoxy and halo;




wherein R<sup>4</sup> is selected from hydrido, halo, lower alkyl, lower alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydrooxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen containing heterocyclosulfonyl; wherein R<sup>5</sup> is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and wherein R<sup>6</sup> is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl, phenylalkynyl, lower alkyl, lower alkoxy, formyl and phenyl; or an isomer or pharmaceutically acceptable salt thereof.

28. Compound of Claim 27 wherein R<sup>3</sup> is selected from hydrido, and chloro; wherein R<sup>4</sup> is selected from chloro, methyl, tert-butyl, methylthio, trifluoromethyl, difluoromethyl, pentafluoromethyl, trifluoromethylsulfide, trifluoromethoxy, cyano, substituted or unsubstituted phenylcarbonyl, and substituted or unsubstituted phenyl; wherein R<sup>5</sup> is selected from hydrido, methyl, tert-butyl, chloro; and wherein R<sup>6</sup> is selected from hydrido, chloro, thienyl, hydroxyiminomethyl, substituted or unsubstituted phenylethynyl, and substituted or unsubstituted phenyl; or an isomer or pharmaceutically acceptable salt thereof.

35

29. A compound of Formula IIb:




wherein R<sup>3</sup> is selected from hydrido, lower alkyl, lower hydroxyalkyl, lower alkoxy and halo; wherein R<sup>4</sup> is selected from hydrido, halo, lower alkyl, lower alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydrooxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen containing heterocyclosulfonyl; wherein R<sup>5</sup> is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and wherein R<sup>6</sup> is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl, phenylalkynyl, lower alkyl, lower alkoxy, formyl and phenyl; or an isomer or pharmaceutically acceptable salt thereof.

30. Compound of Claim 29 wherein R<sup>3</sup> is selected from hydrido, and chloro; wherein R<sup>4</sup> is selected from chloro, methyl, tert-butyl, methylthio, trifluoromethyl, difluoromethyl, pentafluoromethyl, trifluoromethylsulfide, trifluoromethoxy, cyano, substituted or unsubstituted phenylcarbonyl, and substituted or unsubstituted phenyl; wherein R<sup>5</sup> is selected from hydrido, methyl, tert-butyl, chloro;

and wherein R<sup>6</sup> is selected from hydrido, chloro, thienyl, hydroxyiminomethyl, substituted or unsubstituted phenylethynyl, and substituted or unsubstituted phenyl; or an isomer or  
 5 pharmaceutically acceptable salt thereof.

31. A compound of Formula IIc:



10

wherein R<sup>a</sup> is selected from hydrido and lower aralkyl; wherein R<sup>3</sup> is selected from hydrido, lower alkyl, lower hydroxyalkyl, lower alkoxy and halo; wherein R<sup>4</sup> is selected from hydrido, halo, lower alkyl, lower  
 15 alkylthio, lower haloalkyl, amino, aminosulfonyl, lower alkylsulfonyl, lower alkylsulfinyl, lower alkoxyalkyl, lower alkylcarbonyl, formyl, cyano, lower haloalkylthio, substituted or unsubstituted phenylcarbonyl, lower haloalkoxy, lower alkoxy, lower  
 20 aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5- or 6- membered heteroaryl, lower hydroxyalkyl, optionally substituted phenyl and 5- or 6- membered nitrogen  
 25 containing heterocyclosulfonyl; wherein R<sup>5</sup> is selected from hydrido, lower alkyl, halo, lower haloalkyl, lower alkoxy, and phenyl; and wherein R<sup>6</sup> is selected from hydrido, halo, cyano, hydroxyiminomethyl, lower hydroxyalkyl, lower alkynyl, phenylalkynyl, lower  
 30 alkyl, lower alkoxy, formyl and phenyl;

or an isomer or pharmaceutically acceptable salt thereof.

32. Compound of Claim 31 wherein R<sup>a</sup> is hydrido;  
5 wherein R<sup>b</sup> is selected from hydrido, and chloro;  
wherein R<sup>c</sup> is selected from chloro, methyl, tert-  
butyl, methylthio, trifluoromethyl, difluoromethyl,  
pentafluoromethyl, trifluoromethylsulfide,  
trifluoromethoxy, cyano, substituted or  
10 unsubstituted phenylcarbonyl, and substituted or  
unsubstituted phenyl; wherein R<sup>d</sup> is selected from  
hydrido, methyl, tert-butyl, chloro; and wherein R<sup>e</sup> is  
selected from hydrido, chloro, thienyl,  
hydroxyiminomethyl, substituted or unsubstituted  
15 phenylethynyl, and substituted or unsubstituted  
phenyl; or an isomer or pharmaceutically acceptable  
salt thereof.

33. A method of treating a cyclooxygenase-2  
20 mediated disorder in a subject, said method  
comprising treating the subject having or susceptible  
to said disorder with a therapeutically-effective  
amount of a compound of Claims 1-31; or a  
pharmaceutically-acceptable salt thereof.

25

34. The method of Claim 33 wherein the  
cyclooxygenase-2 mediated disorder is inflammation.

35. The method of Claim 33 wherein the  
30 cyclooxygenase-2 mediated disorder is arthritis.

36. The method of Claim 33 wherein the  
cyclooxygenase-2 mediated disorder is pain.

35 37. The method of Claim 33 wherein the  
cyclooxygenase-2 mediated disorder is fever.

38. A pharmaceutical composition comprising a  
therapeutically-effective amount of a compound, said  
compound selected from a family of compounds of  
Claims 1-31; or a pharmaceutically-acceptable salt  
5 thereof.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/07677

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D311/58 C07D335/06 C07D215/54 C07D311/92 C07D493/04  
C07D409/04 C07D491/04 A61K31/35

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. BUNTING ET AL.: "KINETIC A. THERMODYNAMIC CONTROL OF PSEUDOBASE FORMAT. FROM C-3 SUBSTIT. 1-METHYLQUINOLINIUM CATIONS." CANADIAN JOURNAL OF CHEMISTRY., vol. 62, no. 7, 1984, OTTAWA CA, pages 1301-1307, XP002071264 see page 1301 - page 1305; example 6 ---	1, 14, 15, 20
A	EP 0 412 939 A (CIBA-GEIGY) 13 February 1991 cited in the application see claims --- -/--	1-31, 38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 July 1998

Date of mailing of the international search report

30.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07677

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LOIC RENE, R. ROYER: "SUR LA SYNTHÈSE DE DELTA-3-CHROMÈNES SUBSTITUÉS"  EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY. CHIMICA THERAPEUTICA.,  vol. 10, no. 1, January 1975, PARIS FR,  pages 72-78, XP002071265  cited in the application  see page 72 - page 74; table 1</p>	1-31, 38
A	<p>CHEMICAL ABSTRACTS, vol. 125, no. 25, 1996  Columbus, Ohio, US;  abstract no. 328458j,  page 1371; column 1;  XP002071266  see abstract  &amp; UKHIN, L. ET AL.: "FORMATION OF BENZOFURAN AND 2H-CHROMÈNES"  IZV. AKAD. NAUK. SER. KHIM.,  vol. 5, 1996, RUSSIA,  pages 1222-1228,</p>	1

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/07677

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 33-37  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 33-37  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/07677

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 412939 A	13-02-1991	AU 631913 B	10-12-1992
		AU 6089590 A	14-02-1991
		CA 2023016 A	12-02-1991
		JP 3083979 A	09-04-1991
		MX 21917 A	28-02-1994
		PT 94912 A	18-04-1991
		US 5155130 A	13-10-1992